Vol. 93, Sect. 9, 1956

## **MOLLUSCA**

COMPILED BY

A. E. SALISBURY, A.M.I.E.E.

ASSISTED BY

MARCIA A. EDWARDS, B.Sc.

LONDON
PUBLISHED BY
THE ZOOLOGICAL SOCIETY OF LONDON

PRICE TEN SHILLINGS

JANUARY, 1959

# SEPARATE SECTIONS OF THE ZOOLOGICAL RECORD

In addition to the complete bound volume, the separate Sections may be obtained singly, bound in printed paper covers.

The following are the Sections and their prices, viz.:-

								8.	d.	
1.	Comprehens	sive Zo	ology		•••		•••	3	0	
2.	Protozoa							10	0	
3.	Porifera							2	0	
4.	Coelenterate	a.						4	0	
5.	Echinodern	ıa						4	0	
6.	Vermes							10	0	
7.	Brachiopod	a						4	0	
8.	Bryozoa			***				2	6	
9.	Mollusca							10	0	
10.	Crustacea							6	6	
11.	Trilobita							5	0	
12.	Arachnida							10	0	
*13.	Insecta							50	0	
14.	Protochorda	ata						2	0	
	Pisces							8	6	
16A	. Amphibia							6	0	
	TD CTT							17	6	
	Aves							10	0	
	Mammalia				***	•••		10	0	
				•••	***	***				
	List of New	Gener	a and a	Subgen	era	***	•••	4	0	

<sup>\*</sup> Obtainable only from the Commonwealth Institute of Entomology, 56, Queen's Gate, London, S.W.7.

## 9. MOLLUSCA

COMPILED BY

### A. E. SALISBURY, A.M.I.E.E.

ASSISTED BY

## Marcia A. EDWARDS, B.Sc.

### CONTENTS

							PAGE
I.	TITLES	٠.			• •	 ١	2
II.	SUBJECT INDEX			• •		 	65
III.	Systematic Index					 	95
	Polyplacophora					 	95
	Gastropoda					 	96
	Prosobranchiata					 	96
	Docoglossa					 	96
	Opisthobranchiata	à.				 	111
	Nudibranchiata					 	112
	Pulmonata		**			 	113
	Scaphopoda					 	122
	Pelecypoda or Lai	melli	ibranchi	a		 	122
	Cephalopoda		2			 	138

#### I.—TITLES

The year of publication has been omitted where it is the same as the volume year of the 'Record,' namely (1956).

Opinion 386. Validation under the plenary powers of the generic name *Pyramidella* Lamarck, 1799. (Class Gastropoda). Opin. int. Comm. zool. Nom. 12: 231–240.

Opinion 391. Emendation to Muensteroceras of the generic name Munsteroceras Hyatt, 1884. (Class Cephalopoda, Order Ammonoidea). Opin. int. Comm. zool. Nom. 12: 285–294.

Opinion 392. Emendation to Sandbergeroceras of the generic name Sandbergeoceras Hyatt 1844. Opin. int. Comm. zool. Nom. 12: 295–304.

Opinion 420. Addition to the Official List of Specific Names in Zoology of the specific names for eleven species of the Class Brachiopoda and for two species of the Class Cephalopoda originally published by Martin (W.) in 1809 in the nomenclatorially invalid work entitled Petrificata Derbiensia and now available as from the first subsequent date on which they were severally published in conditions satisfying the requirements of the Règles. Opin. int. Comm. zool. Nom. 14: 129–168.

Opinion 422. Determination under the Plenary Powers of the species to which the specific name mammillatus Schlotheim, 1813, as published in the combination Ammonites mammillatus shall apply and designation under the same powers of the foregoing species to be the type species of the genus Douvilleiceras de Grossouvre, 1893 (Class Cephalopoda, Order Ammonoidea). Opin. int. Comm. zool. Nom. 14:181–196.

Opinion 426. Designation under the Plenary Powers of type species in harmony with accustomed usage for the genera *Pictonia* Bayle, 1878, and *Rasenia* Salfeld, 1913 (Class Cephalopoda, Order Ammonoidea) (Jurassic). Opin. int. Comm. zool. Nom. 14: 257–280.

Opinion 427. Rejection for nomenclatorial purposes of the work by Renier (S.A.) known as Tavole per servire alle classificazione e connescenza degli animali and commonly attributed to the year 1807 and addition to the Official Indexes of Rejected and Invalid Names in Zoology of certain names first used in the foregoing work or in two earlier works by the same author commonly known as Tavola alfabetica and the Prospetto respectively and both commonly attributed to the year 1804. Opin. int. Comm. zool. Nom. 14: 281-310.

Opinion 428. Supression under the Plenary Powers of the specific name royerianus d'Orbigny, 1841, as published in the combination Ammonites royerianus and designation under the same Powers of a type species in harmony with the accustomed practice for the genus Cheloniceras, Hyatt, 1903 (Class Cephalopoda, Order Ammonoidea). Opin. int. Comm. zool. Nom. 14: 311-322.

Opinion 429. Direction under the Plenary Powers limiting to suppression for the purposes of the Law of Priority the suppression of the generic name Argus Bohadsch, 1761 (Class Gastropoda) prescribed by the Ruling given in Opinion 185 thereby securing that the generic name Argus Scopoli, 1763 (Class Insecta, Order Lepidoptera) shall remain invalid under the Law of Homonymy. Opin. int. Comm. zool., Nom. 14: 323–337.

Opinion 430. Use of the Plenary Powers for the purpose of making the specific name minimus Miller (J.S.), 1826, as published in the combination Belemnites minimus, the oldest available name for the species concerned (Class Cephalopoda, Order Dibranchia). Opin. int. Comm. zool. Nom. 14: 339–346.

Opinion 431. Use of the Plenary Powers to secure that the generic name *Helicella* Férussac, 1821 (Class Gastropoda) shall be available for use in its accustomed sense. Opin. int. Comm. zool. Nom. 14: 347–372.

Opinion 432. Rejection, as an unpublished proof, of the paper by Binney (W. G.) dated "9th, December 1863" and entitled Synopsis of the Species of Air-breathing Mollusks of North America (confirmation of Ruling given in Opinion 87) and validation under the Plenary Powers of the generic name Carinifex Binney, 1865 (Class Gastropoda). Opin. int. Comm. zool. Nom. 14: 373–392.

Direction 42. Determination of the gender to be attributed to the names of seventy-eight genera of various Classes of Invertebrates and of six genera of the Class Urochorda placed on the "Official List of Generic Names in Zoology" in the period up to the end of 1936. Opin, int. Comm. zool. Nom. 1D: 153–170.

Direction 48. Determination of the authorship to be attributed to the work published anonymously in the year 1798 under the title Museum Boltenianum (Direction supplementary to Opinion 96). Opin. int. Comm. zool. Nom. 1D: 255-264.

Anon, Oyster and mussel culture. [A compilation of various papers.] Rapp. Cons. Explor. Mer 140 3: 1-61.

Anon (1). Pelagic Mollusca: swimmers and drifters. [Report of a meeting of the Malacological Soc. of London] Nature, Lond. 177: 1023-1024.

Anon (2). Calvin Goodrich. [Notice of death.] Nautilus Philad. 68 3 Jan. 1955: 104.

Anon (3). Nenia in the Argentine. Nautilus Philad. 68 3 Jan. 1955: 105.

Anon (4). Family Columbellidae (Con.). [Anachis Strombina etc.] Mins. Conch. Club S. Calif. 154 Jan. 1956: 8-12.

Anon (5). Family Fissurellidae. Mins. Conch. Club S. Calif, 155 Feb. 1956: 3-8.

Anon (6). Family Trochidae. Mins. Conch. Club S. Calif. 156 March-April 1956: 2-8.

Anon (7). Family Bursidae. Family Cyrmatiidae. Family Strombidae. Family Fasciolariidae. Mines, Conch. Club S. Calif. 157 May 1956: 2–6.

Anon (8). Family Epitoniidae Mins. Conch. Club S. Calif. 157 July 1956: 3-6. Anon (9). Family Epitoniidae. Mins. Conch. Club S. Calif., 158 June 1956: 2-6

Anon (10). The feeding of Conus. Mins. Conch Club. S. Calif. 162 Oct. 1956: 9

Anon (11). Taxonomy. Mins. Conch. Club S. Calif. 157 May 1956: 7-8.

Abbott, B. C. & Lowy, J. Mechanical properties of *Pinna* adductor muscle. J. mar. biol. Ass. U.K. **35**; 521-530 6 figs.

Abbott, B. C. & Lowy, J. (1). Resting tension in snail muscle. Nature, Lond. 178: 147-148 1 fig.

Abbott, D. P. Water circulation in the mantle cavity of the owl limpet *Lottia gigantea* Gray. Nautilus, Philad. 693: pp. 79–87 text figs.

A[bbott], R. T. The Kimball Valentine collection [donated to the Academy of National Sciences of Philadelphia. Nautilus, Philad. 69: p. 141.

Abe, N. Colony formation of a limpet, Acmaea dorsursa Gould, and variation of level of the colony. Bull. mar. biol. Sta. Asamushi 7 2-4 1955: 127-132 2 figs.

Ablan, G. L. Lattice method of oyster culture. Philipp. J. Fish, 2 1955: 197-201 3 pls.

Ablan, G. L. (1). Two Philippine boring molluses. Phillip. J. Fish. 2 1955: 191-195 1 pl.

Acolat, L. Nécissité d'une certainé concentration en chlorune de Baryum pour que l'ion Ba soit actif sur le coeur entier d'Escargot (Helix pomatia L.) en survie dans le Ringer. Ann. Sci. Univ. Besançon (2) Zool. & Phys. No. 4 1955: 3-71 fig.

Acolat, L. (1). L'anomie pelure d'Oignou (Anomia ephippium L.), Mollusque interessant au point de vue du mimetisme. Bull. Soc. Hist. nat. Doubs No. 57 1955: 29-32 2 figs.

Acolat, L. (2). Action du chlorure de Baryum sur le coeur entier en survie dans le Ringer, de l'Escargot (*Helix* pomatia L.) et modification de cette action par addition d'hémolymphe. Bull. Soc. Hist. nat. Doubs No. 57 1955: 35–40 2 figs.

- Acolat, L. & Cornier, P. Experimentation sur le coeur de l'Escargot (*Helix pomatia* L.). Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954: 16–19 fig.
- Adam, W. Céphalopodes. Ann. Inst. océanogr. Monaco N.S. **30** 1955: 185–194 l pl.
- Adam, W. (1). Cephalopoda. Partie III. IV. Céphalopodes à l'éxclusion des genres Sepia, Sepiella et Sepioteuthis. Siboga Exped. Monogr. 55C 1954: 119–198 4 pls. 40 figs.
- Adlung, K.-G. & Kauth, H. Prüfung der molluskiziden Figenschaften einiger insektizider Wirkstoffe. Z. angew. Zool. 301–305.
- Aellen, V. & Strinati, P. Matériaux pour une faune cavernicole de la Suisse. Rev. suisse Zool. 63: 183-202.
- Agrawal, S. K. Sur un nouveau sous-geure et deux espèces nouvelles d'Astarte du Jurassique de Kutch (Inde). Bull. Soc. géol. Fr. (6) 5 7–9 1955: 433–439 1 pl.
- Agrawal, S. K. (1). Contribution à l'étude stratigraphique et paléontologique du Jurassique du Kutch (Inde). Ann. cent, et Doc. Paléont. No. 19: pp. 1–188 12 pls. 15 text figs.
- Agrawal, S. K. (2). Deux Lamellibranches intéressants de la série de Katrol dans le Kutch (Inde). Bull. Soc. géol. Fr. (6) 6:13-19 1 pl.

Ahrens, L. see McKerrow, W.

- Aïzenberg, D. E. & Brashnikova, N. E. On the Namur layer in Donetz basin. Bull. Soc. Nat. Moscow geol. 30 1 1955: 37-47. [In Russian.]
- Akimushkin, I. I. New data on geographical distribution of Cephalopods in Far-Eastern seas. C.R. Acad. Sci. URSS N.S. 94 6 1954: 1181–1184. [In Russian.]
- Akimushkin, I. I. (1). Cephelopods in the food of sperm-whale. C.R. Acad. Sci. URSS N.S. 96 3 1954: 665-667. [In Russian.]
- Akimushkin, I. I. (2). Faunistically new Cephalopods for the Far-Eastern Seas of U.S.S.R. C.R. Acad. Sci. URSS 105 2 1955: 367–360. [In Russian.]

Alaphilipps, R. Altération des spermatozoïdes dans la poche copulatrice chez *Planorbarius corneus* (L.) (Mollusca Gastropoda). C.R. Acad. Sci. Paris **241** 1955 : 1225–1227.

[1956]

- Alekgeichik, S. N., Kuzina, I. N., Ratnovski, I. I. Stratigraphy of the Tertinary deposits of Sakhalin. Bull. Soc. Nat. Moscow geol. 29 5 1954: 37-50. [In Russian.]
- Alexeiev, A. K. A Middle Eocene Mollusk Fauna from the Sandstones of Northern Aral Land. Annu. Soc. paléont. russe. 12 [1936–1939] 1945: 17–37 5 pls. [English summary p. 35.]
- Allan, J. The reappearance of a rare Australian volute shell, Amoria grossi Iredale. A new record for New South Wales. Proc. R. zool. Soc. N.S.W. 1954-55 1956: 48-49 l fig.
- Allan, R. S. Report on the Standing Committee on datum-planes in the geological history of the Pacific region. Proc. 8th Pan. Pacif. sci. Congr. 2: 325-423.
- Allen, J. A. On the structure and adaptations of *Pandora inaequivalvis* and *P. pinna*. Quart. J. micr. Sci. **95** 1954: 473–482 5 figs.
- Allen, J. F. The occurrence of estuarine bivalves in an unusual habitat. Nautilus, Philad. 68 1955: 87–90.
- Allen, J. F. (1). A note on Amygdalum papyria Conrad in Maryland waters of Chesapeake Bay. Nautilus, Philad. 68 1955: 83–87.
  - Allison, J. B. see Nelson, T. C.

Almeida, R. C. see Craca, M.

- Altena, C. O. v. R. Bitte um Nachricht. Arch. Molluskenk. 85:199.
- Altena, C. O. v. R. (1). Pyrene scaldensis nova species, a hitherto misunderstood species from the Plio-Pleistocene of Western Europe. Basteria 20: 79–80.
- Altena, C. O. v. R., Bloklander, A. & Pouderoyen, L. P. De fossiele schelpen van de Nederlandse stranden en zeegaten. Basteria 20: 81–90, pls. 9–12.
- Altes, J. Sur le polymorphisme du Gastéropode Leucochroa candidissima Drap. C.R. Soc. Biol. Paris 150: 747-748.

Altes, J. (1). Sur le polymorphisme de la coquille de Leucochroa candidissima modalités et déterminisme. Bull. Mus. Hist. nat. Marseille 16: 53-67 gfis.

Alvarado, R. see Garcia San Nicholás.

Alvarenga, M. see Lopes, H. de S.

Amano, M. Some Upper Cretaceous fossils from southwestern Japan (Part 1). Kumamoto J. Sci. (1B) 2 1: 63-93 2 pls. 1 fig.

Amano, M. see Kobayashi, T.

Ancion, C., Demanet, F. & Leckwijck, W. van. Découverte d'une faune strunienne au toit de la couche d'oligiste oolithique dite famennienne de Couthuin (bord Nord du synclinal de Namur). Bull. Acad. Belg. Cl. Sci. (5) 42:506-514 map.

Anderson, M. M. see Buchanan, J. B.

Andelkovic, M. Z. Quelques espèces caractéristiques d'ammonites et de gastéropodes de l'Albien-Cénomamien et de Senonien à Kosmaj. Ann. géol. Pénin. Balkan 21 1953 : 131–132 I pl.

André, J. L'ultrastructure des mitochondries des spermatocytes de l'Escargot Helix pomatia L. C.R. Acad. Sci. Paris 242: 2048–2051 3 figs.

Andreu, B. & Arté, P. Expériences préalables sur la fixation des larves et la croissance hivernale den jeunes huîtres (Ostrea edulis) den Rias Galiciennes. Rapp. Cons. Explor. Mer 140 3: 17–22 6 figs.

Ansell, A. see Waugh, G. D.

Anteunis, A. Biosocialogische studie van de Belgische Zuduinen. Verband tussen de plantengroei en de Mollusken. fauna. Verh. vlaam. Akad. Wet. te. de Wetensch. Nr. 54: 1-194 figs.

Aoki, S. On the variation of Lima (Acesta) goliath and its allied species, with the description of a new species. Trans. palaeont. Soc. Japan 22: 185–190 l pl. 3 figs.

Archvadze, K. M. Some data on micro-faunistic characteristics of Sakaraul horizon of Abkhazia. Bull. Acad. Sci. U.S.S.R. geol. 6 1954: 101–102. [In Russian.]

Arcisz, W. see Kelly, C. B.

Arkell, W. J. (1). Seven new genera of Jurassic ammonites. Geol. Mag. 90 1 1953: 36-40 pl. 1.

Arkell, W. J. (2). Bajocian ammonites collected by Sir Henry Hayden near Kampadzong, Tibet. Geol. Mag. 90 5 1953: 331–336 pls. 13–14.

Arkell, W. J. & Lucas, G. Découverte récente du genre Ermoceras Douvillé dans l'Atlas sabariens occidental: Conséquences stratigraphiques et paléogéographiques. C.R. Acad. Sci. Paris 236 1953: 2257–2259.

Arsenyev, A. A. Jurassic deposits of Tungussian-Vilyuïsk depression. C.R. Acad. Sci. URSS 105 1 1955: 141-143 map. [In Russian.]

Arté, P. see Andreu, B.

Arvanitaki, A. & Chalazonités, N. Interactions bioélectriques entre aires somatiqués auto-active contiguës (soma géant d'Aplysia?) C.R. Soc. Biol. Paris 150: 700–702 3 figs.

Arvanitaki, A. & Chalazonitis, N. (1). Surexatabilité prénarcotique sur le soma neuronique d'*Aplysia*. (Dérivation endocytaire.) J. Physiol. Paris 48: 374–376.

Arvanitaki, A. & Chalazonitis, N. (2). Biopotentials neuronique à l'éthalle infracellulaire. Stimulation mécanique graducé daus le soma géant d'Aplysia. Bull. Inst. océanogr. Monaco No. 1079: 1–36 figs.

Arvanitaki-Chalazonitis, A. & Chalazonitis, N. Variations lentes et périodiques du potentiel de membrane associées à des groupes de pointes (neurone géant d'Aplysia). C.R. Acad. Sci. Paris 240 1955 : 462–464 3 figs.

Arvanitaki-Chalazonitis, A. & Chalazonitis, N. (1). Les potentiels bioélectriqués endocytainés du neurone géant d'Aplysia en activité autorythmique. C.R. Acad. Sci. Paris 240 1955: 349–351 4 figs.

Arvy, L. & Gaillard, J.-M. Castration parasitairé de Pandora albida (Röding), (P. inaequivalvis L.) mollusque pétécypode eulamellibranche, par Cercaria melanocystea n. sp. cercaire à grande queue résiculeuse. C.R. Acad. Sci. Paris 243 1074–1079 3 figs.

Asano, M., Takayanagi, F. & Kitamura, T. Shellfish poisoning from Callista brevisiphonata Carpenter and its clinical symptons. Tohoku J. agric. Res. 3 1953: 321–330 1 fig.

Astafyeva, K. A. On a new genus of Cardiids. Bull. Soc. Nat. Moscow. geol. 30 3 1955: 94. [In Russian.]

Astakhova, T. V. A contribution to the problem as to the stratigraphic position of *Doricranites*-containing layers. C.R. Acad. Sci. URSS 111: 1065–1067. [In Russian.]

Astre, G. Une date de le paléontologie: La première étude de *Rudistes* par Lapeirouse. Bull. Soc. Hist. nat. Toulouse 89 1954 : 391–400.

Astre, G. (1). Listes stratigraphiques des faunes de Radiolitidés Nord-Pyrénéens. Bull. Soc. Hist. nat. Toulouse 90 1955: 10–14.

Atabekyan, A. A. The discovery of upper Jurassic deposits in the mid-course of the rivers Agstev and Gasan-Tu (Armenia). C.R. Acad. Sci. URSS N.S. 96 3 1954 pp. 589-592. [In Russian.]

Atanasin, L. & Semska, A. Contributions a la connaissance de la faune du calcaire barrémien de Cernavoda-Saliqny. Comun. Acad. Republ. Pop. rom. 1 3 1951: 251–255 3 pls.

Athanassopoulos, & Sur la structure fine des formations dures de certains animaux (coquille des lamellibranches, chitine des malacostracés). Int. Congr. Zool. 14 1953 (1956): 257-260 4 figs.

Atzbach, O. & Geib, K. W. Über einen neuen Fundpunkt im Unteren Meeressand (Mitteloligozän) bei Bretzenheim (Nohe (Mainzer Becken)). Notizbl. hess. Landesamt. Bodenforsch. 83 1955: 237–242 2 figs.

Atzori, M. A. see Bargoni, N.

Aubry, R. La structure de l'acinus et la lignée femelle dans la glande hermaphrodite de *Limnaea stagnalis* adulte. C.R. Soc. Biol. Paris 148 1954: 1498–1580 figs.

Aubry, R. (1). Les éléments nourriciers dans la glande hermaphrodite de Limnaea stagnalis adulte. C.R. Soc. Biol., Paris 148 1954 : 1626-1629 4 figs.

Aubry, R. (2). La lignée mâle dans la glande hermaphrodite de Limnaea stagnalis adulti. C.R. Soc. Biol. Paris 148 1954: 1856–1858 4 figs.

Aubry, R. (3). De la possibilité d'une autofécondation dans l'acinus, hermaphrodite de *Limnaea stagnalis* adulte. C.R. Soc. Biol., Paris 149 1955: 390–392 4 figs.

Aubry, R. (4). Sur le cycle d'élaboration sexuelle dans la glande hermaphrodite de *Limnaea stagnalis* adulti. C.R. Soc. Biol., Paris 149 1955: 2075–2077 5 figs.

Aubry, R. see Bounoure, L.

Audouin, J. see Letaconnoux, R.

Audouin, J. & Letaconnoux, R. Note sur la croissance du *Chlamys varia* L. de l' Ile de Ré. C.R. Ass. franç. Av. Sci. 73 1955: 468-469.

Augustine, D. L. see Chernin, E.

Avnimelech, M. Sur la découverte d'une coquille tropicale lacustre dans des gres Quarternaires de la plaine côtière Palestinienne. C.R. somm. Soc. Géol. Fr. No. 6 : 76-77 I fig.

Avnimelech, M., Parness, A. & Reiss, Z. Mollusca and Foraminifera from the Lower Albian of the Negev (Southern Israel). J. Paleont. 28 1954: 835-839.

Awad, G. H. The Nerineas of Sinai (with a note on the mode of life of these extinct gastropods). Bull. Inst. Desert Egypt 2 2 1952: 23–39 2 pls.

Awad, G. H. (1). Praecardiomya sinaitica a new lamellibranch and genus species from the Middle Cretaceous (Cremonian) of Sinai, Egypt. Bull. Inst. Desert Egypt 2 1 1952: 7–9 6 figs.

Ayers, J. C. Population dynamics of the marine clam, *Mya arenaria*. Limnol. & Oceanogr, 11:26-343 figs.

Azevedo, J. F. de & Faro, M. M. da C. Différence dans en susceptibilité de quelques espèces de *Planorbis* à l'infestion par le *Schistosoma mansoni* de Mozambique. C.R. Soc. Biol., Paris 149: 1848-1851.

Azevedo, J. F. de & Gomes, F. A. C. Utilização dos radioisótopos no esturo do metabolismo em moluscos de água doce (Gastropoda pulmonatu). Garcia de Orta 4: 523-529 8 pls.

Azevedo, J. F. d. & Gomes, F. C. (1). Estudo da eficácia e modo de acção de alguns moluscocidas (Nota previa). An. Inst. Med. trop., Lisbon 13: 579-587 5 pls. [English summary.]

- Azevedo, J. F. d. & Gonçalves, M. M. Ensaios sabre o cotudo da numeração cromosómica de algunas espécies de moluscos de agua doce. An. Inst. Med. trop., Lisbon 13: 569-577 3 pls. [English summary.]
- Azevedo, J. F. de & Medeiros, L. C. M. de. Os moluscos de água doce do Ultramar Português. l.—Introduçãs. Generalidades. Junta Inv. Ultramar. Lisbon 14 1955: 1-95 66 figs. [English summary.]
- Azim, M. A. & Gismann, A. Bilharsiasis survey in South-Western Asia, covering Irak, Israel, Jordan, Lebanon, Saudi Arabia, and Syria in 1950–51. Bull. World Hlth. Org. 14 3: 403–456 7 figs.
- Azuma, M. A new locality for Cypraea (Erosaria) guttata Gmelin. J. Conch. London 24: 146.
- Baba, K. Record of Euselenops (Euselenops) uniceps (Cuvier) from Sagami Bay Japan, Zool. Mag. Tokyo 62 (5) 1953: 189–190. [English summary.]
- Baba, K., Hamatani, I. & Hisai, K. Observations on the spawning habits of some of the Japanese Opisthobranchia. Publ. Seto mar. biol. Lab. 5: 209–219 3 pls., 3 figs.

Bac, M. see Pavlovic.

Bacci, G. Genetical and environmental factors in the sexual differentiation of hermaphrodite molluscs. Ric. sci. 23 Suppl. 1953: 19-31 6 figs. [French, English and German summaries.]

Bădălută, A. see Raileanu, G.

Baden-Powell, D. F. W. The correlation of the Pliocene and Pleistocene marine beds of Britain and the Mediterranean. Proc. Geol. Ass. Lond. 66 271-292 1 map.

- **Bailey, D. W.** Re-examination of the diversity in *Partula taeniata*. Evolution **10**: 360–366 2 figs.
- Baily, J. L. jr. Proposed use of the plenary powers to preserve the generic name *Turbinella* Lamarck, 1799, as the name for the Sacred Chank Shell of India. Bull. zool. Nom. 11: 330-332.
- Baily J. L. jr., (1). Observations on the recently extinct mollusk fauna of Panamint Lake. Nautilus, Philad. 69: 100-103.

- Baird, R. H. A chiton, Callochiton fulvus (Wood), in the River Fal. Nature, Lond. 175 1955: 954.
- Baird, R. H. & Gibson, F. A. Underwater observations on escallop (*Pecten maximus* L.) beds. J. mar, biol. Ass. U.K. 35: 555-562 5 figs.
- Baker, H. B. Familiar names for lands operculates. Nautilus, Philad. 70: 29-31.
- Baker, H. B. (1). Families of Pulmonata [addenda]. Nautilus, Philad. 70: 54.
- B[aker], H. B. (2). Dates of the Nautilus. Nautilus, Philad. 70: 34.
- Baker, H. B. (3). Family names in Pulmonata. Nautilus, Philad. 69: 128-139.
- **B**[aker], **H. B.** (4). Henry Edward Crampton 1875–1956. Nautilus, Philad. **69**: 142.

Bal, D. V. see Palekar, V.

Balashov, Z. G. O sistematicheskom polozhenii i stratighraficheskom znachenii roda *Orchoceras*. Annu. Soc. paléont. russe 15: 223–247 3 pls.

Balashova, E. A. Ghoniatitui Karbona Ber-choghura. Annu. Soc. paléont. russe 14 1953 : 189–205 2 pls.

Balavoine, P. Le gisement fossilifère de Saint-Gervais (Seine-et-Oise). Bull. Mus. Hist. nat. Paris (2) 28: 419-427.

Ballantine, D. & Morton, J. E. Filtering, feeding, and digestion in the Lamellibranch Lasaea rubra. J. Mar. biol. Ass. U.K. 35: 241-274 11 figs.

Bankovski, V. A. & Redichkin, N. A. Schwagerine horizon in the northwest of Donetz basin. C.R. Acad. Sci. URSS 104 3 1955: 456-458 fig. [In Russian.]

Banks, M. R. see Gill, E. D.

- Bargoni, N. & Atzori, M. A. Su alcune proprietà del protidé contrattile del tessuto muscolare del piede di Solen siligua L. R.C. Accad Lincei (8) 21: 210-215 5 figs.
- Barkhatov, B. P. The interrelation between Taurian and Exiordinian layers of the upper Crimea. Bull. Leningrad Univer. 7 1955: 123-136. [In Russian.]
- Barnacle, G. A. S. A new species of *Corilla* from Ceylon. J. Conch. London 24: 95–96 pl. 4.

Basse, E. et Delga, M. D. Presénce en Afrique du nord le Tell constantinois, d'Ammonites nécommienes de la famille des Oostérellidés. C.R. Soc. géol. Fr. 15 1953: 327-330.

Bastide, P see Dastugue, G.

8

Bataller, J. R. Un "Arcomytilus" nuevo del cretácio inferior valenciano. An. Esc. Perit. Agric. Barcelona 2 3 1942: 285-288 figs.

Bataller, J. R. (1). Sinopsis de las especies nuevas del Cretàcico de España. Pars VIII Mollusca 1–3. An. Esc. Perit. Agric. Barcelona 6 1947: 3–186 8 1949: 3–148 figs.; 9 1950: 61–141 figs.

Bataller, J. R. (2). Notes estratigráficos sobre Cardó. An. Esc. Perit. Agric. Barcelona 6 1947: 309–323.

Battaglia, B. Il significato della presenza di polisaccaridi negli spermatozoi atipici dei Gasteropodi Prosobranchi. Ric. sci. 23 Suppl. 1953: 125–129. [French, English & German summaries.]

Baumann, H., see Schwetz, J.

Bayer, F. A. H. Notes on a carnivorous oligochaete commensal on certain fresh-water snails in South Africa. Proc. zool. Soc. Lond. 125 1955: 407-409.

Beets, C. Reconsideration of the so-called Oligocene fauna in the asphaltic deposits of Buton (Malay Archipelago). 1. Mio-Pliocene mollusca. Leid. geol. Meded. 17 1953; 237-256 pl. 1.

Beiftinck, W. G. Vegetatic en Mollusken fauna van de Schorren in het Schelde-Estuarium. Biol. Jaarb. 23: 37-43.

Beldescu, St. & others. L'établissement du poids moyen de quelques espèces du zooplencton de la Mer Noise, par une nouvelle méthode de pesage. Bul. Inst. Cercet. Pisc. Roman. 15: 63-66.

Belskaya, T. N. To the Palaeogeography of Fergan depression at the end of Paleogen. Bull. Acad. Sci. U.S.S.R. geol. 6 1954: 61-74 4 figs. [In Russian.]

Bennett, M. F. see Brown, F. A.

Bensley, C. F. J. The fresh-water Mollusca of the bomb-craters and gunpits on Bookham Common. Lond. Nat. No. 34 1955: 22–28.

Bequaert, J. C. Monograph of the Strophocheilidae, a neotropical family of terrestrial mollusks. Bull. Mus. comp. Zool. 100 1948: 1-210 32 pls.

Berg, K. & Petersen, I. C. Studies on the humic acid Lake Gribsø. Folia limnol. scand. No. 8: 1–273 21 pls. 105 figs.

Bergendahl, E. see Seneca, H.

Bergenhayn, J. R. M. Die fossilen schwedischin Loricaten nebst einer vorlänfigen Revision des Systems der ganzen Klasse Loricata. Acta Univ. Lund. Avd. 2. N.S. 51 8: 1–43 2 pls.

Berkeley-Cotter, J. C. O. Miocénico inarinho de Lisboa. Comun. Serv. geol. Portug. Suplements 36: 1-174.

Berry, S. S. A new West Mexican prosobranch mollusk parasitic on echinoids. Amer. midl. Nat. 56: 355-357 2 figs.

Berry, S. S. (1). A tidal flat on the Vermilion Sea. J. Conch. London 243:81-84.

Beznosov, N. V. Some details in the structure of *Thysanolytoceras* shell. Bull. Soc. Nat. Moscow, geol. **30** 6 1955: 98. [In Russian.]

Bigarella, J. J. Os sambaquis na evolução da paisagen citorânea sulbrasileira. Arq. Biol. Tech. Paraná 9 1954: 199–221 8 figs. [English summary.]

Bigarella, J. J., Tiburtius, G. & Sobanski, H. Contribuição av estudo dos sambaquis do litoral norte de Santa Catarina. I. Situação geográfica e descrição sumária' II. Sambaqui do Rio Pinheiros. Arq. Biol. Tech. Paraná 9 1954: 99–197 102 figs. [English summary.]

Billett, F. & McGee-Russell, S. M. The histochemical localization of  $\beta$  glucuronidase in the digestive gland of the Roman snail (*Helix pomatia*). Quart. J. micr. Sci. 96 1955: 35-48 4 figs.

Binet, L., Jullien, A., Ripplinger, J. & Cardot, J. Sur une substance de choisx, la glucosamine, pour assurer une servie de longue duric du coeur isolé d'Helix pomatia. C.R. Acad. Sci. Paris 240 1955: 1294–1296.

Birkelund, T. Upper Cretaceous Belemnites from West Greenland. Medd. Grønl. 137 9: 1–30 1 pl. 1 f. Blackburn, A. see McKerrow, W.

Blackburn, M. Sonic scattering layers of heteropods. Nature, Lond. 177: 374-375 2 figs.

Blanc, A. C., Settepasci, F. & Tongiargi, E. Excursion on Lac de Massar cinccoli (Plaine côtière de la basse Versilia). IV. Congress internat. étude Quarternaire 1953: 1–29 pls. i–iv.

Blanco, G. J., Villaluz, D. K. & Montalban, H. R. The cultivation and biology of oysters at Baccor Bay. Luzon. Philipp. J. Fish. 1 1951: 35-52 5 pls.

Blaschko, H. & Hope, D. B. The oxidation of L. amino acids by *Mytilus edulis*. Biochem. J. **62**: 335-339.

Blezard, E. Vertical range of snails. North West. Nat. N.S. 3 2 1955: 255–256.

Bloklander, A. see Altena C. O. v. R. others.

Bloklander, A. see Buret L. & ors.

Blom, G. I. To the problem of distinguishing the Vetlyan deposits in the territories of the rivers Suza and Sviyaga in the bounds of Clinvash Republic and the regions adjacent to Tater Republic. C.R. Acad. Sci. URSS 103 5 1955: 887-888. [In Russian.]

Blom, G. I. (1). On apt deposits of the river Kobra basin. C.R. Acad. Sci. URSS 100 6 1955: 1139-1140. [In Russian.]

Blume, W. Systrophia stenogyra zischkai eine neu Lokalrasse aus Bolivien. Archiv Mollusk. 84 4/6 1955: 179–180 text fig.

Bobkova, N. N. Predstaviteli semeřstva Cardiidae Lamarck iz senomanskikh, turonskikh ikon' yakskikh otlozhenii Malogho Kavkaza. Annu. Soc. paléont, russe 15: 125–153 4 pls.

Bodrove, N. V. & Krayukhin, B. K. Intensity of gas-exchange in some Crustacea and Mollusca of the Dnester liman. Bull. Acad. Sci. Kiev. 2 1955: 188-191. [In Ukrainian with Russian summary.]

Boduilevskiř, V. I. K ekologhiř roda Harpaz. Annu. Soc. paléont. russe 15: 167–175 2 pls. Boeters, H. D. Ein Beitrag zur Molluskenfauna der Umgebung von Görlitz. Abh. Naturf. Ges. Görlitz 34 2 1955: 185–187.

Boettger, C. R. Beobachtungen an Nachzuchten von Schlamm-Schneckenarten (Lymnaeidae) in stämmen verschiedener phänotypischer Herkunft. Int. Congr. Zool. 14 1953 (1956): 162– 172 8 figs.

Boettger, C. R. (1). Beiträge zur Systematik der Urmollusken (Amphineura). Verh. dtsch. zool. Ges. 19: 223–256 5 figs.

Boettger, C. R. (2). Objection to A. E. Ellis's proposal to validate *Bithynia* Leach, 1818. (Class Gastropoda). Bull. zool. Nom. 12:21.

Boettger, C. R. (3). Zoogeographische Betrachtungen über die europäischen Süsswasserscknecken der Gattung *Vivi*parus (Montfort). Archiv Mollusk. 84 1/3 1955: 87-95.

Boettger, C. R. (4). Im tropischen Ostafrika eingeschleppte mediterrane Landschnecken. Veröff. Überseemus. Bremen 2A: 331–340 1 pl. 4 figs.

Bogoslovski, B. I. Devonian ammonoide of Rudnoï Altay. Bull. Soc. Nat. Moscow geol. 1 1955: 94-95. [In Russian.]

Bogoslovski, B. I. (1). On the family Pharciceratidae Hyatt, 1900. C.R. Acad. Sci. URSS 103 6 1955: 1103-1106 Pl. and fig. [In Russian.]

Bogoslovski, B. I. (2). On the family Biloclymeniidae, fam. nov.] C.R. Acad. Sci. URSS 104 1 1955: 134–137 fig. [In Russian.]

Bolognari, A. Ulteriori recherche sugli ovociti in accrescimento di *Aplysia depilans* L. (Moll. Gast. Opis.) Arch. Zool. Ital. Napoli 41: 241-259. [English summary.]

Bond, H. W. see Nolan, M.

Bondesen, P. Snegle uden hus. Natur og Mus. 3 2 1954 : 3-10 1 pl. (col.) & figs.

Bonilla, M. G. see Schlocher, J. & others.

Bouchilloux, S. & Roche, J. Sur la pourpae des *Murex* et ses précurseurs. C.R. Soc. Biol., Paris 148 1954: 1583–1587 figs.

Bouchilloux, S. & Roche, J. (1). Contribution à l'étude biochimique de la pourpre des *Murex*. Bull. Inst. océanogr. Monaco No. 1054 1955: 1-23.

Bouchilloux, S. & Roche, J. (2). Sur les prochromogènes et les pigments purpuriques de Murex trunculus Linné. C.R. Soc. Biol., Paris 148 1954: 1732–1734 graph.

Bouillon, J. Influence of temperature on the histological evolution of the ovotestis of *Cepea nemoralis* L. Nature, Lond. 177: 142-143.

Boulet, P. Expériences sur la perception visuelle du mouvement chez Sepia officinalis L. C.R. Soc. Biol. Paris 148 1954: 1486-1489 2 figs.

Boulet, P. (1). Nouvelles expériences sur la perception visuelle du mouvement chez Sepia officinalis L. C.R. Soc. Biol. Paris 149 1955: 1472–1477.

Boullaire, A. C. Observations sur les conditions de vie de *Cypraea tigris* L. en Océanie. J. Conchyliol. Paris **96** 4:159-160.

Bounoure, L & Aubry, R. La structure du canal hermaphrodité de *Limnaea stagnalis* et les possibilité de régénération germinale chez les gastéropodes pulmonés. C.R. Acad. Sci. Paris 243: 1453–1455 4 figs.

Bourdillon, A. Notes sur les Modiolaria habitant la tunique des Ascidies. Rec. Sta. mar. Endoume No. 15 1955: 15-24.

Boury, M. L'Oeuvre de Jean Coste et le développement de l'Ostréiculture. Rapp. Cons. Explor. Mer 140 3 : 7-9.

Bouxin, H. Observations sur le frai de Mytilus edulis var. galloprovincialis (Lmk.), dates précises de frai et facteurs provoquant l'emission de produits génitaux. Rapp. Cons. Explor. Mer 140 3: 43-46.

Bowden, J. & Lowy, J. Innervation. Nature, Lond. 176 1955: 346-347.

Boycott, B. B. & Young, J. Z. Reactions to shape in *Octopus vulgaris* Lamarck. Proc. zool. Soc. Lond. 126: 491-547 28 figs.

Boycott, B. B. & Young, J. Z. (1). The subpedunculate body and nerve and other organs associated with the optic tract of cephalopods. Hanström Festschrift. Lund.: 76–105 f. 1–25.

Boycott, B. B. & Young, J. Z. (2). A memory system in *Octopus vulgaris* Lamarck. Proc. roy. Soc. **143B** 1955: **449**–480 2 pls., 19 figs & tables & graphs.

Bräin, H. Was sind Arion kinkelini Wenz und A. hochheinensis Wenz? Ecl. geol. helv. 49: 593-598 figs.

Brander, T. Zur nomenklatur einiger Skandinavischen Unionazeen. Ark. Zool. 9: 175-185. f. 1-5.

Brandt, R. A. Zur Clausiliidenfauna der Cyrenaika. Arch. Molluskenk. 85 4/6: 121-144 pls. 9-10.

Brashnikova, N. E. see Aizenberg.

Braucourt, J. de H. de. Le fossé tectonique sous le paralléle d'Ishango. [Lists through the text.] Explor. Pare nat. Albert. de Brancourt. No. 1 1955 : 1–150 3 pls., figs.

Brébion, Ph. Au sujet de deux Turritella (Gastropoda) fossiles: T. (Haustator) slansky sp. nov. et T. (Haustator) forgemolli Coq. Bull. Mus. Hist. nat., Paris (2) 27 1955: 414-417 2 figs.

Brébion, P. (1). Turritella (Haustator) doukantensis, nouvelle espèce de gastéropode du Patéocène du Dahomy. Bull. Mus. Hist. nat. Paris (2) 27 1955 : 244– 246 figs.

Breistroffer, M. A propos des genres Jacobella et Hoplites auct. C.R. Soc. géol. Fr. 9 1952: 152-154.

Brien, P. Les oiseaux briseurs de coquilles. Nat. belges 34 9-10 1953: 145-151 1 fig.

Brienne, H. Croissance den moules du Boulonnais. Rapp. Cons. Explor. Mer 140 3:53.

Brown, F. A., Ralph, C. L., Webb, H. M. & Bennett, M. Persistant daily, monthly, and 27-day cycles of activity in the cyster and quahog. J. exp. Zool. 131: 235-262 f. 1-6.

Brown, M. M. & Castell, C. P. Additions to the London Clay fauna of Oxshott, Surrey. Lond. Nat. No. 34 1955: 46-48.

Bruggen, A. C. van. Pyrene sorongensis, a new mollusk (Gastropoda, Prosobranchia) from West New Guinea. Nova Guinea N.S. 7 1:11-14.

- Brunel, A. & Fischer, P.-H. Sur la dégradation des composés d'origine purique chez les Mollusques. Bull. Lab. marit. Dinard No. 40 1954: 33-38.
- Brunnackner M. & K. Die Molluskenfauna einiger Lönprofile im Donautal. Geol. Bl. NO-Bayern 6 3: 91–100 figs.
- Brunson, R. B. The mystery of Discus brunsoni. Nautilus. Philad. 70 1:16-21.
- Bryun, E. S. On the border of the mid and upper Jurassic in the Northern Caucasus. Vestn. Leningr. Univ. biol. geogr. geol. 10 1955: 89-96 fig. [In Russian.]
- Buchanan, J. B. & Anderson, M. M. Additional records to the marine molluscan fauna of the Gold Coast. J. W. Afr. sci. Ass. 1 2 1955: 57-61.
- Buck, H. Zur Verbreitung einiger Gruppen niederer Süsswassertiere in Fliessgewässern Nordwürttembergs (Spongiae, Bryozoa, Turbellaria partim, Oligochaeta partim, Hirudinea, Mollusca, Crustacea partim, Ephemeroptera). Jh. Ver. vaterl. Naturk. Wurttemb. 111 1:153–173.
- Buddenbrack, W. v. Vergleichende Physiologie 3 Basel & Stuttgart (Birkhauser Verlag) pp. 677 205 figs.
- Bukiya, S. G. On the lower Jurassic deposits in the river Okuma basin (Abkhazia.) C.R. Acad. Sci. URSS 100 1 1955: 139-140. [In Russian.]
- Bukiya, S. G. (1). On Datsk stratum in Magrelia. C.R. Acad, Sci. URSS N.S. 99 I 1954: 153-155. [In Russian.]
- Bukiya, S. S. G. (2). On the Lias in North-Western part of Abkhazia. C.R. Acad. Sci. URSS N.S. 99 6 1954: 1073-1075. [In Russian.]
- Bullock, T. H. Temperature sensitivity of some unit synopses [Loligo opalescens]. Publ. Staz. zool. Napoli 28: 305-314.
- Burch, J. B. Distribution of land snails in plant associations in eastern Virginia. Nautilus, Philad. 70 2:60-64 tables.
- Burch, J. Q. Taxonomic characters in Mollusca. Syst. Zool. 5:144.
- Burch, J. Q. (1). The genus in molluscan systematics. Mins. Conch. Club S. Calif. 159 July: 1-2.

- Burch, J. Q. (2), Taxonomic characters in Mollusca. Mins. Conch. Club S. Calif. 160 Aug.: 1-2.
- Burch, J. Q. (3). Some problems in the West Coast Naticidae. Mins. Conch. Club S. Calif. 160 Aug.: 4-7.
- Burch, J. Q. (4). Characters to study in the Classification of Mollusks. Mins. Conch. Club S. Calif. 162 Oct.: 2-8.
- Burch, J. Q. (5). More random notes on molluscan taxonomy. Mins. Conch. Club S. Calif. 163 Nov.: 1–4.
- Bürgl, H. Situacion geologica de la minas de Esmeraldas de Muso, Boyacá. Rev. Acad. colomb. 9:381–388 1 pl. map.
- Bürgl, H. (1), La variabilidad de la Amonita *Dufreynoya texana* Burckhardt. Bol. geol. Colombia 4 2–3: 1–22 3 pls.
- Bürgl, H. (2). Catalogo de las amonitas de Colombia. Part 1. Pulchellüdae. Bol. geol. Colombia 4: 1–119 28 pls. [English summary and descriptions.]
- Burk, C. A. Faunas and age of the Amsden formation in Wyoming. [Moll. pp. 14-15] J. Paleont, 28 1954: 1-16 pl. 1 text-figs. 1-6.
- Butot, L. J. M. Scorpion shells. Penggemar Alam 35 1-3 1955: 71-84 2 pl. [English summary.]
- Cadeo, G. C. & Senna, G. Risultati di alcune ricerche palet nologiche al Buco della Tonda (2005 Lo.) sopra Urio (Como). Natura, Milano 47 4 1956: 168-169.
- Cain, G. L. Studies on cross-fertilization and self-fertilization in *Lymnaea* stagnalis appressa Say. Biol. Bull. Wood's Hole 111: 45-52 2 figs.
- Caire, A. Sur en présence de Creniceras crenatum Boug, daus les marnes oxfordiennes (zone à Cardioceras praecordatum) de Cluey (Jura) et de Turcenay (Doubs). Ann. Sei, Univ. Besançon (2) Géol. No. 2 1955; 129–132.
- Caire, A. (1), Découverte de trois nouveaux échantillons de *Creniceras* crenatum dans la Marnière de Turcenay (Doubs.). Bull. Soc. Hist. nat. Doubs. No. 58:173.

Caire, A. (2). A propos de la découverte d'un rameau de Conifère dans l'Argorien de Dole (Jura). Bull. Soc. Hist. nat. Doubs. No. 57 1955 : 135-144 1 fig.

Caire, A. (3). Sur la présence de Creniceras crenatum Boug. dans les marnes oxfordiennes (zone à Cardioceras praecordatum) de Cluay (Jura) et de Turcenay (Doubs.). Bull. Soc. Hist. nat. Doubs. No. 57 1955: 129–132.

Callomon, J. H. Sections in the Corallian beds at Beckley, Oxfordshire. (Lists.] Proc. Geol. Ass. Lond. 64 2 1953: 83–87.

Callomon, J. H. (1). The Ammonite succession in the Lower Oxford Clay and Kellaways Beds at Kidlington, Oxfordshire, and the zones of the Callovian stage. Philos. Trans. 239B 1955: 215-264 2 pls., 5 figs.

Cambridge, P. G. Notes on Crag palaeontology—1. Trans. Suffolk Nat. Soc. 9: 322-327.

#### Campbell, D. A. see Pratt, D. M.

Campbell, J. D. The Otapirian stage of the Triassic System of New Zealand. Part II. Trans. roy. Soc. N.Z. 84: 45-50 1 fig.

Campbell, J. D. & McKellar, I. C. The Otapirian stage of the Triassic system in New Zealand. Part I. Trans. roy. Soc. N.Z. 83:695-704 1 fig.

Campion, M. A survey of the green algae epiphytic on the shells of some freshwater molluses. Hydrobiologia 8:38-53.

Carasso, N. see Grassé, P.-R.

Carasso, N. see Grassé, P. P. & others.

Cardot, J. see Binet, L. & others.

Cardot, J. see Jullien, A.

Cardot, J. see Jullien, A. & Rerat, D.

Cardot, J. see Jullien & Ripplinger.

Cardot, J. see Jullien, A. & Vieille-Cessay.

Carpenter, L. B. see Miller, A. K.

Carvalho, A. M. d. see Janz, G. J.

Casey, R. Notes on the base of the Gault in Wiltshire. Proc. Geol. Ass. Lond. 66: 231-234.

Castany, G., Coque, R. & Domergue, C. Les plages quaternaires à Cardium des Grands Chott du Sud Tunisien. C.R. Acad. Sci. Paris 236 1953; 2097-2099.

[1956]

Castell, C. P. see Brown, M. M.

Castell, J., Concha, S. de la, & Revilla, J. d. la. Un yacimiento fosilifero del Lias de Maranchón (Guadalajara). Notas Com. Inst. Geol. Min. España No. 40 1955: 25–39 5 pls.

Castellanos, Z. J. A. Catalogo de los Poliplacoforos Argentinos y de aguas vecinas al Estrecho de Magallanes. Rev. Mus. La Plata (Zool.) 6: 465–486 9 pls.

Castellaro, H. A. see Leanza, A. F.

Casteras, M. & Villatte, J. Sur un nouveau gisement de fossiles albiens dans les Corbières Orientales. Bull. Soc. Hist. nat. Toulouse 89 1954: 195–212 7 figs.

Čechovič, V. & Seneš, J. Torton okolia Modrého Kameňa. (Le tortonicon das environs de Modrij Kameň. [Lists.] Geol. Sborn. Slovensk. Akad. 1 1950: 130–166 (Fr. & Russ. summ.) pls. 12–16 5 tables.

Cecioni, G. Leopoldia? paynensis Favre: sua posizione stratigrafica in Patagonia. Atti Soc. ital. Sci. nat. 95: 135-145 1 fig.

Cecioni, G. (1). Distribuzione verticale di Alcune Kossmaticeratidae nella Patagonia Cilena. Bol. Soc. geol. ital 74 1: 141-149 fig.

Chabouis, L. & F. Petite Histoire Naturelle des établishments l'Oceanic. Saint-Amand-Montrond. 2 1954: 1–137 figs. [Moll. pp. 101–121 9 plates.]

Chace, E. P. Unusual high tide collecting. Mins. Conch. Club S. Calif. 162 Oct.: 9.

Chacko, P. I. The first pearl fishery in independent India. Proc. Indian sei. Congr. Ass. 43rd 3: 303 (Abstract).

Chacko, P. I. (1). Prospects for a pearl fishery off Tuticorin, Gulf of Manaar, in 1955. Proc. Indian sci. Congr. Ass. 42nd 3 1955: 309 (Abstract).

Chalazonitis, N. see Arvanitaki-Chalazonitis, A.

Chalazonitis, N. see Arvanitaki, A.

Chambers, T. C. Use of snail stomach cytase in plant cytology. Nature, Lond. 175 1955: 215.

Changeux, J.-P. Melanella comatulicola (Graff) 1874. Vie et Milieu 7. 1956; 105–106.

Chanley, P. E. see Davis, H. C.

Chapman, G. & Newell, G. E. The rôle of the body fluid in the movement of soft-bodied invertebrates. II. The extension of the siphons of Mya arenaria L. and Scrobicularia plana (da Costa). Proc. roy. Soc. 145B: 564-580 11 figs.

Chaput, G. Contribution à l'étude de la faune pliocéne de la Péninsule de Cuide (Turquie). Bull. sci. Bourgogne 15 1955: 39-51.

Chaput, G. & Darkot, B. Sur la Pliocène des environs d'Antalya (Turquie). C.R. Acad. Sci. Paris 236 1953 : 2259-2260.

Chari, S. T. see Venkataraman, R.

Chari, V. K. The horned helmet, Cassis cornula Linn.—an addition to the list of marine gastropods of Bombay. J. Bombay nat. Hist. Soc. 53: 736-737 1 fig.

Charles, R. P. & Maubeuge, P. L. Révision des Liogryphées des Musée d'Histoire Naturelle de Luxembourg. Arch. Inst. Gr.-Duc. Luxumb. N.S. 20 1953: 183-186 l pl.

Charles, R. P. & Manbeuge, P. L. (1). Les Liogryphées jurassiques de l'Est du Bassin Parisien: II. Lyogryphées du Bajocien. Bull. Soc. géol. Fr. 6 2 4-6 1953: 191-196 2 text-plates.

Chavan, A. L'interpretation du genre Ptychomya Agassiz. C.R. Soc. géol. Fr. 9 1952: 161-165 3 text-figs.

Cheesman, D. F. The snail's foot as a langmuir trough. Nature, Lond. 178: 987-988.

Chernin, E., Michelson, E. H. & Augustine, D. L. Studies on the biological control of schistosome-bearing snails. I. The control of Australorbis glabratus populations by the snail, Marsia cornuarietis, under laboratory conditions. II. The control of Australorbis glabratus populations by the leech, Helobdella fusca, under laboratory conditions. Amer. J. trop. Nat. Hyg. 5 2 1956: 297-314 3 figs.

Chernuishev, B. I. Ob *Unio* iz verkhnetriasovuikh otloshenii vostochnoglio sklona Urala. Annu. Soc. paléont. russe 15: 161-165 1 pl. Chétail, M. Caractères histologiques du blastème de régénération tentaculaire chez *Arion rufus*. C.R. Acad. Sci. Paris **242**: 1655–1656.

Chiasson, L. P. see Dickie, L. M.

Chibrikova, E. V. On the age of upper-Saratov deposits. Bull. Acad. Sci. USSR geol. 6 1954: 50-60. [In Russian.]

Chopea, H. G. see Nath, V.

Chow, C. C. see Tang, C.

Chow, S. L. see Tang, C.

Christomanos, A. Nature of the pigment of Aplysia depilaris. Nature, Lond. 175 1955: 310.

**Chubb, L. J.** Rudist assemblages of the Antillean Upper Cretaceous, Bull. Amer. Paleont. **37** 161: 1–23.

Chubb, L. J. Some rarer rudists from Jamaica, B.W.I. Palaeontogr. amer. 4 26:5-30 5 pl.

Chubb, L. J. (1). Thyrastylon, a new rudist genus from the Upper Cretaceous of Guatemala, the Antilles, and Persia, with a discussion of the functions of rudist oscules and pillars. Palaeontogr. amer. 4 27: 35–48 2 pl.

Clarke, jnr. A. H. Two new subspecies of Neptunea decemcostata. Nautilus, Philad. 69 4:115-121 pl. 7 f. 1 2.

Clarke, jnr. A. H. (1). Natural biological control of a  $My\alpha$  predation. Nautilus, Philad. 70 2:37–38.

Clark, R. B. & Milne, A. The sublittoral fauna of two sandy bays on the Isle of Cumbrae, Firth of Clyde. J. Mar. biol. Ass. U.K. 34 1955: 161–180 2 figs.

Claudey, D. see Jullien & others.

Clement, A. C. Experimental studies on germinal localization in *Hyanassa* II. The development of isolated blastomeres. J. exp. Zool. 132: 427-446 f. 1-14.

Clench, W. J. In Johnson F. The Boylston Street Fish weir. Pap. Peabody Foundation 2 1942: 1-212 figs. [Molluscs. p. 45-48.]

Clench, W. J. (1). The occurrence of clines in molluscan populations. Syst. Zool. 3 1954: 122-125

Clench, W. J. (2). Melania cancellata Say. Nautilus, Philad. 68 1955: 107. Clench, W. J. (3). The Unionidae of B.H. and S. H. Wright. Nautilus. Philad. 68 1955 p. 104.

Clench, W. J. (4). A freshwater mollusk survey of North Florida rivers. Nautilus, Philad. 68 1955: 95–98.

Clench, W. J. (5). A new Cerion from Bimini, Bahamas (Mollusca). Amer. Mus. Novit. No. 1794: 1-32 figs.

Clench, W. J. (6). Land shells of Barbada Island, Lesser Antiles. Nautilus. Philad. 70 2:69-70.

Clench, W. J. & Turner, R. D. Freshwater mollusks of Alabama, Georgia, and Florida from the Escumbia to the Suwannee River. Bull. Fla. St. Mus. Biol. Sci. 13:1-239 9 pls.

Clench, W. J. & Turner, R. D. (1). The North American genus *Lioplax* in the Family Viriparidae. Occ. Pap. Mollusks Harvard Univ. 2 1955: 1-20 pl. 1-4.

Coe, W. R. Fluctuations in populations of littoral marine invertebrates. J. mar. Res. 15: 212-232 9 figs.

Cole, H. A. A preliminary study of growth-rate in cockles (*Cardium edule* L.) in relation to commercial exploitation. J. Cons. int. Explor. Mer 22: 77-90 f. 1-5.

Cole, H. A. see Graham, M.

Cole, H. A. & Hancock, D. A. Progress in oyster research in Britain 1949–1954, with special reference to the control of pests and diseases. Rapp. Cons. Explor. Mer 140 3: 24–29.

Cole, H. A. & Hancock, D. A. (1). Odostomia as a pest of oysters and mussels. J. Mar. biol. Ass. U.K. 34 1955: 25-31 2 pls. 1 fig.

Cole, H. A. & Waugh, G. D. Shell disease in Portuguese oysters. Nature, Lond. 178: 422.

Collier, J. R. see Wada, S.

Colman, P. An octopus delicacy and other marine tidbits. Proc. R. zool. Soc. N.S.W. 1954-55 1956: 78-79.

Comfort, A. Mollusca in the alluvia of the River Siagne [List.] [France.] J. Conch. Lond. 24: 97-99.

Concha, S. de la, see Castell, J.

Conley, J. F. see La Rocque.

Coque, R. see Castany, G. & others.

Corbella, C., Croce, N. della & Ravera, O. Plancton, bentos e chimisms delle acque e dei sedimenti in un lago profondo (Lago Maggiore). Mem. Ist. ital. Idrobiol. de Marchi 9 1956: 125-263 26 figs.

Corda, M. Ricerche mecaniche ed elettriche sul cuore isolato di *Helix adspersa*, Boll. Soc. ital. Biol. sper. 31 1955: 1201-1203.

Corda, M. (1). Sul meccanismo dell' azione difasica dell' acetilcolina sul cuore di *Helix adspersa*. Arch. Fisiol. Finenze **56** 1956: 286–291. [English summary.]

Corda, M. (2): Su di un meccanisms di graduazione dell' inotropismo nel cuore di *Helix adspersa*. Arch. Fisiol. Firenze 56: 97-106 6 figs. [English summary.]

Corda, M. see Peruzzi, P.

Cornier, P. see Acolat, L.

Costa, M. see Graca, M.

Cox, L. R. New species of Gastropoda from the Pliocene of Portugal. Mem. Mus. min. geol. Univ. Coimbra 10 1941: 7-12 1 pl.

Cox, L. R. (1). Gastropoda from the Karroo Beds of Southern Rhodesia. Geol. Mag. 90 1953: 201-207 text-figs. 1-2.

Cox, L. R. (2). Lower Cretaceous mollusca from Pointe-à-Pierre, Trinidad. J. Paleont. 28 1954: 622-636 pls. 64-67.

Cox, L. R. (3). Proposed addition to the Official List of Generic Names in Zoology of the generic name Chama Linnaeus 1758. Bull. Zool. Nom. 11: 327-329.

Cox, L. R. (4). Jurassic Mollusca from Peru. J. Paleont. 30: 1179-1186 pls. 127-128.

Cox, L. R. see Eames, F. E.

Cridland, C. C. The experimental infection of several species of African freshwater snails with Schistosoma mansoni and S. haematobium. J. trop. Med. & Hyg. 58 1955: 1-12 map.

Crisp, D. J. see Southward, A. J.

Croce, N. della, see Corbella, C.

Growley, T. E. A new species of Synapterpes from Colombia. J. Conch. Lond. 24: 144 pl. 6.

Cruys, H. Sur Subtissotia peroni Hyatt 1903 = Tissotia africana Peron 1897. C.R. Soc. géol. Fr. 15 1952 : 359–361.

Csepreghyné-Meznerics, I. Die Molluskenfauna von Szob und Letkes. Ann. Hung. Geol. Inst. 45 2: 363–448 14 pls. [German summary.]

Csepreghyné-Meznerics, I. (1). A hazai miocén rétegtani taglatasa az újabb faunavizigálatok alapján. Ann. hist. nat. Mus. hung. N.S. 7: 239–259.

Csutak, V. see Velluda, C.

Cumbie, B. G. see Harry, H. W.

Cunningham, J. A. The horse clam in winter. Victoria Nat. 8 1951: 75-79 2 figs.

Currie, E. D. Scottish Carboniferous Goniatites. Trans. roy. Soc. Edinb. 62 1954: 527-602 4 pls. 9 figs.

Curtis, M. L. K. Type and figured specimens from the Tortworth Inlier, Gloucestershire. Proc. Bristol Nats'. Soc. 29: 147-154.

Cutcliffe, A. S. Rock-ranging along the North Devon coast. No. 2. Marine molluses. Country-Side N.S. 17 11: 466-469.

Cuzzi, G. Su alcune specie di Aptychus del Giura Superiore (Malm) della regione del Canto Alto (Val Brembana). Natura, Milano 47: 22-34 1 pl.

Dan, J. C. see Wada, S.

Dance, S. P. Pisidium obtusale lapponicum Clessin from the lower gravel at Swanscombe, Kent. J. Conch. Lond. 24: 147.

Dance, S. P. (1). A new Sussex locality for *Pisidium pseudosphaerium* Favre. J. Conch. Lond. 24: 91-92.

Danisch, E. Über die Flora und Fauna der Kimmeridge-und Gigas-Schichten zwischen Engter und Venne. Veröff. naturw. Ver. Osnabrück 27 1955: 54-73 1 pl., 2 figs. and map.

Dannevig, A. Lobster and Oyster in Norway. Pp. 92-96 text chart and graphs.

Darkot, B. see Chapeet, E.

Darteville E. & Fischer, P.-H. Une monnaie de coquillage provenant de Formosa. J. Conohyliol. 95 4: 140– 142 text fig. Darteville, E. & Roger, J. Platyodon klinghardti (J. Böhm, 1919), Lamellibranche du Miocène d'Afrique Occidentale. Bull. Mus. Hist. nat. Paris (2) 27 1955: 164-169.

Dastugue, G., Bastide, P. & Moinet, J. C. Déstruction enzymatique de l'ésérine par les humeurs et les organes d'Helix pomatia (L.). C.R. Ass. franç. Av. Sci. 72 1953: 672-674.

Daugherty, J. W. Polyphenol oxidase activity and pigmentation in snail tissues. Biol. Bull. Woods Hole 110: 258-263 1 fig.

David, L. Contribution à la paléontologie stratigraphique du Sénonien de la région de Souk-Ahras (Est-Constantinois). C.R. Soc. géol. Fr. 8 1953: 128–130.

David, L. & Jodot. P. Quelques précisious sur la transgression miocéne dans les environs de Gambetta (Est-constantinois). Bull. Soc. Hist. Nat. Afr. Nord. 46 1955: 121-313 1 pl.

David, L. & Sornay, J. Sur une nouvelle faune d'ammonites du Coniacien de l'Est. Constantinois (Algérie). C.R. Acad. Sci. Paris 242 1956: 532-533.

Davis, A. G. On the occurrence of Pleistocene molluses at Stutton Ness, Suffolk. Proc. Geol. Ass. Lond., 64 1953: 29-30.

Davis, A. G. (1). The Mollusca of Bobbitshole Inter-glacial beds, Ipswich. Trans. Suffolk Nats'. Soc. 9 1955: 110-114.

Davis, A. G. (2). Hygromia (Ponentina) subvirescens (Bellamy) in South Devon. J. Conch. Lond. 24 3:101.

Davis, A. G. (3). Hygromia (Ponentina) subvirescens (Bellamy) in Cornish Holocene deposits. J. Conch. Lond. 24 3: 100.

Davis, A. G. (4). Anodonta minima millet in the Corfe River Dorset. J. Conch. Lond. 24 3: 99.

Davis, A. G. (5). Anodonta minima millet in Dorset. J. Conch. Lond. 24 4: p. 110.

Davis, C. C. The Marine and Freshwater Plankton. [Ann Arbor]. 1955: i-ix, 1-562 681 figs.

Davis, H. C. & Chanley, P. E. Spawning and egg production of oysters and clams. Biol. Bull. Woods Hole 110: 117-128 2 figs.

Dawydoff, C. Contribution à l'étude des invertébrés de la faune marine benthique de l'Indochine. Bull. biol. Suppl. No. 37 1952 : 1–158 2 maps.

Decker, J. M. see Kummel, B.

Dehnel, P. A. Growth rates in latitudinally and vertically separated populations of mytilus californianus. Biol. Bull. Woods Hole 110: 43-53 4 figs.

Delattre, M. Une Bélemnite à épirostre dans le Néocomien. Bull. Soc. géol. Fr. 6 2 4-6 1953 : 283-286 pl. 14a.

Delépine, G. La succession des zones à Goniatites dans les bassins Louillers de l'Afrique du Nord. C.R. Soc. géol. Fr. 10 1952:178-180.

Delga, M. D. see Basse, E.

Dell, R. K. The archibenthal Mollusca of New Zealand. Dom. Mus. Bull., Wellington No. 18: 1-235 27 pls., 6 figs.

Dell, R. K. (1). The fresh-water Mollusca of New Zealand. Part II. The species previously assigned to the genera Limnaea & Myxas. Part III. The genus Physastra. Trans. roy. Soc. N.Z. 84: 71–96 figs.

Demal, J. Essai d'histologie comparée des organes chémorécipteurs des gastéropodes. Mém. Acad. Belg. C. Sci. 8° 29 1 1955: 1–88 7 pls. 12 figs.

Demanet, F. see Ancion, C.

Demoran, W. J. & Gunter, G. Ability of *Thais haemostoma* to regenerate its drilling mechanism. Science 123:1126.

Denis, M. Catalogue des Achatininae existant au museum de Paris. J. Conchyliol. Paris 95 4: 127-139.

Deraniyagala, P. E. P. Some fossils from the Miocene Ampitheatre at Minihagalkauda, Ceylon. Spolia Zeylan. 28: 1-5 3 pls.

**Deschamps, P.** Contribution à l'étude des xylophages marins, Rev. Trav. Inst. Pêches Marit. 20: 185-201 f. 1-7.

Deslandes, N. see Paraense, W. L.

Devidts, J. Sur la présence de *Mitra* (*Episcomitra*) zonata Marryat dans le Golfe de la Ciotat (B.-du.-R.). Bull. Mus. Hist. nat. Marseille 16: 125-129 l pl.

Devries, A. Note préliminaire sur un cas de commensalisme Capulus hemiastericolus nov. sp. chez Hemiaster battensis. C.R. Soc. géol. Fr. 7 1952: 115-116 1 text-fig.

**Dexter, R. W.** Raymond C. Rush. M.D. 1875–1954. Nautilus, Phillad 68 3 1955; 102–104.

Dexter, R. W. (1). Additiona, marine mollusks at Cape Ann. Massachusetts. Nautilus, Philad. 69 4: 140-141.

Dickie, L. M. see Medeof, J. C.

Dickie, L. M. & Chiasson, L. P. Offshore and Newfoundland scallop explorations. Circ. Atlantic biol. Sta. No. 25 1955: 1-4 5 figs.

Dickins, J. M. Permian Pelecypods from the Carnarvon Basin, Western Australia. Aust. Nat. Dept. min. Geol. & Geoph. Bull. 29 1956: 42 pp. text figs & pls. i-vi.

Diehl, M. Die Raubschnecke Velutina velutina als Feind und Bruteinmieter der Ascidie Styela coriacea. Kieler Meeresforsch. 12: 180–185 Pl. 18 19.

Din Sharaf El & H. El Nagar. Control of snails by copper sulphate in the canals of the Gezira irrigated area of the Sudan. J. trop. Med. Hyg. 58 1955: 260-263 map and tables.

Divaris, G. A. see Krijgsman, B. J.

Dobbin, J. E. jr. see Dobrovolny, C.

Dobrovolny, C. G. & Dobbin, J. E., Jr. Trabalhos de campo preliminares realizados com alguns moluscocidas, no Brasil. Bol. Ofic. sanit. panamer. 403: 217-231.

Dodd, J. M. Studies on the biology of limpets. III. Hermaphroditism in the three British species of *Patella*. J. mar. Biol. Ass. U.K. 35: 327-340 1 pl.

Dodd, J. M. see Orton, J.

Dodd, J. M. see Southward, A. J.

Dodge, H. A historical review of the mollusks of Linnaeus. Part 4. The genera *Buccinium* and *Strombus* of the Class Gastropoda. Bull. Amer. Mus. nat. Hist. 111: 153-312.

Domergue, C., see Castany, G. & others.

Dongen, A. v. The preference of Littorina obtusata for Fucaceae. Arch. néerl. Zool. 11: 373-386.

Donovan, D. T. Révision des espèces décrites dans la "Monographie des Ammonites" de P. Reynès (1879). Mém. Soc. géol. Fr. Paléont. 73 1955 : 1–48 2 pls. 4 figs.

Donovan, D. T. (1). The zonal stratigraphy of the Blue Lias around Keynsham, Somerset. Proc. Geol. Ass. Lond. 66: 182–212 text-figs. 1–7.

**Drach, P.** Divers types de croissance des gastéropodes prosobranches. Int. Congr. Zool. **14** 1953 (1956): 442–443.

Drăghindă, I. La présence du Crétacé supérieur dans le bassin de Petroşani. Comun. Acad. Republ. rom. 6: 831-838 10 figs. [French résumé.]

Drushchitz, V. V. Lower Cretaceous Ammonites, Crimea and Northern Caucasus. 1956: 1-150 13 pls. figs. [In Russian.]

Duchateau, G. see Grégoire, C. & others.

Dundee, D. S. see Schalie, H. v. d.

Dupuy, R. see Gaud, J.

Dvořák, J. F. The Cretaceous Formation east of the river Svitava between Letovice and Hradec nad Svit. (Moraria). Rospr. české Akad. 63 1 1953: 1–30 2 pls. [English summary.]

Dzvelaya, M. F. Submerged landslides and crumblings in upper-Miocene deposits of Western Gruzia. C.R. Acad. Sci. URSS N.S. 96 3 1954: 593-596. [In Russian.]

Dzvelaya, M. F. & Maglaperidze, K.S. New data on guriïsk layers of Western Gruzia. C.R. Acad. Sci. URSS N.S. 96 1 1954: 155-157. [In Russian.]

Eager, R. M. C. Naming carboniferous non-marine lamellibranchs. Publ. syst. Ass. No. 2:111-116 1 fig.

Eager, R. M. C. (1). New species of Anthracosiidae in the Lower coal measures of the Pennine region. Mem. Manchester lit. phil. Soc. 95 1954: 40-65 1 pl. 4 figs.

Eames, F. E. Indovoluta, a new Eocene volutid genus. Proc. Malac. Soc. Lond. 32 3:109-110 pl. 22. Eames, F. E. & Cox, L. R. Some Tertiary Pectinacea from East Africa, Persia, and the Mediterranean region. Proc. Malac. Soc. Lond. 22 1/2: 1-68 pls. 1-20.

Ebara, A. Physical studies on the cardiac nerves of a marine mollusc, Dolabella auricula Solander. Sci. Rep. Tokyo Bunrika Daig. 7B 112 1955; 219-230 2 pls. 2 figs.

Eberzin, A. G. Solonovatovodnuie kardiidui pliotzena SSSR. (Cardiidae from the Russian Pliocene.) Trav. Inst. paléont. Acad. Sci. URSS 31 1951: 1-118 17 pls.

Eberzin, A. G. (1). Inserrelationship of the Pliocene lamellibranchiata of Evksin and Caspia. C.R. Acad, Sci. URSS 103 2 1955: 309-312. [In Russian.]

Eberzin, A. G. (2). On some new names for Cardiids for the Pliocene of Ponto-Caspian basin. Bull. Soc. Nat. Moscou geol. 30 3 1955: 93. [In Russian.]

Eckert, H. Die Fauna des oberon Muschelkalkes (Trochitenkalkes) von Wiesloch. I. Pleurotonsariidae und Neritopsidae. Beitr. naturk. Forsch. SudwDtsch. 14 1955: 101-113 3 pls.

Edmondson, C. H. Resistance of woods to marine borers in Hawaiian waters. Bull. Bishop Mus., Honolulu No. 217 1955: 1-91 27 figs.

Eeckhoudt, J. P. Vanden. Quelques aspects de la faune provençale (mammifères - reptiles - batraciens - insectes mollusques). [Moll. pp. 101-103.] Nat. belges 35 4-5 1954: 81-103 9 figs.

Egami, F. see Yagi, Y.

Eglitis, V. K. Soil fauna of the Latvian SSR. Riga 1954.

Eïnor, O. L. The stratigraphy of the Vizeïsk layer of the upper Bashkiria. C.R. Acad. Sci. URSS 103 4 1955: 689-692. [In Russian.]

Eliassen, E. The physiology of the vascular system of invertebrates. I. A monograph on the blood-pigments, their physical-chemical qualities, distribution, and physiological importance. Univ. Bergen Arb. 1953 11 1954: 1-66 25 figs.

Ellis, A. E. Proposed use of the plenary powers to validate the currently accepted usage of the generic name *Unio* Philipsson, 1788 (Class Pelecypoda). Bull. 2001. Nom. 11: 337–343.

[Ellis, A. E.] Editor (1). The Coryndon Museum [E. African marine Bivalves unnamed.] J. Conch. Lond. 24 4: 143.

[Ellis, A. E.] (2). The Hill collection. [William Hill late of Leek, Staffs.] J. Conch. Lond. 24 3:80.

Ellis, E. A. (in paper by J. N. Jennings.) Table of Mollusca from the Alluvium of the Middle Bure Valley. R.G.S. Res. Sci. No. 2 1952: 62-63.

Emerson, W. K. Pleistocene invertebrates from Punta China, Baja, California, Mexico, with remarks on the composition of the Pacific Cosat Quaternary faunas. Bull. Amer. Mus. nat. Hist. 111: 313-342 2 pls. 1 fig.

Emerson, W. K. (1). A new scaphopod mollusk, Dentalium (Tesseracme) hancocki, from the Eastern Pacific. Amer. Mus. Novit. No. 1787: 1-71 fig.

Endeau, R., Stephenson, W. & Kenny, R. The ecology and distribution of intertidal organisms on certain islands off the Queensland coast. Aust. J. mar. & freshw. Res. 7:317-3428 figs.

Entwistle, H. see Johnson, C. G. & others.

Epstein, S. & Lowenstam, H. A. Temperature-shell-growth relations of recent and interglacial Pleistocene shoal-water biota from Bermuda. J. Geol. 61 1953: 424-438 text-figs. 1-8 3 tables.

Erben, H. K. Goniatitacea (Ceph.) aus dem Unterdevon und dem Unteren Mitteldevon. Neues Jb. Geol. Paläont., 98 2 1953: 175–225 pls. 17–19 text-figs. 1–14 1 table.

Erben, H. K. (1). Der Lias und der Dogger Mexikos und ibre intrakontinentalen Beziehungen. Neues Jb. Geol. Paläont. 103: 28-79 4 pls. 4 maps.

Eristavi, M. S. On the subdivision of lower Cretaceous of Crimea. C.R. Acad. Sci. URSS 101 4 1955: 751-753. [In Russian.]

Erlanger, A. A. see Shimanski, V.

Eventov, T. S. see Starobinetz, E.

Fabre, S. Les glauconies turoniennes des bords sud le l'Étang de Berre (B.-du-Rd.) Bull. Soc. lim. Provence 13 2 1942: 10-12.

Fabricius, Å. Anmälan om utplantering av vinbergssnäckan (*Helix pomatia*) in Finland. Memor. Soc. Fauna Flor, fenn. 31: 36.

Fairbanks, L. D. see Fingerman, M.

Farag, I. A. M. Some remarks on the Lower Cretaceous exposures of Gebel Maghara Dome (Northern Sinai). Bull. Inst. Desert Egypt 5 1 1955: 146-163 10 figs.

Farag, J. & Omara, S. On the occurrence of marine Middle Jurassic deposits at Gebel El Minshera (Northern Sinai, Egypt.) Bull. Inst. Desert Egypt 5 1 1955: 165–178 figs.

Farkas, H. & Vágvölgyi, J. On the fauna of the springs of Hungary. I. Mts. Dunazug. Ann. hist. nat. Mus. hung. N.S. 7: 377-384 1 fig.

Faro, M. M. da C. see Azevedo, J. F. de. Faure, L. La coquille saint-Jacques (Pecten maximus L.) de la rode de Brest. Rev. Trav. Inst. Pêches marit. 20:

Favard, P. see Grassé, P.-R.

119-132 f. 1-6.

Favard, P. see Grassé, P. P. & others.

Fedorov, P. V. The modern epoch in the geological history of the Black Sea. C.R. Acad. Sci. URSS 110: 839-841. [In Russian.]

Fedorov, P. V. (1). Some problems connected with Quaternary history of Caspian and Black seas. Bull. Soc. Nat. Moscou geol. 29 5 1954: 21-36. [In Russian.]

Feliksiak, St. Contribution of F. C. Baker to the knowledge of Planorbidae of Poland. Polsk. Arch. Hydrobiol. 3: 163–165. [English summary.]

Fenchel, W. Neue Faunenfundpunkte auf Blatt Wissen/Sieg. Notizbl. hess. Landesamt. Bodenforsch. 83 1955: 92-101 1 fig.

Feniksova, V. V. Stratigraphical significance of the "blue clays" in the Caenozoicun of the mid-Pre-Ob. Bull. Soc. Nat. Moscou geol. 31 (4): 45-54 3 figs. [In Russian.]

Feugueur, L. Présence de caillasses à Potamides tricarinatus au sommet des Calcaires de Saint-Ouen à Herblay (S.-et-O.). Reflexions sur la valeur stratigraphique et la synchronisation de certains termes Bartoniens. Bull. Soc. géol. Fr. 6 2 1953: 373-378 1 text-fig.

Feyling-Hanssen, R. W. Late-Pleistocene deposits at Kapp Wijk, Westspitsbergen. Skr. norsk Polarinst. No. 108 1955: 1–21 3 pls., 7 figs.

Feyling-Hanssen, R. W. (1). Stratigraphy of the marine Late-Pleistocene of Billefjordan, Vestspitsbergen. Skr. norsk Polarinst. No. 107 1955: 1–186 27 pls. 57 figs.

Figueiredo, M. J. R. de. Sobre a associação *Mytilus-Purpura-Balanus*. Naturalia. **5** 1955 : 211–214 2 figs.

Fingerman, M. & Fairbanks, L. D. Osmotic behavior and bleeding of the oyster *Crassostrea virginica*. Tulane Stud. zool. 3: 151–168 11 figs.

Fischer, P.-H. Gastéropodes excavateurs de roches littorales. Int. Congr. Zool. 14 1953 (1956): 455-459.

Fischer, P.-H. (1). Remarques sur la famille des Eustomidae. J. Conchyliol. Paris 96 3: 121-128.

**Fischer, P.-H.** (2). Mollusques récoltés dans la région de Banyuels. J. Conchyliol. Paris **96** 4 : 153–158 fig.

Fischer, P.-H. (3). Note sur Narinea margaritifera D'Arcah. genotype de Campanilopsis N. Gen. J. Conchyliol. Paris 96 1: 3-11.

Fischer, P.-H. (4). Observations sur le compoetement de quelques pulmonés en automne dans le Nord-est de la France. J. Conchyliol. Paris 95 4: 145-153.

Fischer, P.-H. (5). Les Eustoma de la collection de l'École des mines de Paris J. Conchyliol. Paris 96 2 : 49-54 text figs.

Fischer, P.-H. (6). Examen en lumière de Wood de la coquille et de l'épiphragme de quelques pulmoné. J. Conchyliol. Paris 96 1: 46-47.

Fischer, P.-H. see Brunch, A.

Fischer, P.-H. see Dartevelle, E.

Fischer-Piette, E. Répartition. Le long des côtes septentrionale de l'Espagne. Des principles espèces peuplement les roches intercoticaux. [Moll. pp. 96-109.] Ann. Inst. océanogr. Monaco N.S. 31 2 1955: 1-124 5 figs.

Fischer-Piette, E. & Gaillard, J. M. Sur l'ecologie comparée de Gibbula umbilicalis da Costa et Gibbula pennanti Phil. J. Conchyliol. Paris 96 3:115-118 fig.

Fish, G. R. Digestion and the production of sulphuric acid by Mollusca. Nature, Lond. 175 1955: 733-734.

Fisher, L. R., Kon, S. K. & Thompson, S. Y. Vitamin A and carotenoids in certain invertebrates. IV. Mollusca: Loricata, Lamellibranchiata, and Gastropoda. V. Mollusca: Cephalopoda. J. Mar. biol. Ass. U.K. 35: 41-61 63-80 1 fig.

Fleming, C. A. Castecliffim fossils from Ohope Beach, Whakatane (N69). N.Z. J. Sci. Tech. 36B 1955: 511-522 5 figs.

Fliert, J. R. van de. Étude géologique de la region D'Oued Athménia (Algérie). Bull. Cartegéol. Alger, N.S. 3 1955: 1–264 4 pls.

Flint, L. H. Notes on the algal foop of shrimp and oysters. Proc. La. Acad. Sci. 19:11-14.

Florkin, M. see Gregoire, C.

Florschüty F. see Vlerk, I. M. v. d.

Forcart, L. Die nordischen Arten der gattung Vitrina. Arch. Mollusk. 84 4/6 1955: 155-166 3 text figs pl. 12.

Forcart, L. (1). Vitrinidae der Dolomiten. Arch. Molluskenk. 85 4/6: 199.

Forcart, L. (2). Journey to the High Simien (Northern Ethiopia), 1952–3 Three species of *Phenacolimax* (Gastropoda, Vitrinidae), with notes on the taxonomy of the genus. J. Linn. Soc. Zool. 43, 290. 1956 113–122 text figs plates 4.

Forsman, B. Notes on the invertebrate fauna of the Baltic. Ark. Zool. 9 1956: 389-419.

Fort, M. see Schwetz, J.

Fox, D. L. see Goodwin, T. W.

Fraga, F. Average seasonal variation of chemical constituents of the mussel (*Mytilus edulis*). Rapp. Cons. Explor. Mer 140 3:35.

Fraga de Azevedo, J. see Azevedo, J. F. d.

Franc, A. Variations constatées dans la faune des Mollusques de la Baie de Saint-Malo. Bull. Lab. marit. Dinard No. 40 1954: 38–42.

Franc, A. (1). Résultats scientifiques des campagnes de la Calypso. II. Campagne 1951–1952 en Mer Rouge. IX. Mollusques marins. Ann. Inst. océanogr. N.S. 32: 19-60.

- Franc, A. (2). Mollusques terrestres et fluviatiles de l'Archipel Néo-Calédonien, Mém. Mus. Hist. nat. Paris N.S. 13A: 1-200, 24 pls.
- Franzen, A. On the spermiogenesis in Solenogastres. Zool. Bidr. Uppsala 30: 485-491 text-f. 1-13.
- Franzen, A. (1). Comparative morphological investigations into the spermiogenesis among Mollusca. Zool. Bidr. Uppsala 30: 399-456 2 pls. 146 text-f.
- Franzen, A. (2). On spermiogenesis, morphology of the spermatogoon, and biology of fertilization among invertebrates. Zool. Bidr. Uppsala 31: 355-482 6 pls. 223 figs.
- Freanzen, D. S. Types of mollusks described by F. C. Baker part 1, University of Illinois. Nautilus. Philad. 70 1: 21-27.
- Fretag, R. E. see Hunter III G. W. & others.
- Freneix, S. & Magné, J. Une espèce nouvelle de *Pycnodonta* du Crétacé superieur algérien: *Pycnodonta magnei* nov. sp. Bull. Mus. Hist. nat., Paris 28 (2): 244-247 1 pl.
- Fretter, V. The anatomy of the presobranch Circulus striatus (Philippi) and a review of its systematic position. Prec. zool. Soc. Lond. 126: 369-381 5 fgs.
- Friede, R. Weichliere im Röntgenbild. Kosmos, Stuttgart 52: 128–129 7 figs.
  - Frolova, N. V. see Pavlovski, E.
- Frömming, E. Heimische Landschnecken als überträger menschlicher Krankheiten. Kosmos, Stuttgart 51 1955: 535-537 3 figs.
- Frömming, E. (1). Beitrag zur Morphologie und Biologie einer Südwesteuropäischen Süsswasserdeckilschnecke (*Melanopcis dufouri* Fér.). Arch. Hydrobiol. 52: 607–613 3 figs.
- Frömming, E. (2). Experimentelle Untersuchungen über den Einfluss von Umweltfaktoren auf das Gedeihen der Lungenschnecke Ramina decollata (L.) Zool. Jb. (Syst.) 84:577-602 9 figs.
- Frömming, E. (3). Quantitative Untersuchung über die Bedentung bodenbewohnender Landschnecken für den Abbau des Fallaubes. Biol. Zbl. 75:7(5-711 5 figs.

- Frömming, E. (4). Biologie der mitteleuropäischen Süsswasserschnecken. Berlin (Duncker & Humblot): pp. 313 101 figs.
- Frömming, E. (5). Quantitative Untersuchungen über die Nahrungsaufnahme der Landdeckelschnecke *Pomatias elegans* (Müller). Zool, Anz. 157: 59-65.
  - Fujie, T. see Uozumi, S.
  - Furnish, W. M. see Miller, A. K.
- Furon, R. Sur la découverte du Jurassique marin en Afghanistan, C.R. Soc. géol. Fr. 14 1952 : 311-312.
- Gabe, M. Particularités histologiques des cellules neurosécrétrices chez quelques Gastéropodes Opisthobranches. C.R. Acad. Sci. Paris 236 1953: 2166– 2168.
- Gabe, M. (1). La neuro-sécrétion chez les invertébrés. Année biol. (3) 30 1-2 1954 : 5-62 17 figs.
- Gabe, M. (2). Particularités histologiques des cellules neuro-sécrétrices chez quelques lamellibranches. C.R. Acad. Sci. Paris 240 1955: 1810–1812.
  - Gabilly, J. Observations stratigraphiques sur le Lias et le Bajocien inférieur d'Airvault et de Thouars (Deux-Sèvres). C.R. Acad. Sci. Paris 242: 2166-2168.
  - Gabriel, C. J. Mollusca from South East of King Island, Bass Strait. Mem. nat. Mus. Melb. 22 4: 1-6.
- Gabrielyan, A. A. The scheme of Paleogen division in Armenia. C.R. Acad. Sci. URSS 105 4 1955: 790-793. [In Russian.]
- Gaillard, J. M. Lamellibranches recueillis aux Iles Kerguelen et Heard par M.M. Angot, Arétas, Aubert de la Rüe, Brown et Paulian. Bull. Mus. Hist. nat., Paris. (2) 26 1954: 620-623.
- Gaillard, J. M. (1). Gastéropodes recuiellis aux Iles Kerguelen et Heard par MM. Angot, Aubert de la Rüe, Brown et Paulian. Bull. Mus. Hist. nat. Paris (2) 26 1954: 519–525 1 fig.
- Gaillard, J. M. (2). Missions du Batiment Polaire "Commandant Charcot." Récoltés faites en Terre Adélie (1950) par M. Paul Jchernia. III. Mollusques. Bull. Mus. Hist. nat. Paris (2) 26 1955: 678-684 1 fig.

Gaillard, J. M. (3). Variations des peuplements de *Philine aperta* L. et de *Dentalium vulgare* da Costa. Relations avec les parasites de ces espèces. Bull. Lab. marit. Dinard No. 42:86.

Gaillard, J. M. see Arvy, L.

Gaillard, J. M. see Fischer-Piette, E.

Gaillard, J. & J. M. Au sujet de la luminescence chez *Pholas dactylus* L. Bull. Lab. marit. Dinard No. 42 1956: 5-9.

Gajevskaja, N. S. Nutrition and food interrelations of animals inhabiting bottom vegetation and wrecks of the Black Sea. II. Nutrition of the gastropod mollusc Cerithiolum reticulatum (Da Costa). Bull. Soc. Nat. Moscou Biol. 61 5 31-45 3 figs. [English summary.]

Ganapati, P. N. & Nagabhushanam, R. Biology of some marine wood boring organisms of Visakhapatnam Harbour. Proc. Indian sci. Congr. Ass. 42nd 3 1955: 315-316 (Abstract).

Ganapati, P. N. & Nagabhushanam, R. (1). Digestive enzymes of *Martesia strixta* Lim. Curr. Sci. 25 5: 156-157.

Garcia San Nicolás, E. & Alvarado, R. Algunas consideraciones sobre la sistemática del género "Iberus." Bol. Soc. esp. Hist. nat. Biol. 53: 147-149.

Gauzki, R. G., Terektisaisk successions of anticlinals in the Western Primugodzar. Bull. Soc. Nat. Moscou geol. 30 2 1955: 67-82 6 fig. [In Russian.]

Gascoigne, T. Feeding and reproduction in the Limapontiidae. Trans, roy. Soc. Edinb. 63: 129-151 2 pls.-12 figs.

Gasiorowski, S. M. Fauna of Aptychi from the Crinoidal Limestone of the Tithonian-Berriasian near Czorsztyn (Poéning Klippen-Poet). Acta Geol. Polonica 6 3 1956 pp. 71-74 plate and 2 text figs.

Gaud, J. & Dupuy, R. Rhythmes saisonmers de développement de *Bulinus truvcatus* en élevage du Laboratoire. Bull. Inst. Hyg. Maroc 14 1–2 1954: 111–116 2 figs.

Gaud, J. & Dupuy, R. (1). Rythmes de développement de Bulinus truncatus en élevage un laboratoire. Ann. Parasit. hum. comp. 30 1955: 62-68 tables & graphs.

Gautier, Y. Bryozoaires des Gastéropodes de l'herbier de Posidonies, Vie et Milien 6 1955 : 335-341.

Gavrilov, G. B. Multiplication of the Baikal molluse *Baicalia herderiana* Ldh. and others.] Zool. Zh. 32 5 1953; 870–873. [In Russian.]

Gavrish, V. K. The Paleocene deposits of Dnepr-Donetz depression. Bull. Soc. Nat. Moscou biol. 31 (5) 111-116 5 figs. [In Russian.]

Geib, K. W. see Atzbach, O.

Geldiay, R. Studies on local populations of the freshwater limpet Ancylus fluviatilis Müller. J. Anim. Ecol. 25: 389-402 5 figs.

George, J. C. On the significance of the action of the bile salt sodium taurocholate on marine eggs. Proc. Indian sei. Congr. Ass. 42nd 3 1955: 280 (Abstract).

Geyer, O. F. Über Diceras speciosum (Münster) im Weissen Jura von Württemberg. Paläont. Z. 27 1953: 208–211 pl. 13.

Geyer, O. F. (1). Die Fauna der oolithischen Trümmerkalke des oberen Malm in Württemberg und ihre Beziehungen zur korallogenen Fazies des Tithon. [Lists pp. 137–140.] N. Jb. Geol. Paläont. B., 3 1953: 130–140 1 geol. sect. 1 table.

Ghlazunova, A. E. Nekotoruie Ammonitui verkhnegho al'ya Kopet-Dagha. Annu. Soc. paléont. russe 13 1949: 43-53 1 pl. figs.

Ghoretzkii, V. A. Rod Pectunculus iz otlozhenii tortonskogho yarusa yughozapadnoi okranui russkoi platformui. Geol. Sbornik 2–3: 194–202 3 pls. [In Russian.]

Gianfranco, B. Contributo alla conoscenza della Malaco fauna del Marc Toscano-Nota I. -Lamellibranchi. Boll. Pesca Piscic. Idrobiol. 32 fasc. 1 1956 [1957]: 56–82 6 figs.

Giannini, E. Molluschi Eocenici della Migiurtinia. Palaeontogr. Ital. 32 suppl. 7: 23-45 3 pls.

Gibson, F. A. Escallops (Pecten maximus L.) in Irish waters. Sci. Proc. R. Dublin Soc. 27: 253-271 13 figs.

Gibson, F. A. see Baird, R. H.

- Gidon, P. Une Ammonite dans l'Argonien de Grande-Chartreuse. C.R. Soc. géol. Fr. 12 1952 : 237–238.
- Gill, E. D. & Banks, M. R. Cainozoic history of Mowbray Swamp and other areas of North-Western Tasmania. Rec. Q. Vict. Mus. N.S. No. 6: 1-41 6 pls.

Gismann, A. see Asim, M. A.

- Givulescu, R. Contributions à la connaissance de la faune du Crétacé supérieur de Borod-Cornitel (Rég. Bihor). Stud. Cerçet. Acad. Române. Fil. Cluj. 2 1–2 1951: 104–107. [French resumé.]
- Glaçon, G. et J. Etudes de quelques Nérinées et Milioles du Crétacé inférieur des monts du Hodua. Bull. Soc. géol. Fr. 6 3 1-3 1953: 23-34 1 plate 1 text-fig.
- Glahn, P. E., Manchon, P. & Roche, J. Sur les 1-aminoacideoxydases de l'hépatopancréas des lamellibranches, C.R. Soc. Biol. Paris 149 1955: 509-513.
- Glibert, M. Pleurotomes du Miocène de la Belgique et du Bassin de la Loire. Mém. Inst. Sci. nat. Belg. No. 129 1954: 1-75 7 pls.

Glushko, V. V. see Sandler.

- Göke, G. Mikroskopische Untersuchunzen an Belemniten. Mikrokosmos 45: 121–123 5 figs.
  - Gomes, F. A. C. see Azevedo, J. F. de.
  - Goncalves, M. M. see Azevedo, J. F. d.
- Goodhart, C. B. Genetic stability in populations of the polymorphic snail, Cepaea nemoralis (L.). Proc. Linn. Soc. Lond. 167 1:50-67 6 figs.
- Goodwin, T. W. & Fox, D. L. Some unusual caratenoids from two nudibranch slugs and a lampoid fish. Nature, Lond. 175 1955: 1086-1087.
- Goretzki, G. I. The lower border of the Quarternary period. Bull. Soc. Nat. Moscow geol. 31 (4): 29-43 5 figs. [In Russian.]
- Goretzki, G. I. (1). The age relationships of the sediments in Uzunlarsk and Karangat transgression. Bull. Soc. Nat. Moscow. geol. 30 2 1955: 13—29. [In Russian.]
- Goumard, J. Nouvelle espèce d'Unio plissé du Crétacé supérieur du Soudan Francais; *Plicatounio radieri* nov. sp. Bull. Mus. hist. Nat. Paris (2) 28: 248–254 3 figs.

- Graça, M. E. da S., Almeida, R. C. & Costa, M. R. M. The chemical composition and nutritional value of Portugese oysters (*Gryphaea angulata*]. Rapp. Cons. Explor. Mer 140 3: 38-42.
- Gräf, I. E. Die Fährten von *Littorina* littorea Linné (Gastr.) in verschiedenen Sedimenten. Senckenbergiana. leth. 37:305–317 2 pls. 2 figs.
- Graham, M. & others. [Mollusca by H. A. Cole.] Sea fisheries, their investigation in the United Kingdom. London (Edward Arnold) pp. xii + 487 text illust.
- Gramm, M. N. The transsection of the Neogene deposits in North Western part of Minbulak depression (Kizil-Kumi). C.R. Acad. Sci URSS 103 4 1955; 677-680. [In Russian.]
- Grangeon, P. & Jodot, P. Mollusques pliocènes des Sables volcaniques fluviatiles de Mirabel (Ardèche). Ann. Univ. Lyon (3) 9C: 47-72 3 figs.
- Grassé, P.-P., Carasso, N. & Favard, P. L'ultrastructure de la spermatide de l'Escargot. (Helix pomatia L.): Chromosomes, enveloppes nucléaires, centrosome. C.R. Acad. Sci. Paris 241 1955: 1430-1432 2 pls.
- Grassé, P.-P., Carasso, N. & Favard, P. (1). L'ultrastructure des chromosomes et son évolution au cours de la spermiogenèse d'*Helix pomatia*. II. La spermatosoïde, C.R. Acad. Sci. Paris 242: 1395–1398 2 pls.
- Grassé, P.-P., Carasso, N. & Favard, P. (2). Evolution de l'ultrastructure mitochondriale au cours de la formation du segment intermédiaire du spermatozoïde de l'Escargot Helix pomatia L. C.R. Acad. Sci. Paris 242: 1933–1935.
- Grassé, P.-P., Carasso, N., & Favard, P. (3). Les ultrastructures cellulaires du cours de la spermiogenèse de l'escargot (Helix pomatia L.): Évolution des chromosomes, du chondriome, de l'appareil de golgi, etc. Ann. Sci. nat. (Zool.) (11) 18: 339-380 pl. 1-11 f. 1-IX.
- Grayson, J. F. The conversion of calcite to fluorite. Micropaleontology 2 1:71-78 text-figs 1-9 1 table.

Graziadei, P. see Rossi, F.

- Grégoire, C., Duchateau, G. & Florkin, M. La trame des nacres et des perles. Ann. Inst. Océanogr. Monaco 31 1 1955: 1–36 33 pls.
- Grensted, L. W. Effects of a gale on the south Lancashire coast. J. Conch. Lond. 24 4: 145-146.
- Gresson, R. A. R., Threadgold, L. & Stinson, N. The golgi elements of the neurones of *Helix*, *Locusta* and *Lumbricus*. Cellule 58 fau 1:5-16 1 pl.
- Griffiths, R. J. Cypraea in North-West Malaya. J. Conch. Lond. 24 3: 85-90 tables & pl. 3.
- Grijns, A. Bacteriological control of Lealand Oysters and Mussels.' pp. 48-49.
- Grossu, A. V. Potamopyrgus jenkinsi, gastéropode nouveau pour les eaux continentales de la Republique Populaire Roumaine. Comun. Acad. Republ. Pop. rom, 1 7 1951: 593-596 1 fig. [French summary.]
- Grossu, A. V. (1). Contributions systématiques et zoogeographiques à la connaissance des gastéropodes de la vallée du Miniş-Banat. Comun. Acad. Republ. Pop. rom. 5 4 1955: 665-669 1 fig. [French summary.]
- Grossu, A. V. (2). Contribution à la connaissance de la faune des Gastéropodes de la Republique Populaire Roumaine: Hohenwarthia orghidani n.sp. de la famille Ferussaciidae. Comun. Acad. Republ. Pop. rom. 5 11 1955: 1579–1582 1 fig. [French summary.]
- Grossu, A. V. (3). Gastropoda Prosobranchiasi Opisthobranchia. Fauna Rep. Pop. Romine 3 fasc. 2: 1–220.
- Grossu, A. V. & Paladian, G. Contribution à l'étude des mollusques du delta et du bassin du bas Danube. Anal. Inst. Cerc. pisc. Romane N.S. 1 1956: 401-410.
  - Gryaznov, V. I. see Karlov, N.
- Gunter, G. The problem in oyster taxonomy. Syst. Zool. 3 1954:134-137.
  - Gunter, G. see Demoran, W. J.
  - Gupta, B. L. see Nath, V.
  - Gurumani, O. N. see Nair, N. B.
- Gustomesov, V. A. To the ecology of the belemnites in Russian Platform. Bull. Soc. Nat. Moscou geol. 31 (3): 113-114. [In Russian.]

- Guyon, M. see Jullien & Ripplinger.
- Guyon, M. see Jullien, A. & others.
- Haas, F. Binnenschnecken aus einer Karbon-Landschaft im unteren Amazonasgebiete. Arch. Mollusk. 84 1/3 1955: 101–105 6 text figs.
- Haas, F. (1). An Eocene Aturia from Cyrenaica. Amer. Mus. Novit. No. 1747 1955: 1-4 figs. 1-3.
- Habe, T. Deposit of shell remains in Ofanato Bay Iwate Reef, Northern Japan. Venus, Fukuyama 18 4 1955: 244-251. [In Japanese with English summary.]
- Habe, T. (1). Notes on three Tellinid species described by V. Bertin. Venus, Fukuyama 18 4 1955: 240-243 text figs 1-9.
- Habe, T. (2). Anatomical studies on Japanese Land snails (3). Venus, Fukuyama 18 4 1955: 221. [In Japanese with English summary.]
- Habe, T. (3). Spawning of Cerithidea djadjariensis and C. rhizophorarum. Venus, Fukuyama 18 3 1955: 204–205 text figs 1–6. [In Japanese.]
- Habe, T. (4). The breeding of Nodilitorina granularis (Gray). Venus. Fukuyama. 18 3 1955: 206-207 text figs. 1 2. [In Japanese with English abstract.]
- Habe, T. (5). Notes on Pedicularia (Pediculariona) stylasteris Hedley var. Venus, Fukuyama 18 3 1955: 157–160 text figs. 1–6. [In Japanese with English summary.]
- Habe, T. (6). Anatomical studies on the Japanese land snails (6). The superspecific groups of the family Enidae. Zool. Mag. Tokyo 65: 262–266 13 figs.
- Habe, T. (7). Notes on four Korean land snails. Zool. Mag. Tokyo 65: 188-193.
- Habe, T. (8). The floating egg capsules of the Japanese periwinkles (Littorinidae). Venus, Fukuyama 19 2: 117-121 text figs.
- Habe, T. (9). Notes on the systematic position of three American sea shells. Venus, Fukuyama 19 2: 95-100 text figs.

Habe, T. (10). Anatomical studies on the Japanese Land snails (4) Genera of the Family Succinidae. Venus, Fukuyama 19 2: 100-108 text figs.

Habe, T. (11). Anatomical studies on the Japanese Land snails (7). Venus, Fukuyama, 19 2: 109-117.

Habe, T. see Taki, I.

Habe, T. see Yamaguaki, N.

Haffner, K. v. Über Muscheln der Gattung Vulsella als Einmieter in Badeschwämmen und die Bedeutung des Zusammenlebens. Zool. Jb. (Anat.) 75: 1–38 32 figs.

Hagn, H. Sur Kenntnis des Unterea Obercampans (Zone der Belemnitella mucronata [Schloth]. mut. senior Nowak in Südbayern. Neues Jb. Min. Geol. Paläont. 96 2 1953; 304–338 p. 8.

Hall, D. N. F. Ink ejection by Cephalopoda. Nature, Lond. 177: 663.

Hallam, A. The palaeontology and stratigraphy of the Marlstone rock-bed in Leicestershire. [Moll. p. 21.] Trans. Leicester lit. phil. Soc. 49 1955: 17-35.

Halm, H. van. Zeenwse fossielen II. Natura, Amst. 53: 138-142.

Hamai, I. Systematic relative growth in molluscan shells (5) Regularity of direction of variation. Venus, Fukuyama 18 3 1955 pp. 189–197 tables and graphs. [In Japanese with English summary.]

Hamatani, I. see Baba, K.

Hancock, D. A. The structure of the capsule and the hatching process in *Urosalpinx cinerea* (Say). Proc. zool. Soc. Lond., 127: 565-571 1 fig.

Hancock, D. A. see Cole, H. A.

Hancock, D. A. see Mistakidis, M. N.

Hanna, G. D. Preparation of Nudibranchiata. Nautilus, Philad. 68 3 1955: 105–106.

Hanna, G. D. (1). Land and freshwater mollusks of the Arctic slope, Alaska. Nautilus, Philad. 701:10 pl. 1.

Hans, V. Tortouská fauna od Kosihoviee na južnom Slovensku. (La faune tortonienne de kosihavce en slovaquia meridionala). [List.] Geol. Sborn. Slovensk. Akad. 1 1950: 70–73. French and Russian summaries.]

Hara, R. see Hara, T.

Hara, T. & Hara, R. Uber die Pupillarreaktion var der Schnecke Euhadra amaliae (Kobett). Zool. Mag. Tokyo 62 (2) 1953: 48-51 figs.

Haranghy, L. Effect of 3, 4-benzopyrene on fresh-water mussels. Acta biol. Acad. Sci. hung. 7 1 1956:101-108. [English summary.]

Harry, H. W. & Cumbie, B. G. The relation of physiography to the types of freshwater environments and the presence of Australorbis glabratus in Puerto Rico. Amer. J. trop. Med. Hyg. 5: 742-756 6 figs.

Harry, H. W. & Cumbie, B. G. (1). Stream gradient as a criteria of lotic habitats siutable for Australorbis glabratus in Puerto Rico. Amer. J. trop. Med. Hyg. 5: 921-928 4 figs.

Hässlein, L. Mollusken und Molluskengesellschaften der Gewässer des Nördlinger Rieses, Jh. Ver. vaterl. Naturk, Württemb. 111 1:174-199.

Hatanaka, M. see Imai, T.

Havinga, B. Mussel culture in the Dutch Waddensea. Rapp. Cons. Explor. Mer 140 3: 49-52.

Hayasaka, S. A preliminary report on the local variation of a periwinkle, Littorina brevicula (Philippi). Res. Bull. Saito Ho-on Kai Mus. No. 24 1955: 35-37 3 figs.

Hayasaka, I. & S. On a Cretaceous species of Ostrea from Hokkaido, with special reference to its mode of occurrence. Jap. J. Geol. Geogr. 27:161-1651 pl. 3 figs.

Hayashi, Y. On the variation of *Corbicula* due to environmental factors. Venus, Fukuyama 19 1: 54-60 text figs.

Hayward, J. F. A preliminary note on certain granules from port-glacial deposits in the Leas Valley. J. Conch. Lond. 24 4: 139.

Heide, S. van der. Quelques remarques sur *Spirorbis pusillus* (Martin) du Carbonifere des Pays-Bas. Leid. geol. Meded. **20** 1955 [1956]: 100–109 3 pl.

Hemming, C. F. & Verdcourt, B. Notes on the plant and molluscan ecology of a saline desert area in Italian Somaliland including the description of a new species of *Cleopatra* (Mollusca). Rev. zool. Bot. Afr. 53: 57-64 4 figs.

Hemming, F. Note on Zoological nomenclature. J. Paleont. 28 1954:693.

Hepper, B. T. The European flat oyster, Ostrea edulis L., as a host for Mytilicola intestinalis Steuer. J. Anim. Ecol. 25: 144-147.

Hertel, R. Praehistorische Fund von Margaritana auricularia Spglr. und Margaritana margaritifera L. in Sachsen. Abh. Mus. Tierk. Dresden 23: 1-7 8 figs.

Hertlein, L. G. Cretaceous ammorite of Franciscan Group, Marin County, California. Bull. Amer. Assoc. Petrol. Geol. 40: 1985–1988 1 pl.

Hess, O. Beobachtungen zur Normogenese des Süsswässer-Prosobranchiers Bithynia tentaculata L. Biol. Zbl. 75: 664–682 10 figs.

Hess, O. (1). Die Entwicklung von Halbkeimen bei dem süsswasser-Prosobranchier *Bithynia tentaculata* L. Roux. Arch. Entw. Mech. Organ. 148: 336–361 13 figs.

Hess, O. (2). Die Entwicklung von Exogastrulakeimen bei dem Susswasser-Prosobranchier *Bithynia tentaculata* L. Roux. Arch. Entw. Mech. Organ. 148: 474–488 9 figs.

Hidalgo, E. Algunos moluscos de la Isla de Cozumel, Quintana Roo, Mexico. Acta zool. Mexicana 1 No. 10: 1-126 pls. [English summary.]

Hikita, T. & T. On the rare striking squid, Galiteuthis armata Joubin, found in Hokkaido, Japan. Zool. Mag., Tokyo 65: 27-29. [English summary.]

Hipkins, A. K. The genus Rissoina d'Orbigny, 1840. Bull. Conch. Sect. Auck. Mus. 12: 5-7 f. 1-7.

Hirayama, K. Tertiary Mollusca from Hikoshima, Yamaguchi Prefecture, Japan, with remarks on the geological age of the "Ashiya fauna." Sci. Rep. Tokyo Kyoiku Daig. 5C 45: 81-127 3 pls.

Hiregaudar, L. S. see Kathuria, J. B.

Hisai, K. see Baba, K.

Hodgkin, A. L. & Keynes, R. D. Active transport of cations in giant axons from Sepia and Loligo. J. Physiol. 128 1955: 28-60 13 figs.

Hodgkin, A. L. & Keynes, R. D. (1). The potassium permeability of a giant nerve fibre. J. Physiol. 128 1955: 61-88 8 figs. Hölder, H. Oxytoma scanica (Lundgren) in der schwäbischen Planorbiszone. N. Jb. Geol. Paläont. B. 8 1953: 358-364 text-figs 1-6.

Hölder, H. (1). Ein Austernfund im schwäbischen Rhät. N. Jb. Geol. Paläont. B., 4 1953: 185–189 text-figs. 1-2.

Holland, H. Sur le Tournaison de la vallée du Dra (Sud marocain). C.R. Acad. Sci. Paris 242: 2752-2755.

Holme, N. A. see Morton, J. E.

Holmes, W. The colour changes of cephalopods. Endeavour 14 54 1955: 78-82 9 figs. (col.).

Honjo, I. Reizphysiologische Untersuchungen von Pinctata [Pinctada] martensi (Dunken) II. Chemischer Sinn. Zool. Mag. Tokyo 62 (2) 1953: 31–35. [In Japanese.]

Hope, D. B. see Blaschko, H.

Hopkins, S. H. Odostomia imprissa parasitizing Southern oysters. Science 124: 628–629.

Horne, R. see Lacy, D.

Howard, A. D. Pteropod mollusks from the Caribbean. Mins. Conch. Club S. Calif. No. 3 1936: 78.

Howarth, M. K. The Scalpa Sandstones of the Isle of Raasay, Inner Hebrides. Proc. Yorks. geol. Soc. 30: 353-370 figs.

Howarth, M. K. Domerian of the Yorkshire Coast. Proc. Yorks. geol. Soc. 30 1955: 147-1751 pl. figs.

Hoyle, G. & Lowy, J. The paradox of *Mytilus* muscle. A new interpretation. J. exp. Biol. 33: 295-310 13 figs.

Hubbs, C. L. Support for the proposal made by Joshua L. Baily, Jr., to preserve the generic name *Turbinella* Lamarck, 1799, as the sacred chank shell of India. Bull. zool. Nom. 12: 251-.

Hubendick, B. Report from Professor T. Gislen's Expedition to Australia in 1951-1952. 14. On a small material of fresh water molluses collected by Prof. T. Gislen in Australia. Acta Univ. Lund. Avd. 2 N.S. 51 6: 1-11 19 figs.

Hubendick, B. (1). Observations on the occurrence of *Oncomelania quadrasi* Möllendorff, the host-snail of Bilharzia in the Philippines. Int. Congr. Zool. 14 1953 (1956): 511-512. Hubendick, B. (2). Våra Snäckor. Illustrerad Handbok. Snäckor i sött och bräckt vatten. Stockholm (A. Bonniers Förlag) 1949 pp. 100 4 pls. (1 col.) 172 figs.

Hubendick, B. (3). A conchological survey of the genus *Plecotrema*, (Gastropoda, Ellobiidae). Proc. Malac. Soc. Lond. 32 3: 110-126 fig. text maps, pl. 23.

Hubricht, L. Land snails of Shenardoah National Park. Nautilus, Philad. 70 1:15-16.

Hubricht, L. (1). Haplotrema kendeighi Webb. Nautilus, Philad. 69 4: 126-128.

Hubricht, L. (2). Land snails from Louisiana. [Megapallifera subgen. nov.] Nautilus, Philad. 69 4: 124-126.

Hubricht, L. (3). Mesodon normalis (Pils.). Nautilus, Philad. 69 4: 140.

Hubricht, L. see Pilsbry, H. A.

Hudec, V. Die Molluskenfauna im südwestlichen Teil das Javornik-Gebirges. Přírod. Sborn. ostrask. Kraje 173: 390–396. See also p. 425–.

Hudec, V. (1). Delima ornata Rossm. u. Boskovic a u Vratikova. Cas. nár. Mus. 124 2 1955 : 212–214 f. 1.

Hughes, G. M. & Kerkut, G. H. Electrical activity in a slug ganglion in relation to the concentration of Locke Solution. J. exp. Biol. 33: 282-294 10 figs.

Hunter, W. D. R. Studies on the adaptation of freshwater Gastropoda. Summ. Thesis Univ. Glasgow 1953-54 1954: 54-55.

Hunter, W. R. (1). Endemicism in the snails of Jamaica. Glasg. Nat. 17 1955: 173-183.

Hunter, W. R. (2). New and newly-confirmed distribution records of non-marine Molluses in the West of Scotland (IHrd paper). Glasg. Nat. 17 1955: 207–211.

Hunter, W. R. & M. R. Mollusca of Scottish Mountains. J. Conch. Lond. 24 3:80.

Hunter, W. R. & M. R. (1). A note on Plumatella and Ancylus in a mountain Lochan. Glasg. Nat. 17: 241-242.

Hunters, G. W. see Mackie, T. T. & others.

Huvé, P. Contribution à l'étude biologique d'une crique du Golfe de Marseille: L'Anse des cuivres, I.— Repartition des *Monodonta* et *Gibbula* (mollusques prosobranches). Rec. Trav. Sta. mar. Endoume No. 18: 39–573 pls.

Ichikawa, K. Trigonucula (nov.) and other taxodont pelecypods from the Upper Triassic of the Sakawa Basin in Shikoku, Japan. Jap. J. Geol. & Geogr. 21 1949: 267-274 1 pl.

Ichikawa, K. see Kobayashi, T.

Illies, H. Der mittlere Dogger im badischen Oberrheingebiet. Ber. naturf. Ges. Freiburg i. Br. 46: 5-52 1 pl. 11 figs.

Imai, T. Hatanaka, N., Sato, R., Sakai, S. & Yuki, R. Artificial breeding of oysters in tanks. Tohoku J. agric. Res. 1 1950: 69-86 4 figs.

Imai, T., Sakai, S. & Okada, H. Transplantation of European flat oyster, O. edulis into Japanese water and it's breeding in tanks. Tohoku J. agric. Res. 3 1953: 311-320 6 figs.

Imanishi, S. Notes on the Chiebun Formation, a newly found Upper Cretaceous deposit near Nayoro-Machi, North Hokkaido. Kumamoto J. Sci. (1B) 21:57-60 3 figs.

Imanishi, S. On the occurrence of *Trigonia* bearing sandstone at Hôronbetsu, Utanobori-Mura, Esushi-Gun, North Hokkaido. Kumamoto J. Sci. (1B) 2 1:49-55 2 pls.

Imlay, R. W. Barremian ammonites from Trinidad, B.W.I. J. Paleont. 28 1954: 662-667 pls. 74-75.

Imlay, R. W. (1). Stratigraphia and geographic range of the early Cretaceous ammonite *Homolsomites*. J. Paleont. 30: 1143-1146 pl.

Imlay, R. W. see Schlocker, J. & others.

Iredale, T. A Northern Australian volute. Proc. R. zool. Soc. N.S.W. 1954-55 1956: 76-77 1 fig.

Iredale, T. (1). History of New South Wales shells. Part I: Cook and his associates. Proc. R. zool. Soc. N.S.W. 1954-55 1956: 81-83.

- Ishida, S. Metabolic patterns in bivalves. VIII. Distribution of the guanine-deaminating activity in the soft parts of some marine bivalves. IX. Some aspects of nitrogen metabolism in marine bivalves, with special bolism in marine bivalves, with special reference to the mussel, Mytilus edulis. Bull. mar. biol. Sta. Asamushi 8 1: 9-32 5 figs.
- Ishida, S. & Tsuzuki, K. Metabolic patterns in bivalves. V. Distribution of the uricase activity in the bivalves, Meretrix meretrix lusoria and Venerupis phippinarum. VII. Some aspects of the purine metabolism in the clam. Meretrix meretrix lusoria (Gmelin). J. Coll. Sci. Chiba Univ. Nat. Sci. Scr. 1 4 1955: 267-272.
- Ishida, S. Tsuzuki, K. & Nitahara, I. Metabolic patterns in bivalves. X. Distribution of the uricase activity in the bivalves, Mactra sulcataria, Anadara inflata, Mytilus edulis, and certain other species. J. Coll. Sci. Chiba Univ. Nat. Sci. Scr. 21:82-864 figs.
- Itagaki, H. Anatomy of Oncomelania nosophora (Robson) (Gastropoda). Venus, Fukuyama 18 3 1955 pp. 161–168 pl. 1–2. [In English with Japanese summary.]
- Itagaki, H. Anatomy of Lymnaea (Fossaria) truncatula (Müller) from Japan with the comparison with the European material. Venus, Fukuyama 19 1: 25-38 pls. 2-5.
- Itoigawa, J. On the variation of Katelysia (Nipponomarcia) nakamurai Ikebe. Venus, Fukuyama 18 3 1955: 176–184 4 graphs. [In Japanese with English summary.]
- Itoigawa, J. (1) Molluscan fauna of the Tsuzuki group in Kyoto Prefecture, Japan. Mem. Coll. Sci. Univ. Kyoto 23B: 179-192 3 pls. 2 figs.
- Ivanov, A. V. Structure of the parasitic gastropods, Stiliferidae as the result of their mode of life. Trud. Leningrad Soc. Nat. 71 4 1952 86-140 48 figs. [In Russian.]
  - Ivanov, E. see Muravyev, I.
- Iwai, E. Notes on a rare species of Cranchiids squid. Bull. Jap. Soc. sci. Fish. 21: 1210-1214 2 pls.

- Iwai, E. (1) Descriptions on unidentified species of dibranchiate cephalopods. I. An oegopsiden squid belonging to the genus Architeuthis pp. 139–152 5 pls. II. A Cranchiidae squid of the genus Taonius pp. 153–162 3 pls. Sci. Rep. Whales Res. Inst. Tokyo 11.
- Iwanow, A. W. On the ways of penetration into host body as observed in parasitic Gastropoda. Trav. Soc. Nat. St. Pétersb. (Léningr.) 69 4 1947: 88-92. [English summary.]
- Iwanow, A. W. (1). On the degeneration in the structure of the larvae of parasitic Gastropoda, Trav. Soc. Nat. St. Pétersb. (Léningr.) 69 4 1947: 93-99 2 figs. [English summary.]
- Iwata, K. S. Activation of Mactra egg by modification of ionic composition of sea water. Biol. J. Okayama Univ. 1 3 1953: 177–184 2 figs.
- Iwata, K. S. (1). Mechanism of egg maturation in *Mytilus edulis*. Biol. J. Okayama Univ. 1 1-2 1952: 1-11 7 figs.
- Jacob, J. Allopolyplidy in the melaniid snails. Nature, Lond. 174 1954: 1061–1062 2 figs.
- Jacobson, M. K. Malacological notes from western Cuba. Nautilus, Philad. 70 1:1-3.
- Jacobson, M. K. (1) A new Macroceramus from Puerto Rico. Arch. Mollusk. 84 1/3 1955: 97–99.
  - Jacquemart, S. see Lelaup.
- Jacquemart, S. & Leloup, E. Observations écologiques sur une pelouse xérothermique. Le Thiers Pirard (Comblain-au-Pont). Mém. Inst. Sci. nat. Belg. No. 134: 1-99 3 pls. 17 figs.
- Jackel, S. Zur Kenntris de Molluskenfauna von Feldberg in Mecklenburg. Arch. Fr. Naturg. Mecklenburg 2: 273–280.
- Jaeckel, S. (1). Die Weichtiere (Mollusca) der Afghanistan-Expedition (1952 und 1953) J. Klapperichs. Mitt. zool. Mus. Berl. 32 337–353 10 figs.
- Jaeckel, S. (2). Einige Landschnecken vom Kamerun-Berg. Mitt. zool. Mus. Berl. 32: 354.
- Jaeckel, S. G. A. Süsswassermollusken aus spät-und postglazialen Ablagerungen im Schleswig-Holstein. Schr. naturw. Ver. Schl.-Holst. 28: 76–93.

Jaeckel, S. H. Die Wassermollusken der Nuke Niederung und des Raumes zwischen mittlerer Elbe und Warthe. Abb. Ber. Naturk. Vorgesch. Magdeburg 9 1955: 185–217 4 figs.

Jaeckel, S., Sen. & Meise, W. Über Laud und Susswasserschnecken Jugo: slaviens und Albanians. Mitt. hamburg. zool. mus. 54: 21-32 1 pl., 2 figs.

Jakl, H. L. Untersuchungen an Kleinmollusken, Laich, Weichtierschalen und Perlen. Mikrokosmos 45: 160-163 5 figs.

Jánský, L. Suchozemští měkkýši okolí vranovské přehrady. Čas. nár. Mus. Odd. Přírod. No. 125: 48–52. [German summary.]

Janus, H. Muscheln Kopflose Geschöpfe des Wassers. Kosmos, Stuttgart 50 1954: 110-114 2 pls. 7 figs.

Janz, G. J. & Carvalho, A. M. d. Subsidios para o conhecimento das bilharzioses em Angola. An. Inst. Med. trop. Lisbon 13: 597-613 2 maps. [English summary.]

J[enner], H. E. Mass mortality among molluscs off Lowestoft. Rep. Lowestoft Fld. Cl. No. 5 1950 : 152–153.

Jodot, P. see David, L.

Jodot, P. see Grangeon, P.

Jodot, P. Faunule malacologique du Lutétien continental de Sainte-Madeleine (Alpes-Maritimes) [short list of species.] Bull. Soc. géol. Fr. 6 3 1-3 1953: 59-65.

Johansson, J. (1). On the pallial gonoduct of Actaeon tornatilis (L.) and its significance for the phylogeny of the Euthyneura. Zool. Bidr. Uppsala 30: 223-232 1 pl. 3 text-figs.

Johansson, J. (2). On the anatomy of *Tympanotonus fuscatus* (L.) including a survey of the open pallial oviduc s of the Cerithiacea. Atlantic Report No. 4 Danish Exped. Tropical W. Africa 1945–46 1956: 149–166 text figs. & plates.

Johansson, J. (3). On the genital organs of some mesogastropods: Cerithium vulgatum Brug., Triphora perversa (L.) and Melanella (Eulima) intermedia (Cantr.) Contributions to the phylogeny of the pallial gonoducts of the Prosobranchia. Zool. Bidr. Uppsala 30: 1-24 pl. 1.

Johansson, J. (4). Garnault's duct and its significance for the phylogeny of the genital system of *Valvata*. Zool. Bidr. Uppsala 30: 457–464 2 pls. 4 text-figs.

Johansson, J. (5). Genital organs of two Alvania species, and a comparison with related families (Moll. Pros.). Ark. Zool. 9: 377–387 Pl. 1 text f. 1-4.

Johnson, C. G., Southwood, T. R. E. & Entwistle, H. A method of sampling arthropods and molluses from herbage by suction. Nature, Lond. 176 1955: 559.

Johnson, M. C. A living specimen of the East Coast geoduck from St. Augustine, Florida. Nautilus, Philad. 69 4: 121-123 pl. 8.

Johnson, R. I. The Types of Naiades (Mollusca: Unionidae) in the Museum of Comparative Zoology. Bull. Mus. comp. Zool. Harv. 115: 101-142 pl. 12.

Joly, M. see Jullien, A.

Joly, M. see Jullien, A. & others.

Joly, M. see Ripplinger, J.

Joly, T. see Tintant, H.

Jones, A. H. The paper nautilus. Bull. Conch. Sect. Auckl. Mus. 12: 20-21 1 fig.

Jones, N. S. The fauna and biomass of a muddy sand deposit off Port Erin, Isle of Man. J. Anim. Ecol. 25: 217-252.

Jørgensen, C. B. Quantitative aspects of filter feeding in invertebrates. Biol. Rev. 30 1955: 391-454.

Jove, J. A. Use of molluscoiides in the control of bilharsiasis in Venezuela. Equipment and methods of application. Bull. World Hlth Org. 14 4: 617-638.

Judd, W. W. Observations on a second colony of the land snail Cepaea nemoralis (L.) at London, Ontario with a consideration of the banding patterns in the two colonies. Canad. Fld. Nat. 69: 148-150.

Jullien, A. see Binet, L. & others.

Jullien, A., Cardot, J. & Rerat, D. Sur les durées de service du cocur isolé d'Helix pomatia conservé à basse température dans les milieux variés. Ann. Sci. Univ. Besançon (2) Zool. & Phys. No. 5: 23-61 12 figs.

- Jullien, A., Cardot, J., Ripplinger, J. & Claudey, P. Réactions inflammatoires provoquées expérimentalement dans le foie des Céphalopodes décapodes. C.R. Acad. Sci. Paris 243: 456-458.
- Jullien, A., Cardot, J. & Vieille-Cessay, C. Sur quelques particularités fonctionnelles du coeur isolé d'*Helix nemoralis*. C.R. Soc. Biol. Paris 150: 720-722.
- Jullien, A. & Ripplinger, J. (1). Recherches sur la répartition, dans la eveur de l'escargot (Helix pomatia), des fibres chronotropes et tonotropes négatives du nerf viscéral. C.R. Ass. franc. Av. Sci. 72 1953: 470-472.
- Jullien, A. & Ripplinger, J. (2). Recherches sur la structure du nerf visceral de l'escargot (*Helix pomatia*) et sur la distribution de ses fibres constatives à l'interieur du myocarde. Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954: 71-77 4 figs.
- Jullien, A. & Ripplinger, J. (3). Recherches sur la transmission humorale des influx cardio-inhibiteurs chez l'escargot. (Helix pomatia). Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954; 35-70 3 figs.
- Jullien, A. & Ripplinger, J. (4). De l'action des nerfs vagues on de type vagal sur le fonctionnement du coeur chez les mollusques et les poissons et ses rapports avec la production éventuelle d'un médiateur: l'acétylcholine. Bull. biol. 90 1: 33-105 17 figs.
- Jullien, A. & Ripplinger, J. (5). Sur un autagonisme atropine-calcium observé chez *Helix pomatia*. C.R. Soc. Biol. Paris 150: 1209-1211.
- Jullien, A., Ripplinger, J. & Cardot, J. (1). Altérations de l'activité du coeur d'Helix pomatia en transplantation hétéroplastique chez les Vertébris hétérothermes et effets de l'ion calcium sur son retourn l'automatisme. J. Physiol., Paris 47 1955: 206-208.
- Jullien, A., Ripplinger, J. & Cardot, J. (2). Variations de l'automatisme du coeur d'Helix pomatia en transplantations hétéroplastiques sur des vertebrés hétérothermes Actions subséquente des liquides physiologiques et des solutions à ions alcaline-terreux dominants. Ann. Sci. Univ. Besançon (2) Zool. & Phys. No. 4 1955: 59-94 12 figs.

- Jullien, A., Ripplinger, J. & Cardot, J. (3). Sur la production d'une substance équivalente à l'acétylcholine par le coeur d'Helix pomatia perfusé par voie interne et ses variations quantitatives d'après l'état fonctionnil de l'organe. C.R. Soc. Biol., Paris 150 1: 190-192.
- Jullien, A., Ripplinger, J. & Cardot, J, (4). Sur les rapports entre la production d'acétylcholine ou d'une substance équivalente par la coeur isolé d'*Helix pomatia* perfusé par voie interne et lu fréquence de l'organe. J. Physiol., Paris 48: 581-584 2 figs.
- Jullien, A., Ripplinger, J. & Guyon, M. (1). Effets, sur l'automatisme du coeur in situ d'Helix pomatia des variations experimentales de la tension de l'organe. Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954: 85-90.
- Jullien, A., Ripplinger, J. & Guyon, M. (2). Sur les variations de l'automatisme du coeur chez Helix pomatia succédant à des cardiactomés partielles ou subtotales à des remplacements de l'organe et à des changements de tension exercés sur le myocarde. Ann. Sci. Univ. Besançon (2) Zool, & Phys. No. 4 1955: 95-144 9 figs.
- Jullien, A., Ripplinger, J. & Joly, M. (1). L'automatisme du coeur de l'escargot (Helix pomatia) et conditionné par la présence de substances excitomotrices: Leur rôle physiologique. C.R. Acad. Sci. Paris 240 8 1955; 911-913.
- Jullien, A., Ripplinger, J. & Joly, M. (2). Influence de la composition des solutions de perfusion sur l'evolution de l'automatisme ventrialiaire en fonction du temps chez Helix pomatia. Mise en évidence de substances excitomotricés cardiaques. J. Physiol., Paris 47 1955: 200-205 1 fig.
- Jullien, A., Ripplinger, J. R. & Joly, M. (3). Etude comparative des effets, sur le coeur d'Helix pomatia, des extraits et des diffusats de nerf visceral. C.R. Soc. Biol., Paris 149 9-10 1955-964-966.
- Jullien, A., Ripplinger, J. & Joly, M. (4). Action sur le coeur d'Helix pomatia perfusé en situ, de diffusats et d'extracts de différents nerfs. J. Physiol., Paris 42:584-588.

Jullien, A., Ripplinger, J. & Joly, M. (5). Chez l'escargot (Helix pomatia), substance vagale et substance excitomotrice représentent le même produit. Comparaison de la transmission humorale de l'inhibition cardiaque chez l'escargot et chez la grenouille. C.R. Soc. Biol., Paris 150: 192-196.

Jullien, A. Ripplinger, A. & Joly, M. (6). Les échanges ioniques entre les imbeux de perfusion et le coeur d'Helix pomatia, sans les conditions normales et au cours de l'action du neuf viscéral. Rôle des ions sur la substance excitomotrice. Ann. Sci. Univ. Becançon (2) Zool. & Phys. No. 5: 67-123 15 figs.

Jullien, A., Ripplinger, J. & Mercier, J. Sur la formation et la diffusion d'une substance equivalente à l'acetylcholiné par le coeur isole d'Helix pomatia, en fonction de la composition et de la concentration du milieu de perfusion, de la durce, de la temperature et de l'état d'activité on de repos de l'organe. Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954: 91-124 18 figs.

Jullien, A., Ripplinger, J. & Moreau, R. Effets de l'histaminé et des antilistaminuques de synthése sur le comportement des Céphalopodes et de quelques poissons marins. Bull. Soc. Hist. nat. Doubs No. 58: 185–193.

Jutting, W. S. S. v. B. Systematic studies on the non-marine Mollusca of the Indo-Australian archipelago. V. Critical revision of the Javanese freshwater gastropods. Treubia 23: 259-477 135 figs.

Kaas, P. Notes on *Loricata*. Basteria, Lisse 20 4/5: 106 107.

Kajiyama, H. On the distribution of freshwater shells. Venus, Fukuyama 19 1: 49-54 text map. [In Japanese.]

Kakhadze, I. R. The problem on a border between Aalen and Bias in the valley of the river Kuban. Bull. Acad. Sci. U.R.S.S. geol. 3 1955: 119 122 fig. [In Russian.]

Kalabis, V. Zum Vorkommen von Cerithien und Melanopsis martiniana in der Umgebung von Olomonc. Sborn. Sluko 2: 145–151. [German summary.]

Kamada, Y. Japelion yabei a new Miocene Gastropod from the Jobarn Coal-field Japan. Venus, Fukuyama 18 3 1955: 185–188 pl. 3. Kamat, D. N. Intracellular digestion in lamellibranchs. Proc. Indian sci. Congr. Ass. 43rd 3:302-303, [Abstract.]

[1956]

Kamat, D. N. (1). Phosphatases in Lamellidens corrianus (Lea): Histochemical demonstration in the digestive diverticula and the amoebocytes. J. Univ. Bombay N.S. 24B 5: 25-32 5 figs.

Kanakoff, G. P. Two new species of Nassarius from the Pliocene of Los Angeles County, California. Bull. S. Calif. Acad. Sci. 55 2: 110-114 2 pls.

Kanamaru, T. A history of the Japanese Conchology (33). Venus, Fukuyama 18 3 1955: 198-203. [In Japanese.]

Kanamaru, T. (1). A History of the Japanese Conchology (35). Venus, Fukuyama 19 1:65-72. [In Japanese.]

Kanamaru, T. (2). A history of the Japanese Conchology (36). Venus, Fukuyama, 19 2: 124–131. [In Japanese.]

Kanis, J. Geology of the eastern zone of the Sierra del Brezo (Palencia-Spain). Leid. geol. Meded. 21: 375–445 3 pls. figs.

Kanno, S. On some new Glycymerid Molluscs from Japan. Trans palaeont. Soc. Japan 24: 267-272 1 pl.

Kan-o, A. A profile of a neolithic shell mound at Sonno-machi, Chibashi. J. Coll. Sci. Chiba Univ. Nat. Sci. Scr. 1 4 1955: 300–303 3 figs.

Kapler, O. Malakozoologické poznámky z prameniště Moravice. Prirod. Sb. Ostravsk. Kraje 17: 597–598.

Karlin, E. J. Notes on the ecology of Zonitoides arboreus (Say), Opeas pumilum (Pfeiffer), and Lamellaxis gracilis (Hutton) in green houses. Amer. Midl. Nat. 55: 121-125.

Karlov, N. W. The age of Mandrikovsk fauna. C.R. Acad. Sci. URSS N.S. 94 1 1954: 125–127. [In Russian.]

Karlov, N. W. (1). The finding of fossil shells of Caspian mollusca in river-bed deposits of lower Dneper and Ingultz. C.R. Acad. Sci. URSS N.S. 95 6 1954:1283–1285. [In Russian.]

Karlov, N. W. & Gryaznov, V. I. [The age and genesis of homogenous layer of Tertiary sands in the region of the town Zaporozhe. C.R. Acad. Sci. URSS N.S. 94 5 1954: 931-932 fig. [In Russian.]

Karpevich, A. F. Some data on the development of bivalve molluscs. Zool. Zh. 34 1 1955: 46-67 Pl. Sdingrs. [In Russian.]

Kasimova, N. M. see Khaim, V. E. Kašiák, V. Vorläufige Mitteilung über das Auftreten von Fischparasiten in Gewässen der Slowakei. Biológica, Bratislava 11 10 1956: 624-635. [German summary.]

Kathuria, J. B., Rao, S. R. & Hiregaudar, L. S. Some observations on the bionomics of *Indoplanorbis exustus* Desm. (Gastropoda). Proc. Indian sci. Congr. Ass. 43rd 3: 355. [Abstract.]

Katto, J. see Ozaki, H.

Kauth, H. see Adlung, K.

Kawaumura, R see Kuroda, T.

Keen, A. M. Ventricolaria and Similivenes insolida, new names in the Pelecypod family Venerüdae. J. Paleont. 28 1954: 217-218.

Keen, A. M. (1). Comment on a paper by David Nicol [arctic marine pelecypod fauna.] Nautilus, Philad. 69 4: 139-140.

Kellum, L. B. Cretaceous invertebrates of the Aurora Limestone. Pap. Mich. Acad. Sci. 41: 205-231 7 pls. map.

Kelly, C. B. & Arcisz, W. Bacterial control of oysters during processing and marketing. Public Hith. Rep. Wash. 69 1954: 716–720 2 figs.

Kelly, C. B. & Arcisz, W. (1). Survival of enteric organisms in Shellfish. Public Hlth. Rep. Wash. 69 1954: 1205–1209

Kendall, S. B. Liverflukes and snails in Asia. Proc. Linn. Soc. Lond. 166 1-2: 6.

Kennedy, G. Y. & Vevers, H. G. Porphyrin pigments in the tectibranch mollusc Akera bullata O.F. Müller. J. Mar. biol. Ass. U.K. 35: 35-39.

Kenny, R. see Endean & other.

Kerkut, G. H. see Hughes, G. M.

Kerkut, G. A. & Taylor, B. J. R. The sensitivity of the pedal ganglion of the slug to osmotic pressure changes. J. exp. Biol. 33: 493-501 7 figs.

Kerkut, G. A. & Taylor, B. J. R. (1). Effect of temperature on the spontaneous activity from the isolated ganglia of the slug, cockroach and crayfish. Nature, Lond, 178: 426.

Kerney, M. P. Note on the fauna of an early Holocene Tufa at Wateringbury, Kent. Proc. Geol. Ass. Lond. 66: 293-296.

Kershaw, R. C. Victorian fluviatile and lacustrine Mollusca. Part 1. Vict. Nat. Melb. 73 4:51-55.

Keyl, H.-G. Beobachtungen über die J-Meiose der Muschel Sphaerium corneum. Chromosoma 8: 12-17 8 figs.

Keynes, R. D. see Hodgkin, A. L.

Khabakov, A. V. Ancient sinkings and dislocations of the Artinsk layers in the Central Ural. C.R. Acad. Sci. URSS 61 6 1948: 1091-1094 3 figs. [In Russian.]

Khain, V. E., Shardanov, A. N., Kasimova, N. M. The stratigraphy of the middle Jurassic of the South Eastern Caucasus. C.R. Acad. Sci. URSS 100 5 1955: 965-968. [In Russian.]

Khristenko, N. G. On the distribution of *Bithynia leachi* Schepp. in the lakes of the Bol'sheosera-group in the Krasnoyarsk Territory. Zool. Zh. 35: 1583-1584.

Kiefer, F. Ein neuer Fundort von Lartetia rhenana R. Lais. Beitr. naturk. Forsch. SudwDtsch. 15: 125 1 fig.

Kikuchi, H. Ecological survey of aquatic animals in the stream Yokoshibetsu, Sapporo. J. Fac. Sci. Hokkaido Univ. (Zool.) 12: 401-411 f. 1-6.

Kilias, R. Über die Kieme der Teichmuschel (*Anodonta* Lam.). (Ein Beitrag zur Anatomie). Mitt. zool. Mus. Berl. 32: 151–174 17 figs.

Kimura, T. The Torinosu group and the Torinosu limestone in the Togano and Go Basins, Kochi Prefecture. J. Geol. Soc. Japan 26 No. 732: 515-526 7 figs.

Kinoshita, see Kuroda, T.

Kiparisova, L. D. & Popov, Y. N. Segmentation of the lower part of the Triassic system into layers. C.R. Acad. Sci. USSR 109 4: 842-845. [In Russian.]

Kira, T. A remarkable ecological form of *Apollon pusillus*. Venus, Fukuyama 19 2: 122-123 text figs.

Kirpichnikova, E. S. Ring structures in muscle fibres of *Anodonta* heart. C.R. Acad. Sci. URSS 104 6 1955: 919–921 fig. [In Russian.]

Kisch, B.S. L. de Folin (1817–1896). J. Conchyliol. Paris **96** 4: 138–154 portrait.

Kisch, B. S. (1) Les mollusques du Bassin filtrant de l'Aquarium de Biarritz. Bull. Centre Ét. Rech. Sci. Biarritz 11:41-497 figs.

Kitamura, T. see Asano, M.

Kittel, R. Untersuchungen über den Geruchs-und Geschmackssin bei den Gattungen Arion und Limax (Mollusca: Pulmonata). Zool. Anz. 157: 185–195.

Klenova, M. V. Geological structure of Apsheron sill of the Caspian Sea. C.R. Acad. Sci. URSS N.S. 94 2 1954: 311-314. [In Russian.]

Kline, G. F. Notes on the stinging operation of *Conus*. Nautilus, Philad. 69 3 76-78.

Klock, J. W. A field technique for quantitative estimation of the molluscicide sodium pentachlorophenate based on fish mortality rates. Amer. J. trop. med. Hyg. 5: 286–289 1 fig.

Klock, J. W. (1). An automatic molluscicide dispenser for use in flowing water. Bull. World Hlth. Org. 14 4: 639-643 3 figs.

Knight-Jones, E. W. Relations between metachronism and the direction of ciliary beat in Metazoa. [Moll. p. 511-514.] Quart. J. micr. Sci. 95 1954: 503-521 10 figs.

Knipper, H. Neue Achatiniden aus Ostafrika (Moll. Gastrop. Stylomn.) Veröff. Überseemus. Bremen 2A: 341– 356 1 pl. 8 figs.

Knipper, H & Meyer, K. O. Biologische und anatomische Betrachtungen an ostafrikanischen Ellobiiden. Zool. Jb. (Syst.) 84: 99-112 1 pl. 13 figs. [English summary.]

Kobayashi, H. Cytochemische Studien über die Eier der Pelecypoden. J. Sci. Hiroshima Univ. (Zool.) 16: 1-51 5 pls. Kobayashi, T. (1). An interesting new form of the Aturidae from the Palaeogene of Northern Kynshu. Trans. palaeont. Soc. Japan 23: 243-245 4 figs.

Kobayashi, T. (2). Some Jurassic Trigonias from Central and West Japan. Studies on Jurassic Trigonias in Japan, Part V. Jap. J. Geol. & Geogr. 27 1: 1–7 1 pl.

Kobayashi, T. (3). On the dentition of Trigonioides and its relation to similar Pelecypod genera. Japan J. Geol. & Geogr. 27 1: 79-92 1 pl. 3 figs.

Kobayashi, S. see Mawatari, S.

Kobayashi, T. & Amano, M. On the pseudoquadrate trigonians, *Stein-mannella*, in the Indo-Pacific province. Jap. J. Geol. Geog. 26 1955.

Kobayashi, T. & Ichikawa, K. Tosapecten gen. nov. and other Upper Triassic Pectinidae from the Sakawa Basin, in Shikoku, Japan. Jap. J. Geol. & Geogr, 21 1949: 163–174 1 pl.

Kobayashi, T. & Ichikawa, K. (1). Myophoria and other Upper Triassic Pelecypods from the Sakawa Basin in Shikoku, Japan. Jap. J. Geol. & Geogr. 21 1-4 1949: 177-190 1 pl.

Kobayashi, T. & Ichikawa, K. (2). Late Triassic "Pseudomonotis" from the Sakawa Basin in Shikoku, Japan. Jap. J. Geol. & Geogr. 21 1949: 245-264 1 pl.

Kohn, A. J. Feeding in *Conus striatus* and *C. catus*, Proc. Hawaii. Acad. Sci. No. 31: 20.

Koltuipin, S. N. Ob odnom sluchae urodstua rostra belemnitellui. Annu. Soc. paléont. russe 14 1953: 207-212.

Komárek, K. Větrání a poruchy Vápence. [Weathering and limestone treatment.] Čas. nár. Mus. 125 2: 152-155 1 fig.

Kon, S. K. see Fisher.

Kondo, Y. First Helix aspersa in Hawaii, Nautilus, Philad. 69 4: 141-142.

Kondo, Y. (1). Henry Edward Crampton 1875–1956. Nautilus, Philad. 70 1: 31–34.

Kondo, Y. (2). Second *Helix aspersa* in Hawaii and data on carnivorous snails. Nautilus, Philad. **70** 2: 71–72.

Konietzko, B. see Leloup, E.

Korobkov, I. A. Pectinidae from the Paleogene of Middle Asia. Trud. Leningrad. Soc. Nat. 68 2 1951: 64-102 4 pls. [In Russian.]

Korringa, P. Oyster culture and biological productivity. Rapp. Cons. Explor. Mer 140 3: 30-31.

Korringa, P. (1). Oyster culture in South Africa. Invest. Rep. Div. Fish. S. Afr. No. 20 1–85 23 pls.

Koshelkina, Z. V. & Tuchkov, I. I. The age of the Ancella horizon in Prever Khoyansk ridge bend. C.R. Acad. Sci. URSS 102 4 1955: 801-803. [In Russian.]

. Kovalevski, B. A. see Viktorona, R. E.

Kozlovskaya, L. S. Inhabitance of fish-water molluses belonging to the genus *Pisidium* in bogged silvan soils. C.R. Acad. Sci. URSS N.S. 95 6 1954: 1321-1323. [In Russian.]

Kozlovskaya, L. S. (1). Absence of Prosobranchia in the deposits of the lakes in the basin of Tobol and Ishim. C.R. Acad. Sci. URSS N.S. 98 4 1954: 649-651 map. [In Russian.]

Krach, W & Nowak, W. Le Miocène des environs d'Andrychow. Rocznik Ann. Soc. Géol. Pologne 25 1 1955 pp. 9-54 plates and 5 text figs.

Kraemer, A. Von Sagittoceras costulatum Kobald im Sauerland. Abh. Landesmus. Westf. 11 1 1940: 54-55.

Kraus, O. see Zilch, A.

Krayukhin, B. V. see Bodrova, N.

Krijgsman, B. J. & Divaris, G. A. Contractile and pacemaker mechanisms of the heart of molluscs. Biol. Rev. 30 1955: 1-39.

Krimgoltz, G. Y. Some cephalopods from the Jurassic deposits of Transcaucasus. Trud. Leningr. Soc. Nat. 68 2 1951: 26-38 Pl. [In Russian.]

Kristensen, I. De Artemisschelp. Levende Nat. **59**: 82–84 1 fig.

Krömmelbein, K. Die ersten marmen Fossilien (Trigoniidae, Lamellibr.) aus der Cayetano-Formation West-Cubas. Senckenbergiana leth. 37: 331-335 1 pl. Kudrin, L. N. Venus sobieskii Hilb. i Venus sobieskii Hilb. var. media Sok. iz miotzenovuikh otlozheniī yughozapadnoi okrainui russkoi platformui. Geol. Sbornik 2-3: 188-193 6 pls. [In Russian.]

Kudrin, L. N. (1). The age of the sand, clay, coal deposits in the territory of the South-western edge of the Russian Platform. C.R. Acad. Sci. URSS 101 3 1955: 535-536. [In Russian.]

Kudrin, L. N. (2). The absence of paleogene deposits in the South-western ridge of the Russian Platform. C.R. Acad. Sci. URSS 102 2 1955: 347-349. [In Russian.]

Kühnelt, W. Lebensformen und Entwicklungsnichtungen der Muscheln. Verh. zool.-bot. Ges. Wien. 96 16-41 15 figs.

Kuiper, J. G. J. Helicodiscus (Hebetodiscus) singleyanus inermis H. B. Baker neu für die europäiscke Fauna. Arch. Molluskenk. 85 4/6: 163-169 text figs.

Kulikov, M. V. [New data on biostratigraphical segmentation of Permian deposits of Northern Verkhoyan]. C.R. Acad. Sci. URSS. 101 6 1955: 1101–1103. [In Russian].

Kummel, B. Jurassic nautiloids from western North America. J. Paleont. 28 1954: 320–324 pls. 34–35.

Kummel, B. (1). Post-Triassic Nautiloid Genera. Bull. Mus. comp. Zool. Harv. 114 321-494, 28 pls.

Kummel, B. & Decker, J. M. Lower Turonian ammonites from Texas and Mexico. J. Paleont. 28 1954: 310-319 pls. 30-33 text-figs. 1-10.

Kuntz, R. E. Evaluation of sodiumpentachlorophenate as a molluscicide in Egypt. Amer. J. trop. Med. Hyg. 5 2:274-285.

Kuo, Jen-Chiang & Pei, Hung-kang. Studies on the periodic changes of the reproductive glands of the oncomelaniid snails, an intermediate host of Schistosoma japonicum. Acta zool. sinica 8:161-176 16 figs. [English summary].

Kuroda, T. Land Shells fauna of Yakeu Island. In Japanese. Venus, Fukuyama 18 3 1955: 145–157. Kuroda, T (1). New speices of the Conidae (Gastropoda) from Japan. Venus, Fukuyama 19 1: 1-16, text-figs. 1-10 pl. 1.

Kuroda, T. (2). A new *Pleurotomaric* from Japan with a note on a specimen of *P. rumphii* Schipman collected from Taiwan. Venus, Fukuyama 18 4 1955: 211-221 pls. 8, 9.

Kuroda, T. and Kawamura, R. On a new species of *Lischkeia* (Gastropoda): *L. undosa* from Japan. Venus, Fukuyama, 19 1:16-19 text-figs. 1, 2.

Kuroda, T. & Kinoshita, T. A remarkable new Buccinid, *Volutharpa ainos* from Hokkaido. Venus, Fukuyama 19 1: 19–22 text-fig.

Kuzina, I. See Alekseichik, S.

Kuznetsov, V. V. Some regularities of the intraspecific morphological and biological variability on the example of the marine invertebrates. Zool. Zh. 35 1118-1130 4 figs. [English summary].

Kvaliashvili, G. A. Presence of representatives Kulnitzk horizon fauna in guriïsk deposits of Guria. C.R. Acad. Sci. URSS N.S. 96 1 1954:161-162. [In Russian.]

Lacy, D. & Horne, R. A cytological study of the neurones of *Patella vulgata* by light and electron microscopy. Nature, Lond. 178 976-978.

Lacy, D. & Rogers, G. E. Recent observations by light and electron microscopy on the cytoplasmic inclusions of the neurones of *Patella vulgata*. J. roy. micr. Soc. (3) 75 172–175 1 pl.

Laidlaw, F. F. Baiaplecta n. nom. Arch. Molluskenk. 85 1/3 p. 83.

Lal, M. B. & Premvati. Studies in histopathology—Changes induced by a larval monostome in the digestive gland of the snail, Melanoides tuberculatus (Müller). Proc. Indian Acad. Sci. 42B 6 1955: 293-299 5 figs.

Lambert, R. See Maubeuge, L.

Lamesch, M. Les Limacidés et les Arionidés (Mollusques) du Grand-Duché de Luxembourg. Bull. Soc. Nat. luxemb. N.S. 47 1953: 236–286 18 figs. Lamont, A. New lamellibranchs from the Gutterford Burn Flagstones (Gala-Tarannon) of the Pentland Hills, near Edinburgh. Proc. roy. Soc. Edinb. 65B 1958: 271-284 1 pl.

[1956]

Lamy, L. Les bilharsiosis et les moyens nouveaux de la lutte antibilharsienne. Nature, Paris No. 3236 1954: 472–476 12 figs.

Lang, W. D. The Coinstone of the Charmouth Lias. Proc. Dorset nat. Hist. & Archaeol. Soc. 67 1946: 145.

Landman, M. A. Statistics on a colony of *Cepaea nemoralis*. Nautilus, Philad. 69 3 104-106.

Lane, C. E. Recent biological studies on Torido—a marine wood boring molluse. Sci. Monthly 80 1955: 286– 292 4 figs.

Lapkin, I. Y. Lime-dolomite layer of the Donetz Permian. Bull. Soc. Nat. Moscou geol. 29 4 1954: 37-50 3 figs. [In Russian].

Lapparent, A. L. de. Sur la stratigraphie du Tamesna (Sahara méridional) [short list in text.] C.R. Acad. Sci. Paris 236 1953: 2255–2257.

Laptinskaya, E. S. See Vorontzev, N.

Larcher, C. See Tintant, H.

La Roque, A. Variation of carinae in Valvata tricarinata. Nautilus, Philad. 701:13-14.

La Rocque, A. & Conley, J. F. Two Pleistocene Molluscan faunules from Hunter's Run, Fairfield County, Ohio. Ohio J. Sci. 56 325–328 1 fig.

La Rocque, A. & Marple, M. F. Ohio Fossils. Bull-geol. Surv. Ohio 54 1955: 1-152 figs.

Larionova, E. N. & Safonova, T. P. Coal deposits at the town Sovetska at Vyatke elevation. Bull. Acad. Sci. Moscou Geol. 6 1955: 29-41 map. [In Russian.]

Laseron, C. F. The families Rissoinidae and Rissoidae (Mollusca) from the Solanderian and Dampierian zoogeographical provinces. Aust. J. mar. & freshw. Res. 7 384-487 228 figs.

Laseron, C. F. (1). New South Wales mussels. A taxonomic review of the Family Mytilidae from the Peronian zoogeographical Province. Aust. Zool. 12 263–283.

**Laseron C. F. & J.** A new genus of the Nassariidae. Proc. R. zool. Soc. N.S.W. **1954**–**55** 1956: 71–72.

Laursen, D., The genus *Ianthina*. Int. Congr. Zool. 14 1953 (1956): 510

Laursen, D. (1). Emerged Pleistocene marine deposits of Peary Land (North Greenland). Medd. Grønland 127 5 1954: 2nd pag. 1-26 1 pl. 3 figs. map.

Laviolette, P. Role de la gonade dans la maturation glandulaire du tractus génital chez quelques gastéropodes. Ann. Sci. nat. (Zool.) (11) 18 171-173.

Laviolette, P. (1). Regeneration germinale après castration chez un mollusque gastropode pulmone Arion rufus L. Int. Congr. Zool. 14 1953 (1956): 233–235.

Laviolette, P. (2). Role de la gonade dans le déterminisme humoral de la maturité glandulaire du tractus génital chez quelques gastéropodes Arionidae et Limacidae. Bull. biol. 88 3 1954: 310-332 1 pl. 2 figs.

Laviolette, P. (3). Étude cytologique et expérimentale de le régénération germinale après castration chez Arion rufus L. (Gasteropode Pulmoné). Année biol. (3) 30 11-12 1954: 481-492.

Lavoie, M. E. How sea-stars open bivalves. Biol. Bull. Woods Hole 111 114–122 4 figs.

Lavrov, V. V. On two stratigraphic schemes of the continental Tertiary series for the Steppes of Kazakhstan and Western Siberia Bull. Soc. Nat. Moscow Geol. 31 1: 3-19. [In Russian.]

Leanza, A. F. & Castellaro, H. A. Algunos fosiles Cretacicos de Chile. Rev. Asoc. Geol. Argent. 10 3 1955: 179-212 3 pls.

Leckwijck, W. V. See Ancion, C.

Leclercq, J. Invertébrés, surtout mollusques terrestres, cherchant abri sous les schistes houillers aufried des terriés de charbonnages du Pays de Herve (Liège). Nat. belges 35 9-10 1954:159-160.

Le Dantec, J. Observations sur un essai d'elevage d'huitres portugaises en caisses ostreophiles. Rev. Trav. Inst. Pêches marit. 20 171-182 fig. 1-4.

Lefevere, S., Leloup, E. & Meel, L. V. Observations biologiques dans le Port d'Ostende. Mém. Inst. Sci. nat. Belg. No. 133 1-157 3 pls.

Lellák, J. The Chironomidae and other bottom fauna of some stagnant waters in the Central Labe (Elbe) region. Rozpr. české Akad. 63 8-9 1953: 69-142 figs. [English summary.]

**Leloup, E.** A propos les amphineures *Chaetopleura papilio* (Spengler, 1797) et *Chaetopleura apiculata* (Say, 1830). Bull. Inst. Sci. nat. Belg. **32** No. 5 1-6 5 figs.

Leloup, E. (1). See Jacquemart, S.

Leloup, E. (2). See Leferve & Meel.

Leloup, E. & Konietzko, B. Recherches biologiques sur les eaux Saumâtres du Bas-Escaut. Mém. Inst. Sci. nat. Belg. No. 132 1–100 5 pls. 26 figs.

Leloup, E., Van Meel, L. & Jacquemart, S. Recherches hydrobiologiques sur trois mares d'eau douce des environs de Liège. Mem. Inst. Sci. nat. Belge. No. 131 1954: 1-145 4 pls. 18 figs.

Lemche, H. The anatomy and histology of *Cylichna* (Gastropoda Tectibranchia). Spolia zool. Mus. Hauniens. 16 1-.

Lemoine, M., Ricour, J. et Trümpy, R. Données paléontologiques nouvelles sur le gisement triasique de Cervières (Hautea Alpes). C.R. Soc. géol. Fr. 12 1952: 240-242.

Le Roux, P. R., etc. Vectors of human belharzicus in Africa. Trans. R. Soc. trop. Med. Hyg. 49 4 1955: 292-294.

Leroux, S. Phytoplaneton et contenus stomacaux d'huitres portugaises, (Gryphaea angulata Link.) dans le bassin D'Arcachon. Rev. Trav. Inst. Pêches marit., 20 163-170 f. 1-9.

Letaconnoux, R. Compte rendu d'essais pour la protection des bois contre les tarets. Rev. Trav. Inst. Pêches marit. 20 203-214 f. 1-4.

Letaconnoux, R. & Audouin, J. Contribution ou l'etude du pétoncle (*Chlamys varia L.*). Rev. Trav. Inst. Pêches marit. 20: 133-155 f. 1-12.

Letaconnoux, R. See Audouin, J.

**Lewowicki, S.** Stratigraphic position of Clymenia limestone from Dzikowiec Ktodzki (Lower Silesia). Przeglad Geologiczny 5:216–221 figs.

Li, Kie-min. See Tchang, Si.

Lipparini, T., Malatesta, A., Nicosia, M. L. & Valdinucci, A. Contributi alla conoscenza delle faune neogeniche e quaternarie della Sicilia.—V. Pliocene e Quaternario del Capo Milazzo in Sicilia. Boll. Uff. geol. Ital. 77 1955: 579-604 13 figs.

(Litvinenko, A. U.). [Zoomorphosis from Kimmeriisk deposits]. C.R. Acad. Sci. URSS. 102 5 1955:1001-1004 4 figs. [In Russian.]

Liverovskaya, E. On a Cretaceous Fauna from the West Coast of Sakhalin (north of the village Khoy). Annu. Soc. paléont. russe 12 1945: 93– 100 1 pl. [English summary p. 99.]

Liverovskaya, E. V. See Osipova,

Llabador, F. Revision des Néritines vivantes et fossiles du Nord-Ouest de l'Afrique. J. Conchyliol. Paris 96 3: 85-114.

Lloyd-Evans, L. Ena montana (Draparnaud) in Hertfordshire. J. Conch. London 24 3 102.

Lloyd-Evans, L. (1). Acroloxus lacustris (L.) in Merionethshire. J. Conch. Lond. 24 3 104.

Logie, R. R. Mortalités chez les huîtres des provinces maritimes dans le passé et présentement. Circ. Atlantic biol. Sta. No. 27 1–15.

Loosanoff, V. L. On utilization of salt water ponds for shellfish culture, Ecology 37 614-616.

Loosanoff, V. L. & Nomejko, C. A. Relative intensity of oyster setting in different years in the same areas of Long Island Sound. Biol. Bull. Woods Hole 111 387-392 1 fig.

Loosjès, F. E. Eine Clausilien-Ausbente von Kreta. S.B. öst. Akad. Wiss. Abt. I 164 1955: 855–885 5 figs.

Loosjès, F. E. (1). Clausiliidae (Gastropoda, Pulmonata) from Western New Guinea. Zool. Meded. 34 227–230, 2 figs.

Loosjès, F. E. (2), Euphaedusa cumingiana recondita (Sykes) (Moll., Styllon.) auf Flores. Verh. naturf. Ger. Basel 67 1 53-54 2 figs. Lopes, H. de S. Sôbre Pomacea canaliculata (Lamarck, 1822) (Mesogastropoda, Architaenioglossa, Mollusca). Rev. bras. Biol. 16 535–542 46 figs.

Lopes, H. de S. (1). Sôbre *Pomacea lineata* (Spix, 1827) (Mesogastropoda, Architaenioglossa, Mollusca). Rev. bras. Biol. 16 375–380 17 figs.

Lopes, H. de S. & Alvarenga, M. Contribuição ao conhecimento dos Moluscos da ilha Fernando de Noronha-Brasil. Bol. Inst. oceanogr. S. Paulo 6 1–2 1955: 157–196 4 pls.

Lopez, A. O. de Z. y. Observaciones anatómicas y posición sistemática de varios helicidos españoles. Bol. Soc. esp. Hist. nat. Sci. Biol. 54 35-61 30 figs.

Lopez, A. O. d. Z. & Rocandio, A. O. de Z. Contribuciones al conocimento de la fauna malacológica terrestre de la isla de Fernando Póo. Bol. Soc. esp. Hist. nat. Biol. 53 75–140 30 figs.

**López-Benito, M.** Chemical content of scallops (*Pecten jacobeus*). Rapp. Cons. Explor. Mer **140** 3:36–37.

Lörcher, E. Ein unerwarteter Ammoniten-Fund in der *Spinatus*-Zone (Domérien). Neues Jb. Min, Geol. Paläont. 97 1953: 51-56 pl. 6 1 text-fig.

Lowenstam, H. A. Factors affecting the Aragonite: Calcite ratios in Carbonate-secreting marine organisms. J. Geol. 62 1954: 284-322 text-figs. 1-15.

Lowenstam, H. A. See Epstein, G.

Lowy, J. The lamellibranch muscle. Nature, Lond. 176 1955: 345-346 3 figs.

Lowy, J. See Abbott, B. C.

Lowy, J. See Bowden, J.

Lowy, J. See Hoyle, G.

Ložek, V. Měkkyši vrchu Svince a "Tropického údolí" u Nového Jičina. Prirod. Sb. Ostravsk. Kraje 17 596–597. [German summary.]

Ložek, V. (1). Die Weichtiere des Zabó-Tales im Slowakischen Erzgebirge. Biológica, Bratislava 11 8 472– 479 2 figs. [German summary].

Ložek, V. (2). Key to Czechoslovakian Mollusca. Bratislava (Vydav. Slov. Akad.) pp. 437 42 pls. (2 col.) 69 figs.

- Ložek, V. (3). Malakozologische Neuigkeiten aus Tschochoslowakei III. Čas. nár. Mus. 125 2 142–150 7 figs. [German summary].
- Ložek, V. (4). Molluscs in the environment of Lubná near Rakovník (Western Bohemia). Čas. nár. Mus. 124 1955: 86-88. [English summary.]
- Ložek, V. (5). Molluscs in the environment of Skalsko. Čas. nar. Mus. 124 1955: 82-85.
- Lubet, P. Cycle neurosécrétoire chez Chlamys varia L. et Mytilus edulis L. (Mollusques lamellibranches). C.R. Acad. Sci. Paris 241 1955: 119–121 4 figs.
- Lubet, P. (1). Le determinisme de la ponte chez les lamellibranches (*Mytilus* edulis L.) Intervention des gauglions nerveux. C.R. Acad. Sci. Paris 241 1955: 254-256.
- Lubet, P. (2). Effets de l'ablation des centres nerveux sur l'émission des gamètes chez Mytilus edulis L. et Chlamys varia L. Ann. Sci. nat. (Zool.) (11) 18, 175–183, 6 figs.
  - Lucas, G. See Arkell, W. J.
- Ludbrook, N. H. The molluscan fauna of the Pliocene strata underlying the Adelaide Plains Part III—Scaphopoda, Polyplacophora, Gastropoda (Haliotidae to Tornidae). Trans. roy. Soc. S. Aust. 79 1-36 2 pls.
- Ludbrook, N. H. (1). Occurrence of Crespinina kingscotensis Wade. Aust. J. Sci. 18 pp. 164, 165.
- Lundgren, G. Landmollusker från det Jämtländsche Silurområder och augränsande trakter. Göteborgs Vetensk. Samh. Handl. (6) 6B 10 1955: 1– 44 map.
- Luppov, N. P. A Discovery of *Pulchellia* in North-Western Caucasus. Annu. Soc. paléont. russe 12 1945: 133–137 fig. [English summary p. 137.]
- Luppov, N. P. (1). Novuiï vid roda *Cicatrites* iz verkhneaptskikh otlozheniï severnogho Kavkaza. Annu. Soc. paléont. russe 13 1949:71-74 2 figs.
- Lüttig, G. Die Mollusken des Interglazials von Elze. [Lists]. Paläont. Z., 27 1-2 1953: 67-84 text-figs. 1-7.

- Lyashenko, G. P. New date on systematics of Tentaculitida, Stylio-limda and Novakiida. Bull. Soc. Nat. Moscou geol. 30 3 1955: 94-95. [In Russian.]
- Macarovici, N. Sur la fauna fossile du bassin Brad-Baia de Criş (rayon de Brad). An. Sti. Univ. "A. I. Cuza" lasi (2) 2 2 331-337 6 figs. [French summary.]
- Macarovici, N. (1). Sur la faune Silurienne du fondement du plateau Moldane. An. sti. Univ. Iasi. N.S. St. nat. 2 311-314. [French résumé.]
- McCamley, F. The beauty of living molluses. Proc. R. zool, Soc. N.S.W. 1954-55 1956: 70-71.
- McClelland, W. F. J. The differentiation of some East African species of Bulinus likely to transmit Schistosomes. Trans. R. Soc. trop. Med. Hyg. 49 4 1955: 295.
- McClelland, W. F. J. (1). Studies on snail vectors of schistosomiasis in Kenya. J. trop. Med. Hyg. 59 229-242 figs.
- McCullough, F. S. Apparent resistance of a snail to infection in the Gold Coast. Nature, Lond. 176 1955: 981–982.
- McCullough, F. S. (1). Observations on Schistosoma haematobium infection and on the vector snails at Tema in the Gold Coast. W. Afr. med. J. 4 1 1955: 18-24 map.
- McCullough, F. S. (2). Transmission of Schistosoma haematobium by Bulinus sp. in the Ke district of the Gold Coast. Trans. Soc. trop. Med. Hyg. 50 5 449-457.
- Macdonald, R. & MacWilliams, J. G. Raised beach section at Groomsport, Co. Down. J. Conch. Lond. 24 3 103-104.
- McFarland, H. B. See Peck, jr., J. H.
  - McGee-Russell, S. M. See Billett, F. McKellar, I. C. See Campbell, J. D.
- McKerrow, W. S., Taylor, S. R., Blackburn, A. L. & Ahrens, L. H. Occurrence of caesium in fossils. Nature, Lond. 178 204 1 fig.
- Mackie, T. T., Hunter, G. W. & Worth, C. B. Medically imported molluses. A manual of tropical medicine. 2nd edn. Philad. 1954: 625-640.

Mackin, J. G. See Stein, J. E.

Mackintosh, N. A. 2000 feet down. Deep-sea photographs teach us more about the little-known squid. New Scient. No. 1:60 2 figs.

McLean, J. D. The foraminifera of the Yorktown formation in the York-James Peninsula of Virginia, with notes on the associated mollusks. Bull. Amer. Paleont. 36 160: 225-394 19 pl.

McMichael, D. F. Notes on the Freshwater mussels of New Guinea. Nautilus, Philad. 70 2:38-48 text map pl. 2.

McMichael, D. F. (1). Problems of Family Nomenclature. Syst. Zool. 5: 141-142.

McMichael, D. F. (2). Trigonia—a living fossil. Aust. Mus. Mag. 12:82–84 3 figs.

McMichael, D. F. & Whitley, G. P. The published writings of Tom Iredale with an index of his new scientific names. Aust. Zool. 12:211-250.

McMillan, N. F. Notes on local nonmarine Mollusca [Lancashire and Cheshire.] Proc. Lpool Nat. Fld Cl. No 94 1955: 19-20.

McMillan, N. F. (1). The opisthobranchiate Mollusca of Lancashire and Cheshire. II. Rep. Lancs. Chesh. Fauna Comm. 30th. [1955]: 39-41.

McMillan, N. F. (2). Addendum [to paper on result of a gale, by the Rev. Prof. L. W. Grensted.] J. Conch. Lond. 24 4:46.

McMillan, N. F. (3). Marine aliens in Cornwall. J. Conch. Lond. 24:80.

McMillan, N. F. (4). Two forms of Lymnaea auricularia (L.) in one pond. J. Conch. Lond. 24 3:93-94.

McMillan, N. F. (5). Lymnaea stagnalis (L.) in Anglesey. J. Conch. Lond, 24 3:101.

McMillan, N. F. (6). Notes on local Mollusca. Proc. Lpool Nats' Fld Cl. No. 95:13-15.

McMillan, N. F. & Millott, J. O'N. Notes on the non-marine Mollusca of Cheshire and North Wales. Proc. Chester Soc. nat. Sci. [5] 1954: 109-113.

Macneil, F. S. Stenzelia, new name for Trinacriella Macneil. J. Paleont. 28 2 1954:217.

MacWilliams, J. G. See MacDonald, R.

Maghraby, A. M. el & Perkins, E. J. Additions to the marine fauna of Whitstable. Ann. Mag. nat. Hist. (12) 9:481-496.

Maglaperidze, K. S. See Dzvelaya, M.

Magne, A. See Vigneaux, M.

Magné, J. See Freneix, S.

Mahler, F. Die Molluskenfauna der Tiefstein-Schlucht bei Schludorf. Mitt. naturw. Arbeitsgem. 5-6: Zool. 18-25.

Mahler, F. & Sperling, P. Ein Beitrag zur Molluskenfauna der drei Beitrag zur Molluskenfauna der drei Mitt. naturw. Arbeitsgem. 5–6: Zool. 3–17.

Mahmoud, I. G. E.-D. Etudes paleontologiques sur la faune Cretacique du Massif du Moghara (Sinaï-Egypte). Publ. Inst. Desert Egypte No. 8 1955: 1–195 19 pls.

Majidi, P. See Sordi, M.

Major, H. & Winsmer, T. S. Cambrian and Ordovician fossils from Sørkapp Land Spitsbergen. Skr. norsk Polarinst. No. 106 1955: 1–47 5 pls., 8 figs. map.

Makhnach, A. See Stefanenko, A.

Makridin, V. P. & Zinovyev, M. S. New findings of Ammonites in upper Jurassic deposits of north-western edge of Donetz folded erection). C.R. Acad. Sci. URSS 101 3 1955: 537-539 4 figs. [In Russian.]

Maksimora, S. V. Ammonites from the lower part of the Schwagerina layers of the river Yurezan. Trav. Inst. paléont. Acad. Sci. URSS 14 4 1948: 1-43 4 pls. figs. [In Russian.]

Maksimović, B. V. & Marković, B. Nouvelle contribution à la connaissance de l'albien et du cénomanien de la Serbie. Fauna de l'albien et du cénomanien des environs de Kadina Luka et de Rajac. [Lists—Serbian and French résumé p. 208.] Sborn. Radova geol. Inst. Beograd 5 1953: 183–208, pls. 1–10 text-figs. 1–4.

Makuta, T. See Tsuchiya, Y.

Malatesta, A. Contributi alla conoscenza delle faune neogeniche e quaternarie della Sicilia. III. Faune a Cyprina islandica L. tra Piazza Armerina e Mazzarino (Sicilia centro meridionale) IV. Dentilucina doderleini D. St. nel Pliocene inferiore della Sicilia. Boll. Uff. geol. Ital. 77 1955: 447–454; 455–460 1 pl.

Manchon, P. See Glahn, P. E. and others.

Mancuso, V. L'azione dell'azide sodico sullo soiluppo dell' novo di *Physa rivalaris* Ph. R.C. Inst. sup. Saint. 19 6: 549–554 figs.

Mandahl-Borth, G. Acutorbis n. nom. Arch. Molluskenk. 85 1/3:83.

Mann, H. The influence of Mytilicola intestinalis (Copepoda parasitica) on the development of the gonads of Mytilus edulis. Rapp. Cons. Explor. Mer 140 3:57-58.

Mann, H. See Meyer-Waarden.

Marcus, E. & E. Sea-hares and sidegilled slugs from Brazil. Bol. Inst. Oceanog. S. Paulo 6 1955: 3–49 8 pls.

Marcus, E. & E. (1). Zwei atlantische Onchidellen. Kieler Meeresforsch. 12 76-84, 3 pls.

Marcus, E. & E. (2). On two sacoglossan slugs from Brazil. Amer. Mus. Novit. No. 1796: 1–21–23 figs.

Marcuzzi, G. Fauna della Dolomiti. Mem. Ist. veneto Cl. Sci. mat. nat. 31: 1-595 16 pls. 73 figs.

Marinov, N. A. New date on the finding of Triassic deposits in Eastern Trans-Baikal. C.R. Acad. Sci. URSS. N.S. 98 6 1954:1021-1022. [In Russian.]

Märkel, K. Über die Radulafunktion einiger einheimischer hungenschnecken. Verh. dtsch. zool. Ges. 19:453-459, 7 figs.

Marković, B. see Maksimović, B. V.

Marković, O. & Andelkovic, M. Géologie et léctoniques des environs des villages d'Osečenica, Breždje et Struganik (Serbie occidentale). [Lists] [Serbian and French résumé]. Zborn. Radova geol. Inst. Beograd. 5 1953: 111-145 (Fr. rés. 145-49) pls. 1-10, text-figs. 1-10 2 geol maps.

Marković-Stefanović, O. Nouvelle contribution à la connaissance de la situation stratigraphique des couches crétacées aux environs du village Klence près de Beograd. [Serbian and French résumé] short list. Zborn. Radova geol. Inst. Beograd, 4 1953: 193–205 (Fr. rés. p. 203) pls. 1–5.

Marple, M. F. See La Rocque, A.

Mars, P. Observations sur les variations de *Cardium edule*—Linné. Bull. Soc. linn. Provence 16 1948: 5-6.

Mars, P. (1). Observations sur quelques récoltes malacologiques dans les herbiers méditerranéens. Bull. Soc. linn. Provence 18 1951:15–18.

Mars, P. (2). Sur quelques especes du genre Risson (Moll. Gast.). Boll. Mus. Civ. St. nat. Venezia 9: 27-62 l pl.

Mars, P. (3). Faunes malacologiques du Pliocéne et du Quaternaire de Milazzo (Sicile). Bull. Mus. Hist. nat. Marseille 16: 33-52.

Mars, P. & Ottmann, F. Sur quelques gisements de quatenaire marin de Bas-Languedoc. Bull. Mus. Hist. nat. Marseille 15 1955:133-139 2 figs.

Marshall, F. H. A. and others. Physiology of reproduction 1 London (Longmans Green) 3rd edit. 1956, pp. xix + 688, text illust.

Marteil, L. Acclimatation du clam (*Venus* mercenaria L.) en Bretagne. Rev. Trav. Inst. Pêches marit. 20: 157–160 f. 1.

Martinis, B. Contributo alla Paleontologia del huteziono di Rocca Bernarda (Udine). Riv. Ital. Pal. Strat. 61 4 1955: 187–200 1 pl.

Martinson, G. G. Novuie mezozoĭskie presnovodnuie ghastropodui iz vostochnogho zabaĭkal'ya. Annu. Soc. paléont. russe 13 1949: 77–82 figs.

Martinson, G. G. (1). The stratigraphy of Mesozoic continental deposits of Trans Baical. C.R. Acad. Sci. URSS 105 2 1955: 335-338. [In Russian].

Martinson, G. G. (2). The complexes of various types of freshwater molluses in the Tertiary deposits of Sitz-Yan. C.R. Acad. Sci. URSS 102 3 1955:591-593. [In Russian.]

Martinson, G. G. (3). The lake basins in Asia's geological pasted and their fauna. Priroda 4 1955: 78–82 2 figs. [In Russian.]

Martinson, G. G. (4). The agedifferences of the Continental deposits of Transbaical. Bull. Acad. Sci. U.R.S.S. geol. 2 1955: 82-90. [In Russian.]

Martoja, R. & May, R. M. Les cellules neuro-sensorielles et le ganglion sous-acétabulaire dans l'appareil branchial du céphalopode "Octopus vulgaris Lamarck." C.R. Acad. Sci. Paris 240 1955: 2440–2442 2 figs.

Martoja, R. & May, R. M. (1). Comparaison de l'innervation brachiale des céphalopodes Octopus vulgaris Lamarck et Sepiola rondeleti Leach. Arch. Zool. exp. gén. 94 fasc. 1:1-59 f. 1-18.

Martoja-Pierson, M. Aspects du réceptacle séminal chez quelques Prosobranches Sténoglosses. Bull. Soc. zool. Fr. 81:173–174.

Marwick, J. Proposed use of the plenary powers to designate a type species for the genus *Torquesia* Douvillé, 1929. Bull. zool. Nom. 11:353-355.

Masuda, K. Miocene mollusca from Noto Peninsula Japan, part 1. Trans. Proc. palaeont. Soc. Japan N.S. No. 20 1955: 119–127 1 pl.

Masuda, K. (1). On the Miocene Pectinidae from the environs of Sendai: Part 7: On Pecten kaneharai Yokoyama. Trans. palaeont. Soc. Japan 22:175-183 1 pl. fig.

Masuda, K. (2). On the Miocene Pectinidae from the environs of Sendai; Part 8, On Pecten (Patinopecten) kimurai matumoriensis Nakamura. Trans. palaeont. Soc. Japan 23:221–224 1 pl.

Masuda, K. (3). On the Miocene Pectinidae from the environs of Sendai; Part 9, On Pecten (Chlamys) miyatokoensis Nomura and Hatai. Trans. palaeont. Soc. Japan 24: 247–250 1 pl.

Matekin, P. V. The taxonomic position of Succinea chinensis Pf. 1857 (Gastropoda, Pulmonata, Stylommatophora). C.R. Acad. Sci. URSS 111: 901-902 3 figs. [In Russian.]

Matekin, P. V. (1). New data concerning the area of distributino of Succinea elegans Risso (Gastropode, Pulmonate Stylommatophore). C.R. Acad. Sci. URSS 111 5 1956:1144-1146 3 figs. [In Russian.]

Matekin, P. V. (2) Systematic position of some species of land molluscs in the fauna of Tyan-Shan. Bull. Soc. Nat. Moscou Biol. 59 2 1954: p. 90. [In Russian.]

Matsumoto, T. Yebisites, a new Lower Jurassic Ammonite from Japan. Trans. palaeont. Soc. Japan 23:205-212 1 pl.

Matsumoto, T. (1). Further notes on the Kossmaticeratids from Hokkaido. (Studies on the Cretaceous ammonites from Hokkaido and Saghalien). Jap. J. Geol. Geogr. 27:173-187 3 pls. 2 figs.

Matsumoto, T. See Saito, R.

Matsumoto, T. & Obata, I. On some Cretaceous fossils collected by Mr. T. Kubo along the river Ikusyunbeta, Hokkaido. Nat. Sci. & Mus. Tokyo 23 5-6:22-26 2 pls. [In Japanese.]

Maubeuge, P. L. Sur quelques échantillons anormaux d'Ammonites jurassiques. Un cas possible de mutation chez les ammonites. Arch. Inst. Gr.-Duc. Luxemb. N.S. 18 1949: 127-147 3 pls.

Maubeuge, P. L. (1). Notes paléontologiques. Sur quelques ammonites jurassiques rares ou nouvelles de le region frontière franco-luxembergeoise et de la Lorraine centrale. Arch. Inst. Gr.-Duc. Luxemb. N.S. 18 1949:149– 178 17 pls.

Maubeuge, P. L. (2). Sur quelques ammonites de l' "Aalénien ferrugineux" du Luxembourg et sur l'échelle stratigraphique de la formation ferrifère franco-belgo - luxembourgeoise. Arch. Inst. Gr.-Duc. Luxemb. N.S. 17 1947: 73-87 2 pls.

Maubeuge, P. L. (3). Nouvelles recherches stratigraphiques et Paléonitologiques sur l'Aalénien luxembourgeoi geois. (Parties Moyenne et Superieure). Arch. Inst. Gr.-Duc. Luxemb. N.S. 19 1950: 365–397 16 pls.

Maubeuge, P. L. (4). Sur l'Oxfordien supérieur et l'Argovien dans l'Est du Bassin de Paris, C.R. Acad. Sci. Paris 241 1955: 812-814. Maubeuge, P. L. See Charles, R. P.

Maubeuge, P. L. & Lambert, R. Sur quelques ammonites Aalèniennes d' Argentine. Bull. Soc. Belge. Géol. 64 (1955) 1956: 620-624 1 pl.

Maviglia, C. Conchiglie di Limax duplex Poll., di problematica provenienza, rinvenute con resti umain e ceramica preistorica nella grotto del Tamborin (2172 Lo.) in Val Bova (Como). Natura, Milano 47: 164–117 1 fig.

Mawatari, S. & Kobayashi, S. Seasonal settlement of animal fouling organisms in Ago Bay, middle part of Japan. I–II. Misc. Rep. Res. Inst. nat. Resour. Tokyo No. 35 1954:37–47 figs.; No. 36 1954:1–8 pls. figs.

May, R. M. See Martoja, R.

Mayer, G. Eine interessante Schichtfläche aus dem Mittleren Hauptmuschelkalk von Bruchsal. Beitr. naturk. Porsch. Sudw Dtsch. 14 1955: 114–118 2 pls. 3 figs.

Maync, W. (in Rod. E. & Maync, W.). Revision of lower Cretaceous stratigraphy of Venezuela. [Lists of spp. in text]. Bull. Amer. Ass. Petrol. Geol. 38 2 1954: 260-283.

Mayr, E. & Rosen, C. B. Geographic variation and hybridization in populations of Bahama snails (*Cerion*). Amer. Mus. Novit. No. 1806: 1-48 9 figs.

Mazenot, G. Recherches sur les faunas malacologiques du loess recent Wurmien et de divers limons terrestres holocenes dans le Sud-est de la France. Bull. mens. Soc. linn. Lyon 25 1:9-24 3 figs.

Mazenot, G. (1). Fauna malacologique de divers limons terrestres holocènes du Sillon Rhodanien et de ses abords, des Vallies de la Cèze et de l'Aygues à la région du Bas-Rhone. [List]. Bull mens. Soc. linn. Lyon 25 3:73

Mazzotti, L. Lynnaea humilis (Say), huesped intermediario de Fasciola hepatica. Rev. Inst. Salubr. Enferm. trop. Méx. 16 4: 21–23 1 fig. [English summary.]

Mead, A. R. Disease in the giant African snail Achatina fulica Bowditch. Science 123:1130-1131 1 fig.

Mead, A. R. (1). Predators need defending. Nautilus, Philad. 70 2: 65-69.

Medcof, J. C. & Dickie, D. M. Watch for the green crab—a new clam enemy. Circ. Atlantic biol. Sta. No. 26 1955:1 2 figs.

Medeiros, L. C. M. de. See Azevedo, J. F. de.

Medeiros, L. de. See Ranson, M. G.

Medwenitsch, W. Zur Geologie Vardarisch-Makedoniens (Jugoslawien), zum Problem der Pelagoniden. S.B. öst. Akad. Wiss. Abt. I. 165: 397–473 2 pls. I1 figs. [English summary.]

Meel, L. v. See Leferre and Leloup.

Meel, L. v. See Leloup.

Meenakshi, V. R. The excretory spherioles in the digestive gland of *Pila virens*. J. Anim. Morph. Physiol. 3: 75-78.

Meenakshi, V. R. (1). Seasonal variations in the glycogen and fat content in the apple snail, *Pila virens* (Lamarck). J. zool. Soc. India 8: 57-62.

Meenakshi, V. R. (2). Physiology of hibernation of the apple-snail, *Pila virens* (Lamarck). Curr. Sci. 25 10: 321-322 1 fig.

Meenakshi, V. R. (3). Caratenoid in *Pila virens* (Lamarck). Curr. Sei. 25 4:124-125.

Meenakshi, V. R. (4). How Pila without protease meets its protein requirements. Proc. Indian sci. Congr. Ass. 42nd 3 1955: 279 (Abstract).

Meenakshi, V. R. (5). Relative strength of digestive gland anylase in some common south Indian Gastropods. Proc. Indian sci. Congr. Ass. 43rd 3: 303. (Abstract).

Meise, W. See Jaeckel, S.

Melendez, B. Notas paleontologicas regionales. II.—Asturias. An. Asoc. esp. Progr. Cienc. 19 3 1954: 655-662 2 pls., 3 figs.

Mercier, J. see Ripplinger, J.

Mercier, J. See Jullien, A. and others.

Merklin, R. L. Contribution to the knowledge of Tertiary fauna of northeastern Siberia. C.R. Acad. Sci. URSS N.S. 95 2 1954: p. 379-380 2 figs. [In Russian.] Merklin, R. L. & Nevesskaya, L. A. Checklist of bivalve molluses from the Miocene of Turkmenstan and Western Kazakhstan. Trav. Inst. Paléont. Acad. Sci. URSS 59 1955: 1-115 pls. figs.

Mertens, R. See Zilch, A.

Meyer, K. O. Naturgeschichte der Strandschnecke Ovatella myosotis (Draparnaud). Arch. Mollusk. 84 1/3 1955: 1–43 30 text figs pls. 1, 2.

Meyer, K. O. See Knipper, H.

Meyer-Waarden, P. F. Mussel culture in the German Waddensea. Rapp. Cons. Explor. Mer 140 3:47-48.

Meyer-Waarden, P. F. & Mann, H. German investigations with respect to Mytilicola intestinalis in Mytilus edulis in 1953. Rapp. Cons. Explor. Mer 140 3:54-56.

Meznerics. See Csepreghyné.

Michelson, E. H. See Chernin, E.

Miklukho-Maklai, A. D. & Ruzhentzev, V. E. A faunistic description of the Upper Carboniferous from the Kara-Chatir range in Ferghana, Turkestan. C.R. Acad. Sci. URSS 110 427-429. [In Russian.]

Milbourne, R. A. The Gault at Greatness Lane, Sevenoaks, Kent. Proc. Geol. Ass. Lond. 66 235–242 text-fig. 1.

Millar, R. H. Notes on the mechanism of food movements in the gut of the larval oyster, Ostrea edulis. Quart. J. micr. Sci. 96 1955: 539-544 3 fig.

Miller, A. K. & Carpenter, L. B. Cretaceous and Tertiary nautiloids from Angola. Jta. Inv. Ultramar, Lisbon No. 21 1956: 1–56 6 pls. 5 figs.

Miller, A. K. & Furnish, W. M. The classification of the Paleozoic ammonoids. J. Paleont. 28 1954: 685-692 text-figs. 1-2.

Millot, J. O.' N. See McMillan, N.

Milne, A. See Clark, R. B.

Miloslavskaya, N. M. The distribution of pelecypods in eastern Murman in connection with the oscillation of the temperate in the environment. C.R. Acad. Sci. URSS N.S. 99 4 1954: 633-635 diag. [In Russian.]

Miloslavskaya, N. M. (1). Ecology of Macoma baltica. C.R. Acad. Sci. URSS 101 3 1955:565-567 fig. [In Russian.]

Miroshnikov, L. D. On Mesozoic deposits from the Northern Taimyī. C.R. Acad. Sci. URSS 111:676-677. [In Russian.]

Mishima, T. See Yagi, Y.

Mistakidis, M. N. & Hancock, D. A. Reappearance of Ocenebra erinacca (L.) off the East Coast of England. Nature, Lond. 175 1955: 734.

Mizuno, T. Relation between zinc and sperm motility in some marine forms. J. Fac. Sci. Tokyo Univ. (Zool.) 7 Pt. 4:477-487.

Moinet, J. C. See Dastugue, G.

Moiseev, A. S. On the corals and other organisms in the limestones of Primose Land. Trav. Leningrad. Soc. Nat. 68 2 1951: 208-237. [In Russian.]

Molinier, R. Les plate-formes et corniches récifoles de vermets (Vermetus cristatus Biondi) en Méditerranée occidentale. C.R. Acad. Sci. Paris 240 1955: 361–363.

Monakhova, L. P. Contribution to the stratigraphy of the lower part of coal-bearing deposits of Carboniferous of Karagand basin. C.R. Acad. Sci. URSS N.S. 95 4 1954: 865–867 fig. [In Russian.]

Moncharmont Zei, M. Sopra una nueva specie di *Parastrophia* del Quaternario della Punto delle Pietre Nere (Foggia). Boll. Soc. Nat. Napoli 63 1955: 118–120 1 pl.

Moncharmont Zei, M. (1). Contributo alla conoscenza del Calabriano della Valle dell'Ofanto. Boll. Uff. geol. Ital. 77 1955: 499-554 10 pls. fig.

Mongin, D. Note préliminaire sur une faune "sarmatienne" saumâtre recueillie à Bled Tekouka, au S de Mercièr-Lacombe (Algérie). C.R. Soc. géol. Fr. 13 1952: 270-72.

Mongin, D. (1). Révision stratigraphique du Burdigalien de Basse-Provence. Ann. Fac. Sci. Marseille 25 1956: 1–74 18 figs. map.

Mongin, D. et Trouvé, P. Le Valanginien inférieur calcaire du Grand Cañon du Verdon (Basses Alpes). Bull. Soc. géol. Fr. 6 3 1–3 1953: 223–239 pl. 3 text-figs. 1–4.

Montalban, H. R. See Blanco, G.

Moon, H. P. The distribution of Potamopyrgus jenkinsi (E. A. Smith) in Windermere. Proc. Malac. Soc. Lond. 32 3: 105-108 text-map.

Moore, D. R. Observations of predation on echinoderms by three species of Cassididae. Nautilus, Philad. 69 3: 73-76.

Morais, T. de. Subsidios para o estudo dos moluscos vectores da bilharziose na Zambézia (Moçambique). An. Inst. Med. trop., Lisbon 13:883–899 8 figs. [English summary.]

Moreau, R. See Jullien, A.

Mori, K. On the Jurassic formation in the Hashuira district, Province of Pikuzen, Japan. Jap. J. Geol. & Geogr. 21 1949: 315-321 1 fig.

Morton, J. E. The evolution of the Ellobiidae with a discussion on the origin of the Pulmonata. Proc. zool. Soc. Lond. 125 1955: 127-168 15 figs.

Morton, J. E. (1). The functional morphology of *Otina otis*, a primitive marine pulmonate. J. Mar. biol. Ass. U.K. 34 1955:113-150 12 figs.

Morton, J. E. (2)., The functional morphology of the British Ellobiidae (Gastropoda Pulmonata) with special reference to the digestive and reproductive systems. Phil. Trans. 239B 1955: 89–160 31 figs.

Morton, J. E. (3). The tidal rhythm and action of the digestive system of the lamellibranch *Lasaea rubra*. J. mar. biol. Ass. U.K. **35**: 563–586 8 figs.

Morton, J. E. (4). The vascular and nervous systems of *Struthiolaria* (Prosobranchia, Mesogastropoda). Trans. roy. Soc. N.Z. 83: 721-743 7 figs.

Morton, J. E. See Ballantine, D.

Morton, J. E. & Holme, N. A. The occurrence at Plymouth of the Opistobranch Akera bullata, with notes on its habits and relationships. J. Mar. biol. Ass. U.K. 34 1955: 101-112 1 pl. 2 figs.

Motwani, M. P. Experimental and ecological studies on the adaptation of Mytilus edulis L. to salinity fluctuations. Proc. nat. Inst. Sci. India 21B 5-6 1955: 227-246 7 figs.

Müller, K. J. Cephalopodenfauna und Stratigraphie des Oberdevons von Schleiz und Zeulenroda in Thüringen. Beiheft. Geol. Jahrsb. 20: 1–93 2 pls. figs.

Müller-Liebenau, I. Die Besiedlung der Potamogeton-Zone ostholsteinischer Seen. Arch. Hydrobiol. **52**: 470–606 6 pls. 10 figs.

Muraour, P. Contribution a l'etude stratigraphique and sedimentologique de la Basse-Kabylie etc. Bull. Pub. Carte, géol. Alger N.S. 7:1-383 10 pls. figs.

Muravyer, I. S. & Ivanov, E. E. The Kinovsk layers of South-Eastern Tataria, Kazansk Volga course and Idmutria. C.R. Acad. Sci. URSS 101 4 1955: 739-742. [In Russian.]

Muromtzev, V. S. The age of Ostrog suite of Kuzbass. C.R. Acad. Sci. URSS N.S. 95 5 1954:1059-1061. [In Russian.]

Muus, B. J. Development and distribution of a north Atlantic pelagic squid, Family *Cranchiidae*. Medd. Danm. Fisk. Havundersog. N.S. 1 Nr. 15: 1-15 9 figs.

Nagabhushanam, R. On the development of the marine woodboring molluse, *Martesia striata* (Linn.). Sci. & Cult. 22 1:31-33 1 fig.

Nagabhushanam, R. (1). The role of water filtration in the marine woodboring mollusc, *Martesia striata* Linn. Proc. Indian Acad. Sci. **43B** 4: 223-227 3 figs.

Nagabhushanam, R. See Ganapati,

Nair, N. B. (1). Destruction of timber structures by shipworms in Madras waters. J. Sci. Indust. Research 156 3 1957: 81, 82

Nair, N. B. (2). The development of the wood-boring Pelecypod Bankia indica Nair. J. Madras Univ. B26 2 1956: 303-318 text figs. 1-10.

Nair, N. B. (3). Physiology of digestion in Bankia indica. The Enzymatic activity of the Digestive Diverticula. J. Madras Univ. B26 3 1956: 599-627 17 tables.

Nair, N. B. (4). The path of enzymic hydrolysis of cellulose in the woodboring Pelecypod Bankia indica Nair. J. Sci. & Indust. Research 150 6 1956: 155, 156 text figs.

Nair, N. B. (5). On a new species of Shipworm of the subgenus Neobankia from Madras. J. Madras Univ. B25 1 1955 pp. 109-113.

Nair, N. B. (6). Sex changes in the wood-boring Pelecypod Bankia indica Nair. J. Madras Univ. **B26** 2 1956 277-280 2 plates & table.

Nair, N. B. & Gurumani, O. N. new shipworm from Tondi, South India. Curr. Sci. 25: 361-362 3 figs.

Nakazawa, K. A study of the pelecypod-fauna of the Upper Triassic Nabae group in the northern part of Kyoto Prefecture, Japan. Part 4, Cardinioides, Homomya, Pleseromya and others, supplement and brief summary. Mem. Coll. Sci. Univ. Kyoto 23B: 231-253 4 pls.

Nakazima, M. On the structure and function of the midget-gland of mollusca, with a general consideration of the feeding habits and systematic relations. Jap. J. Zool. 11:469-566 10 pls.

Nalivkiu, B. V. Pseud Pelecypoda for the domanic. Pseudosealar Acad. Sci. URSS 111: 188-189 fig. [In Russian.]

Nambudri, C. N. The alkaline phosphatase activity in the gills of freshwater mussel Potamida (Parreysia) corrugata. J. Anim. Morph. Physiol. 3:87-90.

Nath, V. Golgi bodies in the spermatogenesis of pulmonate gastropods. Nature, Lond. 175 1955: 905-906 2 figs.

Nath, V. & Chopra, H. C. On the origin of the golgi Dictyosomes from the mistochondria and acrosome formation in the spermatogenesis of the slug Anadenus altivagus (Theobald). Research Bull. Panjab Univ. Hoshiampur No. 74 pp. 91-100 text-fig. 1A-F. pl. 1.

Nath, V. & Gupta, B. L. The morphology and origin of the Golgi bodies and their role in the secretion of the acrosome in the spermatogenesis of pulmonate gastropods as determined in the living material by phase-contrast microscopy. Quart. J. micr. Sci. 97: 369–377 4 figs.

Nazan, H. El. See Din, Sharaf El.

Neale, J. W. Chlamys (Radulopecten) drewtonensis sp. nov. Proc. Yorks. geol. Soc. 30:371-374 1 pl.

The course of excretion Nejedlýik. in the edible snail (Helix pomatia L.). Rospr. české Akad. 63 5 1953: 45-56 4 pls. [English summary.]

Nelson, T. C. In Johnson F. The Boylston Street Fishweir. Pap. Peabody Foundation 2 1942: 1-212 figs. [Oysters p. 49-64].

Nelson, T. C. & Allison, J. B. The influence of spermatic hormones on spawning and water flow in the oyster, Crassostrea virginica. Int. Congr. Zool. 14 1953 (1956): 435-439.

Nevesskaya, L. A. To the biostratigraphy of the marine Quartenary deposits of Turkmenia. Bull. Soc. Nat. Moscou geol. 31 (3): 65-79 2 plts. 2 figs. [In Russian.]

Nevesskaya, L. A. See Merklin, R. Newell, G. E. See Chapman, G.

Newell, N. D. Permian pelecypods of East Greenland. Medd. Grønland 110 4 1955: 1-36 5 pls.

Nicol, D. Growth and decline of populations and the distribution of J. Paleont. 28 marine pelecypods. 1954: 22-25 text-figs. 1-2.

Nicol, D. (1). Nomenclatural review of genera and subgenera of Cuccullaeidae. J. Paleont. 28 1954: 96–101.

Nicol, D. (2). The taxonomic significance of gaps in Pelecypod morphology. Syst. Zool. 5:143.

Nicol, D. (3). Distribution of living Glycymerids with a new species from Bermuda. Nautilus, Philad. 70 2: 48-53 text map and pl. 3.

Nicolás, E. G. S. See Garcia San Nicolás.

Nüno, H. On the bottom deposits on the banks at the mouth of Wakasa Bay and on the adjacent continental shelf. Jap. J. Geol. and Geogr. 21 1949:193-230 2 pls.

Nilsson, A. Cochlicopa nitens (Kokeil) Gallenstein und C. minima Siemashko, zwei selbständige Artem im Formenkreis der kollektiven C. lubrica (Müller). Ark. Zool. 9:281-304 1 fig. [English summary.]

Nilsson, A. (1). Landmolluskfaunan i kalkkärret vid Örups almskog. Skånes Nat. 43: 103–112 I fig.

Niță, P. P. Sur la présence de Valenciennius annulatus Rouss dans les dépôts pliocènes du Sud Cocorova (district de T. Severin, région de Craiova). Comun. Acad. Republ. rom. 63:459-463 5 figs. [French résumé.]

Nitahara, I. See Ishida, S. and others.

Nolan, M. O. See Weinbach, E.

Nolan, M. O. & Bond, H. W. Results of laboratory screening tests of chemical compounds for molluscoidal activity. III. Derivatives of abietic acid. Amer J. trop. Med. Hyg. 4 1955:152-

Nolte, D. J. Polymorphic coloration in *Donax*. Caryologia 6 Suppl. Atti IX Congr. Internat. Genet. pt. 2 1954: 889–890.

Nomejko, C. A. See Loosanoff, V.

Novik, E. O. Stratigraphy of Devonian deposits in Dneper-Donetz depression. Bull. Acad. Sci. URSS Geol. 2 1954:44-54. [In Russian.]

Nowak, W. See Krach, W.

Obata, I. See Matsumoto, T.

Obr, S. Hydrobiologische Untersuchung der Fauna des Orava-Flussgebietes mit Hinsicht auf die Wasserreinheit. Acta Acad. Sci. Čech. 28: 377–445. [German summary.]

Ockelmann, W. K. On the interrelationship and the zoogeography of Northern species of Yoldia Möller, S. Str. (Mollusca, Fam. Ledidai). Medd. Grønland 107 7 1954: 1-32 2 pls., 2 figs.

Odhner, N. Distinctions between Anisus (Armiger) crista (L.) and Anisus (Gyraulus) riparius (West.) Proc. Malac. Soc. Lond. 32 3:127-131 pl. 24.

Ohba, S. Analysis of activity rhythm in the marine gastropod, Nassarius festivus, inhabiting the tide pool. II. Nocturnal activity and its artificial control by light. Biol. J. Okayama Univ. 1 4 1954: 209-216 4 figs.

Ohba, S. (1). Effects of population density on mortality and growth in an experimental culture of a bivalve, *Venerupis semidecussata*. Biol. J. Okayama Univ. 3 4:169-173 1 fig.

Ohgushi, R. Observations on the behaviour of non-homing Limpets. Venus, Fukuyama 19 1:60-64. [In Japanese.]

Ohm, P. & Remmert, H. Etudes sur les rockpools des Pyrénées-Orientales. [Moll. pp. 199-200.] Vie et Milieu 6 1955: 194-209 5 figs.

Ohsawa, W. & Tsukuda, H. Extruding response of the periwinkle Nodilittorina granularis (Gray). J. Inst. Polyt. Osaka 6D 1955: 71–96 16 figs.

Okada, H. See Imai, T.

Okada, Y., Tsujii, T. Watabe, N. & Yoshii, G. Studies on pearl formation mechanism by radioantography. Rep. Fac. Fish. Univ. Mic. 2:199-204 1 pl.

Okada, Y. Watabe, N. & Tsujii, T. Studies on the cultured pearl in Japan. Int. Congr. Zool. 14 1953 (1956): 440-441.

Okamura, N. See Usuki, I.

**Okutani, T.** Some ecological observations of *Janthina*. Venus, Fukuyama, **19** 1:43-48 text figs.

Okutani, T. See Takemura.

Olivier, L. & Schneiderman, M. A method for estimating the density of aquatic snail populations. Exp. Parasit. 5:109-117.

Olson, O. P. The genus Baryspira (Mollusca) in New Zealand. Palaeont. Bull. N.Z. 24: 1–32 7 pls.

Olsson, A. A. Studies on the genus Olivella. Proc. Acad. nat. Sci. Philad. 108: 155-225 9 pls.

Omara, S. See Farag, J.

Origny, M. R. Observations sur la biologie de Valvata piscinalis. C.R. Ass. franç. Av. Sci. 73 1955: 458-459.

Orr, V. The South African gastropod genus Burnupena (Buccinidae). Proc. Acad. nat. Sci. Philad. 108: 249–263 2 pls. 2 figs.

Ortiz de Lárate. See López and Rocandio.

Orton, J. H., Southward, A. J. & Dodd, J. M. Studies on the biology of limpets. II. The breeding of Patella vulgata L. in Britain. J. Mar. biol. Ass. U.K. 35: 149-176 2 pls. 4 figs.

Osipova, A. I. On the stratigraphic importance of the fauna and the Kaplanbek complex from the Paleogene of Central Asia. C.R. Acad. Sci. URSS 111:1076-1079. [In Russian.]

Osipova, A. I. & Liverovskaya, E. V. To the characteristics of the Tedzhiksk Paleogene Sea and its fauna during the Suzak and Alay period. Bull. Akad. Nauk. geol. 12 82-90. [In Russian.]

Óskarsson, I. Íslenskar saeskeljar. Náttúrufraedinguriun 22 1 1952: 1–18 figs. maps and graphs.

óskarsson, I. (1). Saeskelin Cardium edule L. fundin vid Ísland. Náttúru-fraedinguriun 23 1 1953: 40-41 1 fig. [English summary.]

Óskarsson, I. (2). A spècies new to Iceland. Náttúrufraedinguriun 25 2 1955: 103.

Ottmann, F. See Mars, P.

Otuka, Y. Fossil Mollusca and rocks of the Kiyosami group exposed at Minoto-machi, Chiba Prefecture, and its environs (1st paper). Jap. J. Geol. and Geogr. 21 1949: 295-312 1 pl. 3 figs.

Owen, G. Observations on the stomach and digestive diverticula of the Lamellibranchia. I. Quart. J. micr. Sci. 96 1955:517-537 13 figs.

Owen, G. (1). Observations on the stomach and digestive diverticula of the Lamellibranchia. II. The Nuculidae. Quart. J. micr. Sci. 97:541-567 9 figs.

Oyama, K. Fossil communities of the oceanic waters. Misc. Rep. Res. Inst. nat. Resour. Tokyo No. 34 1954: 89-100. [English summary.]

Oyama, K. (1). Mollusca of Aogasima. Misc. Rep. Res. Inst. nat. Resour. Tokyo No. 38 1955: 145–149. [In Japanese.]

Oyama, K. (2). A contribution to the Phylogeny of the Umboniinae. Venus, Fukuyama 18 3 1955: 169-176 text figs. 1-2. [In Japanese.]

Oyama, K. (3). A developmental history of Japanese conchology. III. Conchological works on palaeontology. Nat. Sci. & Mus. Tokyo 23 56:1-21. [In Japanese.]

Ozaki, H. & Katto, J. On a new species of *Cymatoceras* from Nagase, Kôte Prefecture (Sikaku). Bull. nat. Sci. Mus. Tokyo 3:61-66 2 pls. map.

Packard, E. I. An Engonoceras from Central Oregon. J. Paleont. 30: 398– 402 1 fig.

Padget, P. The Stratigraphy of Cuilcagh, Ireland. Geol. Mag. 90 1953: 17-26 text-figs. 1-2.

Pain, T. On a collection of *Pomacea* from Colombia, with description of a new subspecimen. J. Conch. Lond. 24 3:73-79; text fig.

Pain, T. (1). Notes on the generic names *Pomacea* and *Ampullarius*. J. Conch. Lond. 24 3: 79.

Pain, T. (2). Further notes on the non-marine mollusca from semi-arid areas of East Africa. J. Conch. Lond. 24 4:142, 143.

Pain, T. (3). Revision of the Melaniidae of British Guiana and Sarinam. Basteria, Leiden, 20 4/5: 91-105.

Paladian, G. See Grossu, A. V.

Palekar, V. C. & Bal, D. V. A preliminary note on the marine organisms injurious to submerged timber in the Bombay Harbour. Proc. Indian sci. Congr. Ass. 42nd 3 1955: 313. (Abstract).

Palmer, K. v. W. Muséum Demidoff. Nautilus, Philad. 70 1:35.

Papp, A. Die Molluskenfauna des Pannon im Wiener Becken. Mitt. geol Ges. Wien 44 1953: 85-222 pls. 1-25 1 text-fig.

Papp, A. (1). Landschnecken aus Žihlava bei Videm (Slowenien) und bei Veli Opačae SW Zadar (Dalmatian, Jugoslawien). Rozpr. Akad. Ljubljana (4) 3 1955: 425-430.

Papp, A. (2). Ueber den Byssus fossiler Anomyen aus Sanden des Tortons von Haschendorf in Burgenland. Burgenländ. Hermit Bl. 17 1955: 145–147 5 figs.

Paproth, E. Eine Kohlenkalkfauna aus dem Kulmkonglomerat von Frankenberg au des Eder. [Moll. Lists pp. 186-.] Paläont. Z. 27 3-4 1953: 169-207 pls. 11-12 text-figs. 1-2.

Paraense, W. L. A genetic approach to the systematics of Planorbid Molluscs. Evolution 10:403-407.

Paraense, W. L. & Deslandes, N. Studies on Australorbis centimetralis. II. Biospecific characterization. III. Generic status. Rev. bras. Biol. 15 1955: 341-348.

Paraense, W. L. & Deslandes, N. (1). The Brazilian species of Drepanotrema. I. D. anatinum (Orbigny, 1835) II. D. melleum (Lutz, 1918). Rev. bras. Biol. 16:491-499 18 figs. 527-534 4 figs.

Paraense, W. L. & Deslandes, N. (2). Australorbis inflexus sp. n. from Brazil. Rev. bras. Biol. 16:149-158 7 figs.

Paraense, W. L. & Deslandes, N. (3). Diagnostic characters of the Brazilian species of "Australorbis" (Pulmonata, Planorbidae). Rev. bras. Biol. 16: 281–286 4 figs.

Paris, J. Contribution à la connaissance de la "zone nord des cannalots". [List of Moll. p. 497.] Vie et Milieu 5 1954: 469-511 9 figs.

Parmalee, P. W. An analysis of faunal remains found in the Modoc rock shelter, Randolph County, Illinois. Rep. Inv. Ill. St. Mus. No. 4:47-53.

Parness, A. See Avnimeloch, M. and others.

Parodiz, J. J. Araucania twomeyi n. gen. n. sp. del sur de Chile. Neotropica 1 (2) 1954:17-18 1 fig. [English résumé.]

Parodiz, J. J. (1). Cuatio nuevas especies de *Scansicochlea* de Argentina. 1a parte. Neotropica 2 (8): 59-64 5 figs.

Parodiz, J. J. (2). Cuatro nuevas especies de *Scansicochlea* de Argentine. Neotropica 2:77-80 2 figs.

Parodiz, J. J. (3). Dos nuevas *Epi-phragmophora* de la Argentina. Neotropica 1 (6) 1955: 93-95 1 fig.

Parodiz, J. J. (4). Una nueva especie de *Triphora* de Uruguay. Neotropica 1 (4) 1955: 59-60 1 fig. [English summary.]

Parodiz, J. J. (5). La volidez del nombre Paludestrina d'Orbigny 1839. Neotropica 1 (6) 1955: 95-96.

Pasic, M. Contribution à la connaissance de la faune sénonienne aux environs de Beograd (Mašin Majdan). [Serbian with French résumé.] Zborn. Radova geol. Inst. Beograd, 3 1952: 111-114 (Fr. rés. p. 114) pl. 1.

Pasić, M. (1). Geologische und faunistioche Darstellung der Verhältnisse des Čerevićki Potok und seiner Quallenarme (Fruška Gora). [List p. 173, in Serbian with German summary.] Zborn. Radova geol. Inst. Beograd, 4: 1952: 169–192 (Germ. summ. p. 191) pls. 1–4.

Pasteels, J. Évolution de la métachromasie au bleu de toluidine suivre sur le vivant, au cours des premières phases du développement de l'Oursin et de la Pholade. Bull. Acad. Belg. Cl. Sci. (5) 41 7 1955: 761-768 figs.

Pasternak, S. I. See Vyalov, O. S.

Paucă, M. Deux espèces de fossiles rares du pliocène du bassin de Sălaj. Comun. Acad. Republ. Pop. rom. 47-8 1954: 397-400 2 figs. [French summary.]

Paulus, M. Contribution à l'étude de la faune malacologique des anciens cordons littoraux du delta du Rhône. I. Bull. Soc. linn. Provence 13 1 1942: 7-10.

Paulus, M. (1). Contribution à l'étude du Quatermaire du Sénégal. Sur un gisement de mollusques pleistocènes à l'anse Bernard, près de Dakar. Bull. Soc. linn. Provence 17 1949: 17–20.

Pavlović, V. & Bać, M. Adaptation at the outer temperature and glyamic of the snail (Helix pomatia). Glasn. sopsk. Akad. Nauk 222 6a Prirod. Mat. Nauk N.S. 10 1956: 93-96. [English summary.]

Pavlovski, E. V. & Frolova, N. V. Organic remains in metamorphic complexes. Bull. Acad. Sci. U.S.S.R. geol. 6 1954: 15–29 9 figs. [In Russian.]

Peck, Jr. J. H. & McFarland, H. B. Whitfield Collection types at the University of California. J. Paleont. 28 1954: 297-309 pl. 29.

Pei, Hung-kang. See Kuo, J.-C.

Pejović, D. The Senonian fauna from Leštar, not far from the mine Rtanj in East Serbia. Zborn. Radova geol. Inst. Beograd. 4 1952: 113-119 (Engl. summ. 119) pl. 1.

Pejović, D. (1). La faune sinonienne aux environs de Novi Pazar. Zborn. Radova geol. Inst. Beograd, 5 1953: 79-85 (Fr. rés. 86) pls. 1-4.

Pelin, M. See Raileanu, G.

Pemberton, C. E. Defence of a predator [ref. to a paper by Dr. A. R. Mead Naut. 69 2 Oct. 1955 pp. 37-40 a criticism.] Nautilus, Philad. 69 4: 142-144.

48

Perkins, E. J. The fauna of a sandbank in the mouth of the Dee Estuary. Ann. Mag. nat. Hist. (12) 9:112-128 3 figs.

Perkins, E. J. (1). Notes on the fauna of the River Lugg at Shelwick July-August, 1953. Trans. Woolhope Nat. Fld. Cl. 35 1:57-61.

Perkins, E. J. See Maghraby, A.

Permyakov, V. The stratigraphy of the lower-Peruvian deposits of Digoria. Bull. Soc. Nat. Moscou geol. 29 4 1954:51-59. [In Russian.]

Peruzzi, P. & Corda, M. Azione difasica dell'acetilcolina e condizioni di lavoro nel cuore di *Helix adspersa*. Arch. Fisiol. Firenze 56:107-121 10 figs. [English summary.]

Petersen, I. C. See Berg, K.

Petković, K. V. Lumachelle des Cephalopodes et des Inocerames dans les conches sénoniennes de la rivière Osmakovska Reka, son importance biostratigraphique et l'explication di ce phénomène (Serbie Orientale). [Lists and figs. in Serbian with French résumé.]. Zborn. Radova geol. Inst. Beograd 6 1953: 2-54 (Fr. résumé 54-66) pls. 1-17 text-figs. 1-4.

Petrauskas, L. E. A case of cone shell poisoning by "bite" in Manas Island. Papua and New Guinea med. J. 1 2 1955: 67-68.

Petrbok, J. Najady rěky Latorice u Velkých Kapušan na Slovensku. Čas. nár. Mus. 125 : 198–202 figs.

Petrbok, J. (1). The molluses of the Bohemian Holocene. Acta Mus. nat. Prag 12B 1: 1-43 1 pl.

Petrbok, J. (2). The Pleistocene molluses of Sarovce at Hron-river, the vicinity and the supplements to knowledge of the malacofauna of south Slovakia. Geol. Práce Slov. Akad. Bratislava 7:99-100 1 fig.

Petrbok, J. (3). Stratigraphische Chronologie und Fauna Cardenoisienischer Schichten der Matschauschen Hügel bei Sened in der Serwaka. Biológica, Bratislava 10 1 1955: 33–37. [German summary.] Péwé, T. L. Evidence for possible marine Tertiary (?) sediments in central Alaska. Bull. Amer. Ass. Petrol. Geol. 38 10 1954: 2222-2225.

Pfeiffer, K. L. Die Albinarien des Dodekans Moll. Clausiliidae. Arch. Mollusk. 84 4/6 1955: 109-154 3 maps 6 text figs. & pls. 8-11.

Pfeiffer, K. L. (1). Die Albinarien des Dodekanes [Moll. Clausiliidae]. 2. Arch. Mollusk. 85 4/6: 87-119 pls. 5-8.

Phear, E. A. Gut haems in the invertebrates, Proc. zool, Soc. Lond. 125 1955: 383-406 7 figs, tables & graphs.

Philip, G. M. See Talent, J. A.

Pierson, M. Particularités histologies de la glande brune chez Nucella lapillus (L.) (Gastéropode prosobranche) C.R. Acad. Sci. Paris 241 1955: 1168–1170.

Pierson, M. M. See Martoja-Pierson. Pilsbry, H. A. A new Bostryx from Peru. Nautilus, Philad. 69 3:92-93 pl. 5 f. 4, 5.

Pilsbry, H. A. (1). A gastropod domicillary in sea urchin spines. Nautilus, Philad. 69 4:109, 110 pl. 6 f. 4–8.

P[ilsbry], H. A. (2). Southern record for *Pomacea cumingi* (King). Nautilus, Philad. **69** 4:140.

Pilsbry, H. A. (3). Inland Mollusca of Northern Mexico III. Polygyridae and Potadominae. Proc. Acad. nat. Sci. Philad. 108: 19–40 3 pls. 4 figs.

Pilsbry, H. A. & Hubricht, L. Beach drift Polygyridae from Southern Texas. Nautilus, Philad. 69 3:93-96. pl. 5 f. 1-3a.

Pitchford, G. W. Agriolimax reticulatus (Müller) feeding on carrion. J. Conch. Lond. 24 3:96.

Plate, H.-P. Zur Molluskenfauna der Insel Hiddensee. Arch. Fr. Naturg. Mecklenburg 2:307-335.

Pleničar, M. Cretaceous Fauna at Jelšane near Ilirska Bistrica. Rozp. Geol. Poročila 3 1955: 204–207 3 pls. [English summary p. 207.]

Plotnikov, M. A. A new genus of Lamellibranchiata from the Tatarian Stage of the lower course of Sukhona River. Annu. Soc. paléont. russe 12 1945: 138-144 l pl. [English summary p. 143.] (Plotnikov, M. A.) (1). [K poznaniyn faunui tatarskogho yarusa rr. Sukhonui i maloi severnoi dvinui.] Annu. Soc. paléont. russe 13 1949: 91-97 1 pl.

Poel, L. v. d. Faune malacologique du Hervien. Deuxième note. Bull. Inst. Sci. nat. Belg. 32 No. 19:1-23 l pl.

Poel, L. v. d. (1). Faune malacologique du Hervien Premier note. Bull. Inst. Sci. nat. Belg. 32 N 18: 1-12.

Polster, M. The exchange of phosphorous in shells of the aquatic snail *Physa acuta*. Experientia 12:310-311.

Pomeyrol, R. Sur les 'cardiidae' crétacés de la Nouvelle Caledonie. C.R. somm. Soc. Géol. Fr. No. 5:69.

Popov, G. I. The stratigraphical separation and comparison of Black-Sea and Caspian Quatenary deposits. C.R. Acad. Sci. URSS 101 1 1955: 143-146. [In Russian.]

Popov, G. I. (1). History of Manichok straits in connection with the stratigraphy of the Black-Sea and Caspian Quatenary deposits. Bull. Soc. Nat. Moscou geol. 30 2 1955: 31-49 4 figs. [In Russian.]

**Popov, G. I.** (2). Apsheron deposits of Ala-Dag. C.R. Acad. Sci. URSS **103 4** 1955: 681–683. [In Russian.]

Popov, Y. N. The ecology of the Tertiary Teredidae. C.R. Acad. Sci. URSS 100 4 1955: 789-791 3 figs. [In Russian.]

Popov, Y. N. (1). Asymmetrical splitting in the shell line in Triassic ammonites. C.R. Acad. Sci. URSS N.S. 95 2 1954: 381–383 2 figs. [In Russian.]

Popov, T. N. See Kiparisova, L. D.

Popescu, L. "Carii-de-Mare" (Teredinidae) și problema combaterii lor [Les Teredinidae et les moyens de les combattre.] Bul. Inst. Cercet. Pisc. Roman. 15 No. 4:69-76 f. 1-7. [French résumé.]

Pora, A. E. & Roşca, D. I. L'influence de l'extrait d'*Ecbalium elaterium* Rich. sur la coeur isolé de *Helix pomatia* L. Notul. biol. Bucureşti 5 1947: 199-204 2 figs. Pora, F. & Roşca, D. Le comportement aux variations de salinité. Dreissena polymorpha et Theodoxus fluviatilis. Stud. Cercet. Acad. Române. Fil. Cluj 6 1–2 1955: 27–41 6 figs. [French résumé.]

Postma, N. The lengthening reaction (Bingham Curve) of smooth muscles of hollow organs is characteristic for some gastropods only and is due to lengthening and tension reflexes. Int. Congr. Zool. 14 1953 (1956): 310-311.

Potts, W. T. W. The rate of urine production of *Anodonta cygnea*. J. exp. Biol. 31 1954:614-617.

Potts, W. T. W. (1). Osmotic and ionic regulation in a marine and freshwater lamellibranch. Abstr. Diss. Univ. Camb. 1952-53 1955: 32-33.

Pouderoyen, L. P. A new fossil *Natica* from the Netherlands. Basteria, Lisse 20 4/5:78.

Pouderoyen, L. P. See Altena, C. O. v. R. and others.

Prantl, F. & Ružička, B. Mnataia nov. gen., a new Devonian Pelecypod from Bohemia. Acta Mus. nat. Prag. 11B 5 1955: 1-17 1 pl. 9 figs. [English summary.]

Pratt, D. M. & Campbell, D. A. Environmental factors affecting growth in *Venus mercenaria*. Limnol. & Oceanogr. 11:2-1710 figs.

Pravoslavlev, P. A. Traces of fossil volcanic mud around Elton lake and the same formation in the lower Transvolga territory. Trud. Leningrad. Soc. Nat. 68 2 1951: 39-46 fig. [In Russian.]

Premvati. See Lal, M.

Price, L. Notes on small land snails found near Kaitaia, Northland [N.Z.]. Bull. Conch. Sect. Auck. Mus. 12:8-10.

Pronin, A. A. Namurian deposits of the western slope of the Middle Ural. C.R. Acad. Sci. URSS N.S. 90 6 1953:1123-1125. [In Russian.]

Pruvot-Fol, A. Revision de la Famille des Phyllidiadae. J. Conchyliol., Paris 96 2:55-80 text-figs.

Pruvot-Fol, A. (1). Les Arminiadae. (Pleurophyllidiadae ou Diphyllidiadae des anciens auteurs.) Bull. Mus. Hist. nat. Paris (2) 27:462-468 10 figs.

Pruvot-Fol, A. (2). Un Aeolidien nouveau des mers tropicales: Aeolidiopsis ransoni n.g., n.sp. Bull Mus. Hist. nat., Paris (2) 28:228-231 11 figs.

Pruvot-Fol, A. (3). Note à propos de l' Acteon. Bull. Soc. zool. Fr. 80 301-307 2 figs.

Pruvot-Fol, A. (4). Note sur deux Nudibranches attribués à la famille les Polyceradae. Bull. Soc. zool. Fr. 80:350-359 3 figs.

Purchon, R. D. The functional morphology of the rock-boring lamellibranch *Petricola pholadiformis* Lamarck. J. mar. Biol. Ass. U.K. **34** 1955: 257–278 16 figs.

Purchon, R. D. (1). A note on the biology of *Brechites penis* (L.). J. Linn. Soc. (Zool.) 43:43-54 8 figs.

Purchon, R. D. (2). A note on the biology of *Martesia striata* L. Proc. zool. Soc. Lond. 126: 245-258 7 figs.

Purchon, R. D. (3). The stomach in the Protobranchia and Septibranchia (Lamellibranchia). Proc. zool. Soc. Lond. 127: 511-525 f. 1-5.

Pushcharovski, Y. M. The tectonic structure of Priverkhoyansk ridge depression. Bull. Acad. Sci. U.S.S.R. geol. 5 1955: 37-53. [In Russian.]

Quick, H. E. A Review of "Sites of Infection" by A. Mozley. J. Conch. Lond. 24 3:104.

Quideau, Y. Contribution à la faune malacologique du Massif de Fontainebleau et de la Basse Volte de Loing, [List in text.] Trav. Nat. Loing No. 12 1955: 93-101.

Quideau, Y. (1). Malacofaune du Massif de Fontainebleau. Especes nouvelles observees en 1955. Bull. Ass. Nat. Loing 32 1 1955: 5–6.

Raabe, Z. Investigations on the parasitofauna of freshwater molluses in the brackish waters. Acta Parasit. polon. 4:375-406 7 figs. [English summary.]

Radoman, P. Contribution à la connaissance des Gastéropodes du Bassin d'Ohrid. Rec. Trav. Sta. Hydrobiol. Ohrid 3 1955: 25-39 41 figs. [French summary.] Răileanu, G. La présence d'un geure Antaracosia dans le permien inférieur de la zone de Svinita-Svinacea Mare, Comun. Acad. Republ. Pap. rom. 2 9-10 1952:581-585 3 figs. [French summarv.]

[1956]

Răileanu, G., Bădălută, A. & Pelin, M. L'étude de la faune d'Aptychus des calcaires du Jurassique supérieur de la zone de Svinita-Svinécea Mare (Barat). Ann. Univ. "C. J. Parhon" București: Ser Sci: Nat. No. 11:223—231.

Ralph, C. L. See Brown, F. A.

Rammelzhmeier E. S. [Mollynski iz mezozoïskikh ozernuikh Otlozheni khrebta Kara-Tan.] Trav. Inst. paléozool. Acad. Sci. URSS. 15 1 1948: 99–101 figs. [In Russian.]

Rangneker, P. V. Oxygen-carrying capacity of the blood of some crustaceans and molluses. Proc. Indian sci. Congr. Ass. 42nd 3 1955: 280–281

Ranson, G. Note sur Cypraea cicercula Linné 1758. Bull. Mus. Hist. nat., Paris (2) 26 1954: 612-619.

Ranson, G. (1). Les eaux apparemment vertes, mais en réalité lactescentes, du lagon de l'île Anaa (Tuamotu). [Moll. p. 31.] Bull. Lab. marit. Dinard No. 40 1954: 27–32 2 figs.

Ranson, G. (2). Observations sur l'anatomie de *Gyraulus costulatus* (Krauss), (Planorbe d'Afrique). Bull. Mus. Hist. nat., Paris (2) 27 1955: 374-379 3 figs.

Ranson, G. (3). Observations préliminaire sur quelques Planorbes américains. Bull. Mus. Hist. nat. Paris (2) 27 1955: 220-225 7 figs.

Ranson, G. (4). Observations sur l'appareil génital de Biomphalaria smithi Preston. Bull. Soc. zool. Fr. 81:35-43, 4 figs.

Ranson, M. G. & Medeiros, L. de. Observations sur l'anatomie de *Plan*orbis planorbis (Linné). Bull. Mus. Hist. nat. Paris (2) 27: 454-461 5 figs.

Rao, K. V. Seasonal gonad changes in the adult backwater oyster, Ostrea (Crassostrea) madrasensis Preston from Ennur near Madras. Proc. Indian Acad. Sci. 44B: 332–356 3 pls. 3 figs.

Rao, S. R. See Kathuria, J.

Rasetti, F. Internal shell structures in the middle Cambrian gastropod Scenella and the problematic genus Stenothecoides. J. Paleont. 28 1954: 59-66 pls. 11-12.

Rasmussen, L. B. The Marine Upper Miocene of South Jutland and its Molluscan Fauna. Danm. geol. Unders. (11) 81:1-166 10 pls. figs.

Ratnovski, I. See Alekseichik.

Raven, C. P. Effects of monovalent cations on the eggs of *Limnaea*. Pubbl. Staz. zool. Napoli 28:136-168 f. 1-3.

Ravera, O. See Corbella, C.

Ray, H. C. Mitres of Indian Waters (Mollusca, Gastropoda: family Mitridae). Mem. Indian. Mus. 14:1-72 3 pl.

Rayner, D. H. The lower Carboniferous rocks in the north of England: a review. [Lists pp. 285-287.] Proc. Yorks. geol. Soc. 28 4 1953: pp. 231-315 maps (pls.) xix, xx.

Redichkin, N. A. See Bankovski, V.

Rees, C. B. Continuous plankton records; the distribution of lamellibranch larvae in the North Sea, 1950–51. Hull Bull. Mar. Ecol. 4 27 1954: 21-46 4 pls. 13 figs.

Rees, W. J. Escaped Escargots. Nautilus, Philad. 68 3:90-94.

Rees, W. J. (1). The *Macrotritopus* problem. Int. Congr. Zool. 14 1953 (1956): 509.

Reiss, Z. See Avnimelech, M. and others.

Remmert, H. See Ohm, P.

Rendall, R. Mollusca Orcadensia. Proc. roy. Soc. Edinb. 66B: 131-201.

Requate, H. Die Jagdtiere in dem Nahrungsresten einiger frühgeschichtlicher Siedlungen in Scheswig-Holstein: Ein Beitrog zur Faunengeschichte des Landes. Schr. naturw. Ver. Schl.-Holst. 28 1:21—41.

Rerat, D. See Jullien, A.

Revilla, J. de la. See Castell, J.

Rey, L. L. Sobre el fomento de la producción pesquera de las aquas marinas y salobres confinadas; en el litoral de España y en el de sus territorios de Soberania y de Protectorado. Bol. Inst. esp. Oceanogr. No. 74:1-62 f. 1-37.

Reyment, R. A. Some new Upper Cretaceous ammonites from Nigeria. Colon. geol. & min. Res. 4 1954: 248-270 5 pls. 11 figs.

Reyment, R. (1). Über den bau von Spectoniceras versicolor (Trautschold) aus dem Neokom Russlands. Neues Jb Geol. Paläontol. Mh. Heft. 2 1956: 101–105 f. 1–4.

Ricci, E. Quelques remarques sur la sexualité chez la moule (Mytilus galloprovincialis Lmk.). Bull. Soc. linn. Provence 14 2 1943: 37–39.

Ricour, J. See Lemoine, M.

Riggs, C. D. & Webb, G. R. The mussel population of an area of loamy-sand bottom of Lake Texoma. Amer. midl. Nat. 56:197-203.

Rikhter, V. G. The history of karabogaz tongues of land. C.R. Acad. Sci. URSS 105 3 1955:543-546. [In Russian.]

Ripplinger, J. Application sur le ventricule d'Helix pomatia in situ, d'un procede de double inscription simultanu nouvelles observation au sujer de l'antomatisme ventriculaire de l'escargot. Ann. Sci. Univ. Besançon (2) Zool. et phys. No. 1 1954: 125–127.

Ripplinger, J. (1). Application à un organe donne, d'un systeme de double inscription simultanee des activites partielles respectives de territoires physiologiques ment interdependants. Ann. Sci. Univ. Besançon (2) Zool. et Phys. No. 1 1954: 129–131.

Ripplinger, J. See Binet, L. and others.

Ripplinger, J. See Jullien, A.

Ripplinger, J. R. See Jullien, A. and others.

Ripplinger, J. See Jullien A. & Cardot.

Ripplinger, J. See Jullien A. & Guyon.

Ripplinger, J. See Jullien, A. & Joly.

Ripplinger, J. & Joly, M. Sur une hypothèse de l'action des ions K+ et Ca++ sur l'activité du coeur de l'escargot (Helix pomatia). C.R. Soc. Biol., Paris 149 1955: 969-971.

Ripplinger, J. & Joly, M. (1). Etude des facteurs à l'origine des modifications de la réponse du myocarde d' Helix pomatia aux influx cardio-inhibiteurs. C.R. Soc. Biol. Paris 150 1956: 717-719.

Ripplinger, J. & Mercier, J. Etude de la production d'equivalent-acetylcholine au cours de diverses conditions experimentales, dans l'intimite du myocarde d'Helix pomatia sous l'influence de excitation du nerf visceral. Ann. Sci. Univ. Besançon (2) Zool. et phys. No. 1 1954:133-155 8 figs.

Risbec, J. Nudibranches du Viet-Nam. Arch. Mus. Hist. nat. Paris (6) 4 5-78 22 pls.

Risbec, J. (1). Etude anatomique de Naticidae de Nouvelle-Calédonie. J. Conchyliol. Paris 96 1:12-45 textfigs.

Rishi, R. The morpholy of the Golgi bodies with reference to secretion in the liver-cells of the slug, *Anadenus altivagus*, as seen under the phase-contrast microscope. Quart. J. micr. Sci. 97:171-176 l pl. 1 fig.

Ritchie, J. A Double Centenary— Two notable naturalists, Robert Jameson and Edward Forbes. Proc. roy. Soc. Edinb. 66B 1 (3): 29-57 Text-figs. and pls. i-iv.

Ritter, L. Ein Beitrag zur Molluskenfauna der Görlitzer Umgebung. Abh. Naturk. Mus. Görlitz 35 1:129-133.

Ritter, M. Les limaces et les escargots. Importance économique et moyens de lutte. Rev. Zool. agric. 54 4-6 1955 : 41-50 4 figs.

Rocandio, A. O. d. Z. See López, A. O. d. Z.

Roche, J. See Bouchilloux, S.

Roche, J. See Glahn, P. E. and others.

Roger, J. Paleobiogeographie des Invertébres fossiles de Madagascar. Mém. Inst. Si. Madagascar 1D 1945: 97-115.

Roger, J. See Darteville, E.

Rogers, G. E. See Lacy, D.

Rogue, A. L. The morphology of the cytoplasmic inclusions of the living spermatocytes, spermatids, and neurones of *Helix* as revealed by phase-contrast and interference microscopy and by suprarital dyes. J.R. micr. Soc. (3) 74 1955: 188-206 2 pls. 5 figs.

Rosca, D. I. See Pora, A.

Roşculescu-Soverth, E. Contributions àl'étude de la faune du pliocéne d'Olténie. Ann. Univ. "C. J. Parhon'. Bucureşti Ser. Sti. Nat. No. 9 1956 : 205-211 2 pls. 3 figs. [French résumé.]

Rosen, C. B. See Mayr, E.

Rosenkrantz, A. A large Velata from the lower Jurassic of East Greenland. Medd. dansk. geol. Foren. 13 2:79–84 f. 1–7.

Rossi, F. & Graziadei, P. Nouvelles contributions à la connaissance du système nerveux du tentacule des cephalopodes. II. Cordons ganglionnaires ou moelles périphériques du tentacule d'octopus. Acta anat. 26: 165-174 1 pl.

Rossi, F. et Graziadei, P. (1). Nouvelles contributions à la connaissance du système nerveux du tentacule des céphalopodes. III. Le ganglion sousacétabulaire et ses rapports avec une pseudo-articulation de la ventouse. Acta anat. 28: 206-216 f. 1-3.

Rost, H. & Soot-Ryen, T. Pelecypods from the Seychells Islands collected by Mr. W. V. Hasselberg. Acta Borealia (A) No. 8 1955: 1-23 3 figs.

Rothschild, Lord. Fertilization, London (Methuen) 1956, pp. ix + 170, 5 pls. 30 figs.

Roy, A. B. The steroid sulphatase of *Patella vulgata*. Biochem. J. **62**: 41-50 9 figs.

Rudkevich, M. Y.). [Some peculiarities in the history of Neogene of the western part of the Black Sea valley.] Bull. Soc. Nat. Moscou geol. 1 1955: 3-20 7 figs.

Rüefli, W. Ein Aufschluss von Aalénien-Schiefern im emgewickelten Glarner Flysch. Ecl. geol, helv. 49: 486–494 figs.

Rusconi, C. Fosiles ordovicios de la Quebrada de los Bueyes, (Mendoza). Rev. Mus. Hist. nat. Mendoza 9 1-2: 3-15 1 pl.

Rusconi, C. (1). Fosiles cambricos al sud de El Totoral Mendoza. Rev. Mus. Hist. nat. Mendoza 9 3-4:115-119 5 figs.

Rusconi, C. (2). Mares y organismos extinguidos de Mendoza. Rev. Mus. Hist. nat. Mendoza 9 3-4: 2-88 21 pls. 111 figs.

Russell, L. S. Nonmarine mollusca from the North Park formation of Saratoga Valley, Wyoming. J. Paleont. 30:1260-1263 4 text-figs.

Russell Hunter. See Hunter.

Rutsch, R. F. Remarks on I. Born's Index rerum naturalium Musei Caesarie Vindobonensis 1778. Nautilus, Philad. 69 3: 78-79.

Rutsch, R. F. & Salvador, A. Mollusks from the Cogollo and La Luna formations (Cretaceous) of the Chejendé area, western Venezuela. J. Paleont. 28 1954: 417-426 pl. 40 text-figs. 1-2.

Ruzhentzev, V. Asselsk stratum in Permian system. C.R. Acad. Sci. URSS N.S. 99 6 1954:1079–1082. [In Russian.]

Ruzhentzev, V. E. (1). On the family of Cyclolobidae Zittel. C.R. Acad. Sci. URSS 103 4 1955: 701-703 fig. [In Russian.]

Ruzhentzev, V. E. (2). The basic and stratigraphic complexes of Ammonites of the Permian. J. Acad. Sci. U.S.S.R. biol. 4 1955: 120–132.

Ruzhentzev, V. E. (3). Systematics and evolution of the families Pronoritidae Frech and Medlicottidae Karpinsky. Trav. Inst. paléozool. Acad. Sci. URSS 19 1949: 1–206 17 pls. figs.

Ruzhentzev, V. E. (4). Bio-stratigraphy of the Sakmara deposit in the Aktyubinsk Province of the Kazakh SSR. Trav. Inst. paléozool. Acad. Sci. URSS 42 1952: 1-90 6 pls. figs. [In Russian.]

Ruzhentzev, V. E. See Miklukho-Maklai, A.

Ružička, B. See Prantl, F.

Ryabikh, O. F. The problem of the chalk deposits surface of the Oskolsk layer. C.R. Acad. Sci. URSS 100 3. 1955: 541-544 2 figs. [In Russian.]

Sacarrão, G. P. A "linguagem" das cores nos cephalópodos. Naturalia  $65-6:14-20\ 3$  figs.

Sacarrão, G. da F. Quelques aspects sur l'origine et le développement du type d'oeil des Céphalopodes. Rev. Fac. Ciênc. Univ. Lisboa (2C) 4 1 1954:123– 158 21 pls. [English summary.] Sacarrão, G. da F. (1). Sur la genèse des chromatophores de *Tremoctopus*. Rev. Fac. Ciênc. Lisboa (2) 4C 2 1955: 295-304 5 pls.

Sacchi, C. F. Relations entre les groupements malacologiques terrestres et les conditions biogéographiques et biohistoriques des étangs roussillonnais. Vie et Milieu 5 1954: 529-564 1 fig.

Sacchi, C. F. (1). Contributo alla conoscenza dei popolamenti della piccoli isole Mediterrana. V.—Sulla malacofauna terrestre di Pantelleria. Atti Soc. ital. Sci. nat. 95:33–44 map. [English summary.]

Sacchi, C. F. (2). Il genere Opica (Stylomm.-Helicidae) in Sicilia. Annu. Ist. Zool. Univ. Napoli 8 (7): 1–6 2 figs. [English summary.]

Sacchi, C. F. (3). Ricerche su Euparypha arietina (Rossmaesster). I. Posiziome sistematica. Annu. Ist. Zool. Univ. Napoli 8 (8): 2 figs.

Safonova, T. P. See Larinova, E.

(Saidov, M. N.). [Meso-Caenozoic Continental deposits of the Dzhunger depression (basic features for stretigraphy.] Bull. Akad. Nauk. Geol. 10: 85-97 map. [In Russian.]

Saito, R. & Matsumoto, T. A new species of *Damesites* from the Cenomanian of Hokkaido, Japan. (Studies on the Cretaceous ammonites from Hokkaido and Saghalien—9). Trans. palaeont. Soc. Japan 22:191-194 1 fig.

Sakai, S. See Imai, T.

Salairanca, E. S. The effect of salt, vinegar, and heat on the coliforms in oysters. Philipp. J. Fish. 2 1954:93-107 4 figs.

Salmon, E. Catalogie des Strophocheilidae existant au Muséum National d'Histoire naturelle de Paris. J. Conchyliol., Paris 96 4:129-137.

Salvador, A. See Rutsch, R. F.

Salzmann, P.-E. Faunistisch-ökologische Untersuchungen über Süsswasser-Mollusken im Verlandungsgebiet am Südende des Ammersees. Veröff, zool. Staatsamml. Münch. 4:1-115 6 pls.

- Samodurov, V. I. Stratigraphy of the Mesozoic deposits of the Sir-Darya plain, Bull. Soc. Nat. Moscou geol. 30 3 1955: 39–56 5 figs. [In Russian.]
- Sánchez, J. A. Las biocenosis costeras de gastrópodos terrestres del litoral de Santander (nota previa). Bol. Soc. esp. Hist. nat. Secc. Biol. 54 1956; 33-34.
- Sandler, Y. M. & Glushko, V. V. Folded Silurian in North-Western part of Lwów province. C.R. Acad. Sci. URSS 103 4 1955: 685-688 map. [In Russian.]
- Sarkar, S. S. Révision des Ammonites dérontées du Crétacé inférieur du SE de la France. Mém. Soc. géol. Fr. Paléont. 72 1955 : 1-176 11 pts. 23 figs.
- Sarlet, L. Les mollusques de la région verviétoise. Rev. verviét. Hist. nat. 12 10-12 1955 : 100-103.
  - Sato, K. See Yagi, Y.
  - Sato, R. See Imai, T.
- Sato, T. Corrélation du Jurassique inférieur Japonais en basant sur les ammonites fossiles. J. Geol. Soc. Japan 62 No. 732: 490-503 f. 1-7.
- Sato, T. (1). Révision chronologique de la série de Karakuwa (Jurassique moyen). Jap. J. Geol. Geogr. 27: 167-171 1 pl. 5 figs.
- Sauerzopf, F. Interessante Mollusken aus dem Burgenland, Bürgenland, Heimat. 18:30–32.
- Saumont, R. La dessication, comme procédé permettant la mesure en vraie grandeur des phénomènes électriques du nerf. C.R. Acad. Sci. Paris 240 1955: 232-234 2 figs.
- Savilov, A. I. The size and its variation in the invertebrates of the White Sea, Mytilus eclatis Mya areurria, Balams Calanoides. Trud. Inst. Okeanol. 7 1953: 198–258 17 diagr. [In Russian.]
- Savilov, A. I. (1). Comparison of the size of *Mytilus edalis* in White and Okhotsk seas. Trud. Inst. Okeanol. 11 1954: 246–257 6 graphs. [In Russian.]
- Sawada, N. Artificial activation of the eggs of the bivalve, Caecella chinensis. Mem. Ehime Univ. (2B) 2 1955: 223-228 2 figs.

- Saxena, B. B. Excretory constituents of the blood of *Pila globosa* (Swainson). Arch. internat. Physiol. **64**: 578-582.
- Saxena, B. B. (1). Nutritive constituents of the blood of *Pila globosa* (Swainson). Naturwissenschaften **43** 9:201-202.
- Saxena, B. B. (2). Some observations on the ecology and behaviour of the common Indian apple-snail *Pila globosa* (Swainson). J. Bombay nat. Hist. Soc. 53:733-736.
- Sazonov, N. T. New data on systematics of upper-Jurassic ammonites. Bull. Soc. Nat. Moscou 30 4 1955:117. [In Russian.]
- Sazonova, I. G. Stratigraphy of Aptian deposits of the central part of the Russian platform. Bull. Soc. Nat. Moscou geol. 29 4 1954: 97-100 fig. [In Russian.]
- Schäfer, W. Wirkungen der Benthos-Organismen auf den jungen Schichtverband. Senckenbergiana leth. 37 183–263 2 pls. 35 figs.
- Schalie, H. v. d. & Dundee, D. S. The morphology of *Pomatiopsis cincinnatiensis* (Lea), an amphibious prosobranch snail. Occ. Pap. Mus. Zool. Univ. Mich. No. 579, 1-17 pl. 1-7.
- Scheltema, R. S. & Truitt, R. V. The shipworm *Teredo navalis* in Maryland coastal waters. Ecology 37:841-843 4 figs.
- Schermer, E. Hydrobiologische Untersuchungen im Travegebiet. Forsch. geogr. Ges. Lübeck (2) No. 42 1950: 35–131 8 pls. maps.
- Schermer, E. (1). Die Molluskenfauna der Nebengewässer. Forsch. geogr. Ges. Lübeck (2) No. 43 1951: 7–89 figs. map.
- Schindewolf, O. H. Über Strenoceras und andere Dogger-Ammoniten. N. Jb. Geol. Paläont. B., 3 1953: 119–130 text-figs. 1–10.
- Schlesch, H. Produkter från molluskriket. Svensk Faun. Revy 17 1955: 30–38.
- Schlesch, H. (1). Sinistral Oxychilus cellarius (Müller) in Sweden. J. Conch. Lond. 24 4:138.
- Schlickum, W. R. Daudebardia brevipes im Bergirchen Land und im angrenzenden Sauerland. Arch. Mollusk. 84 4/6: 181–182.

Schlocker, J. Bonilla, M. G. & Imlay, R. W. Ammonite indicates Cretaceous age for part of Franciscan Group in San Francisco Bay area, California. Bull. Amer. Ass. Petrol. Geol. 38 11 1954: 2372–2381 pl. 1 text-figs. 1–2.

Schmandt, W. & Sleator, W., Jr. Deviations from all-or-none behaviour in a molluscan unstriated muscle: decremental conduction and augmentation of action potentials. J. cell. comp. Physiol. 46 1955: 439–475 1 pl. 10 figs.

Schmidt, H. A. Zur Abhängigkeit der Entwicklung von Gehäuseschnecken von Kalkgehalt des Bodens. Dargestellt bei Oxychilus draparnaldi. Arch. Mollusk. 84 4/6 1955: 167–177 textfigs. 1 2.

Schmidt, H. A. (1). Theba carthusiana O. F. Müller lebond in Mecklenburg. Arch. Fr. Naturg. Mecklenburg 2 1956: 366-372.

Schneider, E. Über das Auftreten von Ceratiten im Trochitenkalk des Saarlandes, Ann. Univ. Sarav. Sci. 5 1 1956: 17-19 1 pl.

Schneiderman, M. See Olivier, L.

Schouten, A. R. Landslakken. Naaktslakken. Wet. Meded. Ned. naturw. Ver. No. 10 1954:1–19 16 figs.

Schüller, L. Zum Gedenken an Friedrich Mahler. Mitt. naturw. Arbeitsgem: 5-6 Zool.: 1-2.

Schwarz, E. The shape of the prostate gland in geographical forms of *Biomphalaria alexandrina* Ehrenburg. Trans. R. Soc. trop. Med. Hyg. 48 6 1954:537-538. (See also p. 540.)

Schwetz, J. Sur le réexamen partiel de trois agglomérations congolaises, examinées six aus auparavant au point de vue molaw-schistomique: Bunia et Irumu (Haut-Ituri) et Kasenzi (lac Albert). Bull. Acad. Sci. colon. Brux. N.S. 2 6: 1084–1095 2 figs.

Schwetz, J. (1). The identification of snails and the mysterious schistosome vectors in the Gambia. Trans. Soc. trop. Med. Hyg. 50: 616-617.

Schwetz, J. (2). The effect of climate variations on pulmonate Gastropoda, compared with that of molluscarides. Trans. Soc. trop. Med. Hyg. 50:579–582 map.

Schwetz, J., Fort, M. & Baumann, H. Double probleme de l'infection bisexuée, d'un seul planorbe par des cercaires de Schistosoma rodhaini et hybridation de S. rodhaini et de S. mansoni. C.R. Soc. Biol. Paris 148 1956: 1507-1509 1 fig.

Schwetz, J., Fort, M. & Baumann, H. (1). Sur le problème d'une infection uni-ou bisexuée transmise par les cercaires de *Schistosoma* émiser par un seul mollusque. C.R. Soc. Biol., Paris 149: 2026–2028 figs.

Schwinck, I. Vergleich des Redox-Pigmentes aus Chromatophoren und Retina von Sepia officinalis mit Insektenpigmenten der Ömmochromgruppe. Verh. dtsch. zool. Ges. 19:71-75 1 fig.

Scott, M. I. H. Dos nuevos Melanidos del Alto Paraná. Neotropica 1 (3) 1954: 45-48 3 figs. [English summary.]

Scott, M. I. H. (1). Ulpia nuevo género de Gastropoda terrestre. Neotropica 1 (5) 1955: 65-68 4 figs.

Segal, E. Adaptive differences in water-holding capacity in an intertidal gastropod. Ecology 37:174-178 4 figs.

Segal, E. (1). Microgeographic variation as thermal acclimation in an intertidal molluse [Acmaea limatula]. Biol. Bull. Woods Hole 111: 129–152 8 figs.

Seilacher, A. Studien gur Palichnologie II. Die fossilen Ruhespuren (Cubichnia). Neues Jb. Geol. Paläont. 98 1 1953: 87–124 pls. 7–13 text-figs. 1–5.

Sellmer, G. P. A method for the separation of small bivalve molluscs from sediments. Ecology 37: 206.

Semaka, A. See Atanasiu, L.

Semikhetova, S. V. Namurian layer and its importance in stratigraphy of the U.S.S.R. Carboniferous. Bull. Soc. Nat. Moscou. geol. 30 (5) 1955: 75–104. [In Russian.]

Seneca, H. & Bergendahl, E. Toxicity of antibiotics to snails. Antibit. & Chemother, N.Y. 5 1955: 737-741.

Seneš, J. See Čechovič, V.

Seneš, J. (1). Helvétska slirova fauna z okolia Modiého Kameňa. (Lay faumits du schlier helvétien des environs de Moorý Kamen (Slovaquie du Sud) [List]. Geol. Sborn. Slovensk. Akad. 1 1950: 110–129. (Fr. & Russ. summaries) pl. 11.

Seneš, J. (2). Nouvelle travailles de faune helvétien en Solvaquie du Sud. [List.] Geol. Sborn. Slovensk. Akad. 2 1951:71-74 (Fr. & Russ. summ.) pl. 4.

Seneš, J. (3). Geologický profil neogénon východnéko Bulkarska. [List.] Geol. Sborn. Slovensk. Akad. 2 1951: 85-89, pls. 22-30.

Seneš, J. (4). Oligozan zwischen Šaby und Lučenec in der Südslowakei. [List]. Geol. Sborn. 3 1952: 103—133. (Russ. & Germ. summ.) pl. 19.

Sénesse, P. Les trois facies du Coniacien des Corbieres meridionales. Bull. mus. Soc. linn. Lyon 25: 161-168.

Senna, G. See Cadeo, G. C.

Sergeev, V. A. On stratigraphy and tectonics of the western coast, of Karabogaz bay. Trud. Leningr. Soc. Nat. 68 2 1951: 7-25 3 figs. [In Russian.]

Setteparsi, F. See Blanc, A. G and others.

Shardanov, A. N. See Khaim, V. E.

Sharman, M. Note on Capulus ungaricus (L.) J. mar. Biol. Ass. U.K. 35:445-450 3 figs.

Shatski, N. S. On phosphorite bearing deposits of Karagandrinsk Basin. Bull. Acad. Sci. URSS geol. 3 1955: 48-56. [In Russian.]

Shaw, R. F. The polymorphism of the Japanese Littleneck clam. Nautilus, Philad. 70 2:53-59 tables, graph, pls. 45.

Shibata, M. A Fossil Fauna from the Northern part of the Tanzawa massif. Trans. palaeont. Soc. Japan 23: 229-234 1 pl.

Shimanski, V. N. On the principles of separating of the largest systematic groups of Cephalopods. Bull. Soc. Nat. Moscou geol. 29 5, 1954: 98-99. [In Russian.]

Shimanski, V. N. (1). Revision of some groups of Cephalopods. Bull. Soc. Nat. Moscou geol. 1 1955: 98-99. [In Russian.]

Shimanski, V. N. & Erlanger, A. A. On the findings of the Triconic nautiloidea. Bull. Soc. Nat. Moscou geol. 30 3 1955 95-96. [In Russian.]

Shimanski, V. N. & Zhuravleva, F. A. Stratigraphic importance of Nautiloidei. Bull. Soc. Nat. Moscou geol. 31 (3):112–113. [In Russian.]

Shulga, P. L. Pelecypods of the Carboniferous in Galizia Volhymian depression and their stratigraphic significance. Bull. Acad. U.S.S.R. geol. 6 1954: 75-84. [In Russian.]

Shuto, T. Umbonünae from Miyazaki Group (Palaeontological study of the Tertiary Muyazaki Group I). Japan J. Geol. & Geogr. 27:47-66 1 pl., 8 figs.

Si, T. See Tchang-Si.

Sieber, R. Fossilführung und Molluskenfaunen in Quatär Österreich. Anz. Öst. Akad. Wiss. 93:37-42.

Sieber, R. (1). Systematische Übersicht der jungtertiären Bivalven des Wiener Beckens. Ann. naturh. Mus. Wien 60 1955: 169–201.

Sieh, P. K. See Tang. C.

Silberling, N. J. 'Trachyceras' zone in the upper Triassic of the Western United States. J. Paleont. 30:1147-1153 2 text-figs.

Silva, G. H. da. Contribution à la connaissance de la faune fossile de l'île de Porto Santo. Rev. Fac. Ciênc. Coimbra 25: 33–35.

Silva, G. H. da (1). La faune Miocène de l'île du Prince. Rev. Fac. Ciênc. Coimbra 25 : 36-60 7 pls.

Sinizin, N. M. The elements of the Palaeozoic relief in Tyan-Shan. Trud. Leningrad. Soc. Nat. 68 2 1951: 47-56 6 figs. [In Russian.]

Sivertsen, E. Blekkspout (squids). K. norske vidensk. Selsk. Årbok. 1954 1955: 5–15 4 figs. [English summary.]

Sleator, W. jr. See Schmandt, W.

Slentzov, M. M. The biology of Cephalopods of the Far Eastern seas and for the North Western part of the Pacific. Trud. Inst. Okeanol. 18 1955: 69-77. [In Russian.]

Smekhov, E. M. Sarisuïsk dislocations. C.R. Acad. Sci. URSS N.S. N.S. 92 6 1953:1209-1211. [In Russian.]

Smith, E. K. Notes and records from Stewart Island. Bull. Conch. Sect. Auck. Mus. 12: 2-4.

Smith, F. G. W. Shipworm, saboteurs of the sea. Nat. geogr. Mag. 110 4: 559-566 figs.

Smithers, S. R. On the ecology of Schistosome vectors in the Gambia, with evidence of their role in transmission. Trans. Soc. trop. Med. Hyg. 50: 354-365 2 pls map.

Sobanski, A. See Bigarella, J.

Solem, A. The helicoid cyclophorid mollusks of Mexico. Proc. Acad. nat. Sci. Philad. 108:41-59 2 pls.

Solem, A. (1). Non-marine mollusca from Galobra, Matto Grosso Brazil and a collection of South Brazilian Artemon. Notulae Naturae No. 287: 1-14 pl. 1.

Soot-Ryen, T. New records on the distribution of marine Mollusca in Northern Norway. Astarte No. 1 1951:1-11.

Soot-Ryen, T. See Rost, H.

Sordi, M. & Majidi, P. Osservazioni sui nudibranchi e gli Ascoglossi (Gasteropodi Opistobranchi) del litorale livornese. Boll. Pesca Piscic. Idrobiol. (N.S.11) 32: 235–245 2 pls.

Sornay, J. Nautile fossile à coquille anormale. Bull. Mus. Hist. nat. Paris (2) 27 1955: 260–261 figs.

Sornay, J. See David, L.

Southward, A. J. The population balance between limpets and seaweeds on wave-beaten rocky shores. Rep. Mar. biol. Sta. Pt. Erin No. 68, 1955, [1956]: 20-29.

Southward, A. J. See Orton, J.

Southward, A. J. & Crisp, D. J. The distribution of certain intertidal animals around the Irish coast. Proc. R. Irish Acad. 57B 1 1954:1-29 10 figs. maps and tables.

Southward, A. J. & Dodd, J. M. Studies on the biology of limpets. I. The late J. H. Orton's work on Patella. J. Mar. biol. Ass. U.K. 35: 145-147.

Southwood, T. R. E. See Johnson, C. G. and others.

Soyer, T. Sur une faunale malacologique terrestre de Biarritz. Bull. Centre Ét. Rech. Sci. Biarritz 11:51-52.

Sparks, D. W. The non-marine Mollusca of the Hoxne interglacial. Phil. Trans. 239B: 351-354.

Sperling, P. See Mahler, F.

Staadt, J. L. Observations on the subgeneric classification of *Calliostoma*. J. Conchyliol. Paris 96 3:119-120.

Staiger, H. Chromosomal dimorphism and structural heterozygosity in *Pur*pura lapillus (Prosobranchia) with relation to its ecology. Caryologia 6 Suppl. Atti. IX Congr. Internat. Genet. pt. 2 1954: 892.

Starmühlner, F. Beiträge zur Mikroanatomia und Histologie der Darmkanals einiger Opisthobranchier, I. S.B. öst. Akad. Wiss. Math. Nat. Kl. Abt. I. 165: 93–152 36 figs.

Starmühlner, F. (1). Zur Molluskenfauna des Felslitorals und submariner Höhlen am Capo di Sorrento (2 Teil). Ergebnisse der Österreichischen Tyrrhenia-Expedition 1952, Teil IV. Öst. zool. Z. 6: 631-713, 15 figs.

Starobinetz, E. Y. & Eventov, Y. S. The age of Paleogene in the midstream of Amu-Darya river. C.R. Acad. Sci. URSS 58 3 1948: 313-315. [In Russian.]

Stehepinsky, V. "Belemnites Cornuclianus" D'Orbigny (Hauterivien). Cahiers Géologiques 38:383-384.

Stefaneuks, A. & Makhnach, A. S. Stratigraphy of the Devonian deposits and the age of the salt-bearing layer of the Pripet bend. J. Acad. Sci. Beloruss. Repub. 2 1955: 87-100. [In Russian.]

Stein, J. E. & Mackin, J. G. A study of the nature of pigment cells of oysters and the relation of their numbers to the fungus disease caused by *Dermocystidium marinum*. Texas J. Sci. 7 4 1955: 422-429 1 fig.

Steinberg, J. E. The pelagic nudibranch, Cephalopyge trematoides (Chun, 1889), in New South Wales, with a note on other species in this genus. Proc. Linn. Soc. N.S.W. 81:184-1928 figs. 1 pl.

Stephenson, L. W. A new bivalve genus and species from the Ripley formation [Cretaceous] of Mississippi. J. Paleont. 30:752-753 l text-fig.

Stephenson, L. W. (1). Fossils from the Eutaw Formation Chattahoochee River Region, Alabama-Georgia. Prof. Pap. U.S. geol. Surv. 274-J: 223-250 8 pls. fig.

Stephenson, W. See Endean and other.

Steusloff, U. Neue Beobachtungen und Erkenntrisse über Flora (Potamogeton vaginatus; Armeria inverseni; Moose; Zwergweiden), Fauna (Mollusken; Ovibos) und Klimageschichte (Allerödphase) des Würnperiglasiales in der Niederterrase der Emscher und der Lippe. Abh. Landesmus. Westf. 14 2 1951: 8-45.

Stevanović, P. M. Faunistische neuigkeiten aus den unteren Congerianschichten von Donja Tuzla in N.E. Bosnien—mit einer rückschau auf die Sog. Karagač-Endemiten. Ann. Géol. Penin. Balkan 21 1953: 25–28.

Stinson, N. See Gresson, R.

**Stock, J. H.** Landslakken. Huisjesslakken, deel 1. Wet. Meded. Ned. naturk. Ver. No. 5 1953:1-15 51 figs. No. 7 1953:1-28 46 figs.

Stolyarov, V. P. Parasitic fauna of fishes in Ribinsk water-reservoir. Trud. Leningrad Soc. Nat. 21 4 1952: 261–285. [In Russian.]

Stratton, L. W. The Mollusca of Hertfordshire. Trans. Herts. nat. Hist. Soc. 24 1955; 81-118.

Stratton, L. W. (1). Report on Mollusca. Trans. Herts. nat. Hist. Soc. 24 197.

Stratton, L. W. (2). The Mollusca of the Malham area (Presidential Address). J. Conch. Lond. 24 4:111-138. Lists and text maps.

Stratton, L. W. (3). Comments on introduced land molluscs of New Zealand. Bull. Conch. Sect. Auck. Mus. 12:11-12.

Strausz, L. Adatok Felsőoligocén Cerithiumfélék változékonysaga. Foldt. Kozl. 86: 274–283 1 pl. (French summary pp. 282–283).

Strausz, L. (1). Sur les Cerithidae de l'Oligocène Superieur. Bull. Hungarian Geol. Soc. Köt 86 3:274–283, plate.

Strinati, P. See Aellen, V.

Sukharevich, P. M. New data on the stratigraphy of Jurassic deposits in the south western part of the depression adjoining the Black Sea. C.R. Acad. Sci. URSS 111:681-684. [In Russian.]

Sukhova, A. I. Some representatives of the family Valencienniidae from the Pliocene of the Black Sea basin. Bull. Soc. Nat. Moscou geol. 30 3 1955: 93. [In Russian.]

Sultanaev, A. A. Stratigraphy of the upper-Paleozoic of Kolvu-vishersk Land. C.R. Acad. Sci. URSS N.S. 98 2 1954: 257–258. [In Russian.]

Sultanov, K. M. Oysters of the Tarkan horizon of Asserbaidzhan. C.R. Acad. Sci. URSS 100 3 1955: 547-549. [In Russian.]

Suzuki, K. Development of the fossil non-marine mollusca faunas in Eastern Asia. Jap. J. Geol. & Geogr. 21 1949: 91-133.

Svagrovský, J. Cerithium (Ptychocerithium) procrenatum Sacco aus dem Tortonien der Ostslawakei. Geol. Sbornik Slov. Akad. Bratislavia 7 3-4: 311-320 1 fig.

Sylvester-Bradley, P. C. The superspecies. Syst. Zool. 3 1954:145-146.

Takayanagi, F. See Asano, M.

Takemura, Y. & Okutani, K. Boring animals and their damages inflicted on the shells of silver-lip pearl oyster [Pinctada maxima] collected from the Arafura Sea. Bull. Tokai Fish. Res. Lab. 14:21-29. [English summary.]

Taki, I. Anatomical study on Japanese Epitoniidae (I) *Epitonium, Amaea* and *Papyriscala*. Bull. nat. Sci. Mus. Tokyo 3:71-79 3 pls.

Taki, I. & Habe, T. Pholadidae in Japan Pholadidae, Glauconomidae, Fimbrüdae. Illustrated Catalogue of Japanese Shells. 2 2 Sep. 1955: 7–16 pl. 2 & 3 text-figs.

Talent, J. A. Devonian brachiopods and pelecypods of the Buchan caves limestone, Victoria. Proc. roy. Soc. Vict. N.S. 68: 1-56 5 pls. 15 figs.

Talent, J. A. & Philip, G. M. Siluro-Devonian Mollusca from Marble Creek, Thomson River, Victoria. Proc. roy. Soc. Vict. N.S. 68: 57-71 2 pls. 4 figs. Talmadge, R. Notes on an Ecological Study [Halictis]. Mins. Conch. Club S. Calif. 154 Jan. 1956: 6-7.

Talmadge, R. (1). Haliotis varia. Mins. Conch. Club S. Calif. 161:3-7.

Talmadge, R. (2). Haliotis pustulata. Mins. Conch. Club S. Calif. 164:56.

Tanaka, J. Anatomical observations on *Anomia cytaeum* Gray. Bull. mar. biol. Sta. Asamushi 7 2-4 1955:111-119 5 figs.

Tang, C. C., Chow, C. C., Wang, P. C., Sieh, P. K. & Chow, S. L. Epidemiology of Schistosomiasis japonica in Futsing, Fukien Province. Peking nat. Hist. Bull. 19 2-3 1951: 225-247.

Tauc, L. Divers aspects de l'activité électrique spontanée de la cellule nerveuse du ganglion abdominal de l' Aplysia. C.R. Acad. Sci. Paris 240 1955 : 672–674.

Tauc, L. (1). Activités électriques fractionnées observés dans des cellules ganglionnaires de l'Escargot (*Helix* pomatia). C.R. Acad. Sci. Paris 241 1955: 1070-1073 figs.

Tavolga, W. N. The Old Man of the Sea, *Octopus vulgaris*. Biol. Rev. City Coll. N.Y. 15 1 1953: 3-5.

Coll. N.Y. **15** 1 1953 : 3-5. **Taylor, B. J. R.** See Kerkut, G. A.

Taylor, M. Variation of the shells of Murex tribulus from the Great Bitter Lake, Egypt. Nature, Lond. 174 1954: 1111-1112 1 fig.

Taylor, S. R. See McKerrow, W.

Tchang, Si, Tsi, Chung-yen & Li, Kie-min. Les tarets des côtes du nord de la Chine et leurs variations morphologiques. Acta zool. sinica 7 (1) 1955: 1-16 4 pls. [French résumé.]

Tchang-Si, Tsi, Chung-yen & Li, Kie-min (1). Recherches sur la reproduction et la croissance chez un pétoncle courestible—Chlamys farreri (Jones & Preston). Acta zool. sinica 8:235-253, 4 pls. [French résumé.]

Teesdale, C. Freshwater molluses in the coast province of Kenya with notes on an indigenous plant and its possible use in the control of bilharsia. E. Afr. med. J. 31 1954: 351-365.

Termier, G. Gasteropodes du Crétacé supérieur dans le sud-ouest de la France (groupe II). Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 : 323-382 74 figs. Teskey, M. C. Ventridens intertextus eutropis lives here [Cheatham Co., Tennessee.] Nautilus Philad. 69 4: 123-124.

Thiéblemont, M. Utilisation en Morbihan du collecteur-carton en vue du captage du naissain (Ostrea edulis). [Résumé]. Rapp. Cons. Explor. Mer 140 3: 23.

Thompson, S. Y. See Fisher, L. R.

Thorarinsson, S. The Nucella shore line at Húnafloi in the light of tephrochronological and radiacarbon datings. Náttúrufraedinguriun 25 3 1955:172–186 2 figs. [English summary.]

Thorpe, W. H. Learning and instinct in animals, London (Methuen), pp. viii + 493 9 pls. 71 figs.

Threadgold, L. See Gresson, R.

Tiburtius, G. See Bigarella, J. J.

Ticsa, I. See Velluda, C.

Tikhomirov, S. V. Devonian of Pripyatsk depression. Bull. Soc. Nat. Moscou geol. 30 6 1955: 93.

Tintant, H. & Joly, T. Observations sur la stratigraphie du Bathonien en Côte-d'Or. Bull. sci. Bourgogne 15 1955: 25-38 1 pl.

Tintant, H. & Larcher, C. Nouvelles observations sur l'Oxfordien aux environs de Dijon. Bull. sci. Bourgogne 16: 165-171 1 fig.

Tólp, Ö. Über das Benthos des Flusses Emajõgi. Loodus. Seltsi Aust. 49: 143–160 2 figs. [German summary p. 17.]

Tongiosgi, E. See Blanc, A. C. and others.

Trochon, P. Étude sur la reproduction de l'huître plate Ostrea edulis dans les claires de la région Mareunes-Oléron. Rapp. Cons. Explor. Mer 140 3: 14-16.

Tröger, K.-A. Über die Kreideablagerungen des Planenschen Grundes. Jb. Staatl. Mus. Min. Geol. 2:22–124 10 pls. figs.

Trousil-Linhart, W. Entwicklung der Schlammschuecke. Mikrokosmos 46 4:91-93 3 figs.

Trouve, P. See Mongin, D.

Truitt, R. V. See Scheltema, R. S.

Trümpy, R. Notizen zur mesozoischen Fauna der inner-schweizerischen Klippen (1-11). Ecl. geol. helv. 49: 573-591 2 pls. figs.

Trümpy, R. See Lemoine, M.

Tsi, Chung-yen. See Tchang-Si.

Tsuchiya, Y. & Makuta, T. The occurrence of cobalt in marine products. Tohoku J. agric. Res. 2 1951:113-117.

Tsujii, T. See Okada, Y.

Tsujimura, T. See Yagi, Y.

Tsukuda, H. See Ohsawa, W.

Tsuzuki, K. See Ishida, S.

Tsuzuki, K. See Ishida and others.

Tuaev, N. P. Lower Cretaceous of adjacent Dzhungaria. C.R. Acad. Sci. URSS 100 2 1955:351-354. [In Russian.]

Tuchkov, I. I. Fauna morskogho reta Severo-vostoka Azii. Annu. Soc. paléont. russe 15:177–221 5 pls.

Tuchkov, I. I. (1). The fauna of Pseudomonotis of the Noriïsk layer in north-eastern part of Siberia. C.R. Acad. Sci. URSS 104 4 1955: 608-610 Pl. [In Russian.]

Tuchkov, I. See Koshelkina, Z.

Turishchev, I. E. Lower Cretaceous deposits of South-eastern part of the Mongolian Republic. C.R. Acad. Sci. URSS N.S. 99 (3) 1954: 445-448. [In Russian.]

Turner, J. S. Nautiloid nomenclature: Coelonautilus and Cyrtoceras. Proc. Leeds phil. lit. Soc. Sci. Sect. 6 4 1954: 243–250 2 pls.

Turner, R. D. Notes on Xylophaga washingtona Bartsch and on the genus. Nautilus, Philad. 70 1:10-12.

Turner, R. D. (1). The Eastern Pacific marine mollusks described by C. B. Adams. Occ. Pprs. Mollusks Harvard Univ. Cambridge 2 20: 21-135 pls. 5-20.

Turner, R. D. (2). Additions to the Western Atlantic Marine Mollusks described by C. B. Adams. Occ. Pprs. Harvard Univ. Cambridge 2 20:136 pl. 21.

Turner, R. D. See Clench, W. J.

Twarog, B. M. Responses of molluscan smooth muscle to acetylcholine and 5-hydroxytryptamine. J. cell. comp. Physiol. 44 1954:141-163 5 figs.

Unklesbay, A. G. Distribution of American Pennsylvanian cephalopods. J. Paleont. 28 1 1954: 84-95 text-figs. 1-3.

Unklesbay, A. G. (1). Nautiloids from the Tanyard formation of central Texas. J. Paleont. 28 5 1954: 637-656 pls. 68-71.

[1956]

Uozumi, S. & Fujie, T. The sandpipe, created by the pelecypods: Platyodon nipponica n. sp. and Pholadidea (Penitella) kamakurensis (Yokoyama). J. Fac. Sci. Hokkaido Univ. (4) 9: 351-369 5 pls. 7 figs.

Ursin, E. Distribution and growth of the Queen, *Chlamys opercularis* (Lamellibranchiata), in Danish and Faroese waters. Medd. Danm. Fisk. Havundersøg. N.S. 1 Nr. 13:1–32 9 figs.

Usuki, I. Effects of some inhibitors of anaerobic glycolysis on the ciliary activity of the oyster gills. Sci. Rep. Tôhoku Univ. (4) 22 2:49-56 5 figs.

Usuki, I. (1). A comparison of the effects of cyanide and azide on the ciliary activity of the oyster gill [Ostrea gigas]. Sci. Rep. Tôhoku Univ. (4) 22 3:137-142 4 figs.

Usuki, I. (2). Effects of malonate and 2, 4-dinitrophenol on the ciliary activity of the oyster gill. Sci. Rep. Tôhoku Univ. (4) 22 3: 143-148 2 figs.

Usuki, I. & Okamura, N. Glycolytic intermediates in the oyster gills. Sci. Rep. Tôhoku Univ. (4) 22 4:225-232 1 fig.

Vágvölgyi, J. Malacfaunistical data. Ann. hist. nat. Mus. hung. N.S. 7: 451-453 1 fig.

Vágvölgyi, J. See Farkas, H.

Vaillant, F. Recharches sur la faune madicole (hygropétrique S.L.) de France, de loose et d'Afrique du Nord. Mém. Mus. Hist. nat. Paris 11A 1956: 1-258 6 pls. 35 figs.

Vannini, E. Comportamento dei nucleoli e delle ribonucleoproteine negli ovociti in accrescimento (Behaviour of nucleoli and ribonucleo-proteins in growing occytes). Ric. sci. 23 Suppl. 1953: 81-86 4 figs. [French, English & German summaries.]

Varentzov, I. M. The stratigraphy of the mid-Devonian deposits of Tnua. C.R. Acad. Sci. URSS 104 3 1955: 459-461. [In Russian.]

Vassoevich, N. B. The stratigraphy of the Mesozoic deposits of the flish zone in the South Eastern Caucasus. Trud. Leningrad. Soc. Nat. 68 2 1951: 168-207 2 figs. [In Russian.]

Velluda, C. C., Ticşa, I. & Csutak, V. La réaction du coeur isolé d'escargot (*Helix pomatia*) en présence de l'acétylcholine. Stud. Cercet. Acad. Române. Fil. Cluj 4 3–4 1953 : 205–219. [French résumé.]

Venkataraman, R. & Chari, S. T. Chemical investigations on the formation of pearls in the Indian pearl oyster (Margaritifera vulgaris). Proc. Indian sci. Congr. Ass. 43rd 4:29. (Abstract).

Venmans, L. W. A. C. Notes on Alycaeus. Proc. Malac. Soc. London 32 3:81-87 7 text-figs.

Venmans, L. W. A. C. (1). A new subspecies of *Pila africana* (Martens). J. Conch. London 24 4: 140-141 pl. 5.

Venmans, L. W. A. C. (2). Land and freshwater mollusks from the Dolomites. Atti Ist. veneto Cl. Sci. Mat. Nat. 112 1954:59-96 map.

Venmans, L. A. W. C. See Verdcourt, B.

Venugopalan, V. K. The amino acids of the crystalline style of Sanguinolaria diphos. J. Anim. Morph. Physiol. 3:109-113.

Verástegui, P. The Pelecypod genus Venericardia in the Paleocene and Eocene of western North America. Palaeontogr. Amer. 3 25 1953:5-112 pls. 1-22 2 charts.

Verdcourt, B. Notes on some land and freshwater mollusca from Ethiopia and Somalia (including some records from Kenya). Proc. Malac. Soc. London 32 1/2:72-79 text-figs.

Verdcourt, B. (1). Notes on the snails of North East Tanganyika Territory. 7. The variation of Gulella conradti (von Marts) and the description of Gulella intrusa, a closely allied new species. Proc. Malac. Soc. London 32 1/2:69-71 text-figs.

Verdcourt, B. & Venmani, L. A. W. C. A synopsis of the section Primigulella Pilsbry of Gulella Pfr. (Mollusca Streptaxidae. Basteria, Lisse 20 4/5: 65-77 text map and figs.

Verdcourt, B. See Hemming, C. F.

Veselinović, D. Die Ergebnisse der geologischen Untersuchungen im Gebeit von Vratarnica und M. Izvor in Ostserbien. [In Serbian with German summary.] Zborn. Radova geol. Inst. Beograd 4 1952: 121–143 (Germ. summ. p. 140) pls. 1–2 geol. map.

Vevers, H. G. See Kennedy, G. Y.

Vieille-Cessay, C. See Jullien, A.

Vigneaux, M. et Magne, A. Observations récentes sur le Burdigalien de Cestas (Gironde). [Lists.] C.R. Soc. géol. Fr. 15 1952: 351-3.

Viktorova, R. E. & Kovalevski, B. A. Aktchagil in Moldavia. C.R. Acad. Sci. URSS N.S. 94 4 1954: 737-740. [In Russian.]

Vilela, H. Contribution pour la connaissance de la reproduction des huîtres portugasis—Gryphaea angulata Lamck. Rev. Fac. Ciènc. Univ. Lisboa (2C) 4 1 1954: 187–211 5 figs.

Vilela, H. (1). Sobre ostreicultura. Naturalia 6 5-6:1-13 13 figs.

Vilela, H. (2) État de maturation sexuelle des huîtres du Tage, *Gryphaea* angulata Lmk. Rapp. Cons. Explor. Mer 140 3:32–34.

Villaluz, D. K. See Blanco, G.

Villatte, J. See Casteras, M.

Villela, G. G. Caratenoids of some Brazilian freshwater Gastropods of the genus *Pomacea*. Nature, Lond. 178:93.

Vinogredova, N. G. Material on the quantitative composition of the bottom fauna of some bays in Okhotsk and Behring seas. Trud. Inst. Okeanol. 9 1954:136-158 13 figs. [In Russian.]

Vistelius, A. B. Mineral associations and typical paragnesis of Apt-Senoman terrigene layer of Transcaspian. C.R. Acad. Sci. URSS N.S. 97 3 1957: 503-506 fig. [In Russian.]

Vitagliano-Tadini, G. Sul differenziamento sessuali dei goni in rapporto alle cellule follicolari. Caryologia 6 Suppl. Atti IX Congr. Internat. Genet. pt. 2 1954: 948–953 3 figs.

Vlerk, I. M. van der & Florschütz, F. The paleontological base of the subdivision of the Pleistocene in the Netherlands. Verh. Akad. Wet. Amst. Afd. Natuwrk. (1) 20 2 1953:1-58 4 pls map.

- Vokes, H. E. Notes on the Nucinellidae [Pelecypoda] with description of a new species from the Eocene of Oregon. J. Paleont. 30:652-671 2 figs.
- Vokes, H. E. (1). Notes on, and rectification of, pelecypod nomenclature. J. Paleont. 30: 762-765.
- Vologdin, A. G. Cambrian Solenopora and Mollusca of the northern Tyan-Shan. C.R. Acad. Sci. URSS. 105 2 1955: 354–356 3 figs. [In Russian.]
- Vorontzev, N. S. & Laptinskaya, E. S. New data about the age of *Inoceramus*, belonging to the group *Retrorsus* keys. C.R. Acad. Sci. URSS N.S. **96** 1 1954: 145-146. [In Russian.]
- Vorontzev, N. S. & Laptinskaya, E. S. (1). New data on lower Jurassic of Anaber region. C.R. Acad. Sci. URSS 100 5 1955: 955-956. [In Russian.]
- Voss, G. L. Checklist of the Cephalopods of Florida. Quart. J., Fla. Acad. Sci. 19: 274-282.
- Vostryakov, A. V. The northern border of the distribution of the marine Apsheron deposits. C.R. Acad. Sci. URSS 103 6 1955: 1081-1084 map. [In Russian.]
- Vyalov, O. S. New data on Paleogene oysters from Kashgar. C.R. Acad. Sci. URSS 101 5 1955: 921-923. [In Russian.]
- Vyalov, O. S. (1). Paleogene of the south east of Kopetdag. Bull. Soc. Nat. Moscou geol. 31 (5): 75-83. [In Russian.]
- Vyalov, O. S. & Pasternak, S. I. Novuie nakhodki inotzeramov v Zakarpatskom flishe. Geol. Sbornik 2-3: 203-209 1 pl.

Wada, K. See Watabe.

- **Wada, S. K.** Spawning in the Tridacnid clams. Jap. J. Zool. **11** N.Z. 1954: 273–285.
- Wada, S. K., Collier, J. R. & Dan, J. C. Studies on the acrosome V. An egg-membrane lysin from the acrosomes of Mytilus edulis spermatozoa. Exp. Cell Res. 10 1:168-180 11 figs.

Wagner, E. D. See Wong, L.

Wajdi, N. A quantitative survey of Bulinus snails in the winter season. Bull. endemic. Dis., Baghdad 1 3 1955: 257-264 2 figs.

Waldén, H. W. The land gastropoda of the vicinity of Stockholm. Ark. Zool. (2) 7 No. 21 1955: 391-448 text-figs. pl. i.

[1956]

- Waldén, H. W. (1). Notes on some Siberian slugs, especially on the type of Limax hyperboreus Westerlund. Ark. Zool. (2) 10 5:347-365 text-figs. & plate.
- Walliser, O. H. Chronologie des Lias  $\alpha_s$  zwischen Fildern und Klettgau (Arietenschichten, Südwestdeutschland). Neues Jb. Geol. Paläont. 103: 181–222 2 pls. 5 text-figs.
- Walne, P. R. Bacteria in experiments on rearing oyster larvae. Nature, Lond. 178:91.
- Walne, P. R. (1). The biology and distribution of the slipper-limpet Crepidula fornicata in Essex rivers: with notes on the distribution of the larger epi-benthic invertebrates. Fish. Invest. Lond. (2) 20 No. 6, 1-50 33 figs.
- Walne, P. R. (2). Observations on the oyster (Ostrea edulis) breeding experiments at Conway, 1939–1953. Rapp. Cons. Explor. Mer 140 3: 10–13.
- Walne, P. R. (3). Experimental rearing of the larvae of Ostrea edulis L. in the laboratory. Fish. Invest. Lond. (2) 20 No. 9:1-23 2 pls.

Wang, P. C. See Tang, C.

- Wangaard, F. F. The natural marine borer resistance of tropical American woods. Trop. Woods No. 105:38-43.
- Warren, P. Collecting notes from Leigh. Bull. Conch. Sect. Auck. Mus. 12:22-24.
- Watabe, N. Dahllite identified as a constituent of prodissoconch I of Pinctata martensii (Dunker). Science 124:630.

Watabe, N. See Okada, Y.

- Watabe, N. & Wada, K. On the shell structures of Japanese pearl oyster, *Pinctada martensii* (Dunker). Rep. Fac. Fish. Univ. Mic. 2: 227-231, 2 pls. 7 figs.
- Watson, H. Support for A. E. Ellis's application regarding the name *Unio* Philipsson, 1788 (Class Pelecypoda). Bull. zool. Nom. 12:308.

Waugh, G. D. See Cole, H. A.

Waugh, G. D. & Ansell, A. The effect on oyster spatfall, of controlling barnacle settlement with DDT. Ann. appl. Biol. 44:619-625 1 pl.

Wautier, V. & J. Le cancer au les invertébrés. Seconde partie. Reactions provoquees par les substances cancerigenes [Moll. pp. 109-111]. Bull. mens. Soc. Linn. Lyon 24 5 1955:107-123

Webb, G. R. See Riggs, C. D.

Webb, H. M. See Brown, F. A.

Webb, J. H. A new species of *Pecten* (*Chlamys*) from Ryukyu. Venus, Fukuyama, 19 1:22–24 text-figs.

Weel, P. B. van. Osmoregulation in *Tethys* (Aplysüdae, Mollusca). Preliminary note. Proc. Hawaii. Acad. Sci. No. 31:8–9.

Weerekoon, A. C. J. Studies on the biology of Loch Lomond. I. The benthos of Cluchentullich Bay. Ceylon J. Sci. 7C 1: 1-94 70 figs.

Weinbach, E. C. & Nolan, M. O. The effect of pentachlorophenol on the metabolism of the snail Australorbis glabratus. Exp. Parasit. 5:276-284 2 figs.

Wells, J. See Wells, M. J.

Wells, M. J. & J. Tactile discrimination and the behaviour of blind *Octopus*. Pubbl. Staz. zool. Napoli 28:94-126.

Welsh, J. H. Acetylcholine and the heart of *Venus mercenaria*: A study of molecular configuration and physiological activity. Int. Congr. Zool. 14 1953 (1956): 277.

Welsh, J. H. (1). Neurohormones of invertebrates. I. Cardio-regulators of *Cyprina* and *Buccinum*. J. Mar. biol. Ass. U.K. **35** 6:193-201 5 figs.

Wendt, H. L. Das Haupt der Medusa. Kosmos, Stuttgart 48 1952:15-21 10 figs.

Wenger, R. Über einige Aberrationen bei Muschelkalk-Ceratiten. Neues Jb. Geol. Paläont. 103: 223–232 2 pls. 1 text-fig.

Westermann, G. Phylogenie der Stephanocerataceae und Perisphinctaceae des Dogger. Neues Jb. Geol. Paläont. 103: 233–279 9 text-figs.

Westermann, G. (1). Monographie de Bajocien-Gattungen Sphaeroceras und Chondroceras (Ammonoidea). Beihft. Geol. Jahrb. 24: 1-125 14 pls. figs. Weyrauch, W. The genus Naesiotus, with descriptions of new species and notes on other Peruvian Bulimulidae. Proc. Acad. nat. Sci. Philad. 108: 1-17 l pl.

Weyrauch, W. (1). Two new species of Clausiliidae from Peru and Bolivia. Nautilus, Philad. 69 4:110-115 pl. 6 f. 2-3a.

Weyrauch, W. K. (2). Neue Landschnecken aus Peru. Arch. Molluskenk. 85 4/6:145-162 pl. 11.

Whitley, G. P. See McMichael, D. F.

Whitten, H. E. Some variations in the species, *Charopa coma* (Gray). Bull. Conch. Sect. Auck. Mus. 12:17-20 3 figs.

Wickstead, J. An unusual method of capturing prey by a cuttlefish. Nature, Lond. 178: 929.

Wilde, J. de. Koordinationen im gefässsystem von Octopus vulgaris L. Pubbl. Staz. zool. Napoli 28: 359–366. [English summary.]

Wilkins, G. L. A review of "Introducing Sea Shells" by R. Tucker Abbott. J. Conch. Lond. 24 4:148.

Williams, G. Fauna of Strangford Lough and neighbouring coasts. Proc. R. Irish Acad. 56B 1954: 29-133 3 pls.

Wilson, D. P. A contribution to the biology of *Ianthina janthina* (L.). J. mar. Biol. Ass. U.K. **35**: 291-305 1 pl. 2 figs.

Winsmer, T. G. See Major H.

Wirz, K. Contribution à l'étude des octopodes de profondeur I. Bathypolypus sponsalis P. & H. Fischer espècé commune dans la partie ouest de la Mediterranée. Vie de Milieu 6 1955: 129-147 12 figs.

Wittmann, O. Lur Stratigraphie und Bildungsgeschichte der Meeressandbildungen entlang der Rheintalflexus bei Lörrach. [Short lists in text]. Jber. oberrhein. geol. Ver. NF.33 1953: 44-77 text-figs. 1-5. [Map & graphs.]

Wong, L. W. & Wagner, E. D. Some effects of ultraviolet radiation on Oncomelania nosophora and Oncomelania quadrasi, snail intermediate hosts of Schistosoma japonicum. Trans. Amer. micr. Soc. 75: 204-210 7 figs.

Wood, B. J. A study of the colour variation in the slug Agriolimax reticulatus Mull. Durham Coll. nat. Hist. J. 1954-55 [1956]: 15-18.

Woolacott, L. Coral Dwellers. Proc. R. zool. Soc. N.S.W. 1953-54 1955: 79-81 pl. 2.

Woolacott, L. (1). Notes on Australian shells. Proc. R. zool. Soc. N.S.W. 1954-55 1956; 72-75 5 figs.

Worth, C. B. See Mackie, T. T. and others.

Wright, C. A. Bulinus (Pyrgophysa) forskalii (Ehrenberg) as a vector of Schistosoma haematobium. Nature, Lond. 177: 43.

Wright, C. A. (1). A note on the ecology of some molluscan intermediate hosts of African schistosomiasis. Ann. trop. Med. Parasit. 50:346-349.

Wright, C. A. (2). The generic names, *Biomphalaria* and *Planorbis*. Trans. Soc. trop. Med. Hyg. **50**: 509

Wright, C. A. (3). The anatomy of the six species of the molluscan genus Bulinus (Planorbidae) from Senegambia. Proc. Malac. Soc. Lond. 32 3: 88-104, 18 text-figs. & pls. 21.

Wright, C. W. Notes on Cretaceous ammonites.—III *Utaturiceras* gen. nov. and the Metoicoceratinae. Ann. Mag. nat. Hist. (12) 9:391-393.

Wright, C. W. (1). A propos du genre Jacobella (Résponse à M. Jean Mercier). C.R. Soc. géol. Fr. 11 1952 : 217.

Wurtz, C. B. Fresh-water mollusks and stream pollution. Nautilus, Philad. 69 3: 96–100.

Yagi, Y., Mishima, T., Tsujimura, T., Sato, K. & Egami, F. Recherches sur l'hémoglobine d' Anadara inflata (Reeve). I. Purification et propriétés. II. Acides aminés N-et C-terminaux. C.R. Soc. Biol., Paris 149: 2285-2291.

Yakovlev, N. N. Rodui Cyrtodontarca Yakovlew i Dozierella Newell. Annu. Soc. paléont. russe 15: 158–160 fig.

Yamaguchi, N. & Habe, T. Studies in Japanese slugs. [In Japanese & English résumé.] Venus, Fukuyama 18 4 1955 : 234-240.

Yamamoto, G. Habitats of spats of the scallop, *Pecten yessoensis* Jay, which turned to bottom life. Sci. Rep. Töhoku Univ. (4) 22:149-156 3 figs. Yonge, C. M. Giant clams. Discovery 16 4 1955: 154-158 6 figs.

Yonge, C. M. (1). Adaptation to rock boring in Botula and Lithophaga (Lamellibranchia, Mytilidae) with a discussion on the evolution of this habit. Quart. J. micr. Sci. 96 1955: 383-410 19 figs.

Yoshida, H. Early life history of useful bivalves in the Ariake Sea. J. Shimonoseki Coll. Fish. 6:115-122 4 figs. [English summary.]

Yoshii, G. See Okada, Y.

Young, J. Z. Visual responses by octopus to crabs and other figures before and after training. J. exp. Biol. 33: 709-729 11 figs.

Young, J. Z. See Boycott, B. B.

Yuki, R. See Imai, T.

Zacks, S. I. The cytochemistry of the amoebocytis and intestinal epithelium of *Venus mercenaria* (Lamellibranchiata), with remarks on a pigment resembling ceroid. Quart. J. micr. Sci. 96 1955: 57-71 2 figs.

Zakaria, H. Further study on the ecology of the intermediate host of Schistosoma haematobium, Bulinus truncatus Baylis. Bull. endemic Dis., Baghdad 1 2 1955: 123-155 4 figs.

Zänkert, A. Drei Grabmäler aus Perlmutter. Kosmos, Stuttgart 48 1952:524-525 2 figs.

Zárate, O. de. See López & Rocandio.

Zavattari, E. Malacofauna e schistosomiasi nel medio e basso Giuba [Italian Somaliland.] Riv. Parassit. 17 No. 4: 193-202 f. 1-8. [English summary.]

Zázvorka, V. Chlamys (Gigantopecten) latissima (Brocchi) aus dem Neogen Mährens. Čas. nár. Mus. 125 1956: 177–179 2 figs. [German summary.]

Zei, M. M. See Moncharmont Zei.

Zeiss, A. Hecticoceras und Reineckeia im Mittel-und Ober-Callorim von Blumberg (Südbaden). Abh. Bayer-Akad. Wiss. Math.-Nat. Kl. N.S. No. 80:1-1014 pls. 4 figs.

Zeissler, H. Pisidium vincentianum im Ehringsdorfer Pleistozän. Arch. Molluskenk. 85 4/6:197-198.

Zeukevitch, L. Neue Vertreter der Mittelmeer-fauna im Kaspischen Meer. Int. Congr. Zool. 14 1953 (1956):113–118.

Zhgenti, E. M. Deposits of Konsk stage, formed in normal condition of salinity of the sea. C.R. Acad. Sci. Moscou N.S. 96 1 1954: 159-160. [In Russian.]

Zhuravleva, F. A. See Shimanski, V. N.

Zilch, A. Die Typen und Typoide des Natur-Museums Senckenberg 15 Mollusca, Cyclophoridae, Cyclophorinae-Cyclophoreae (2). Arch. Mollusk. 84 4/6 1955: 183-210 pls. 13-15.

Zilch, A. (1). Die Typen und Typoide des Natur-Museums Senckenberg. 14: Mollusca, Viviparidae. Arch. Mollusk. 84 1/3 1955; 45-85 pls. 3-7.

Zilch, A. (2). Die Typen und Typoide des Natur-Museums Senckenberg, 18: Mollusca Cyclophoridae, Cyclophorinae (4). Arch. Mollusk. 85 4/6:171-195 pls. 12 13.

Zilch, A., Mertens, R. & Kraus, O. Comments on application by Joshua L. Baily, Jr., regarding the preservation of the generic name *Turbinella* Lamarck 1799, as the name for the sacred chank shell of India. Bull. zool. Nom. 12:183.

Zinovyev, M. S. See Makridin, V. Žlebnik, L. Triassic cephalopods from Peca. Razp. Geol. Poročila 3 1955: 216-219 1 pl.

## II.—SUBJECT INDEX

N.B.—A number in parenthesis following an author's name refers to the particular paper in Titles so numbered when more than one work by the same author is recorded.

## GENERAL LITERATURE

General Works and Treatises.—Mollusca of Portuguese overseas territories —General Introduction, Azevedo & Medeiros; Spanish Cretaceous Mollusca, J. Bataller (1); Gas exchange in mollusca, Bodrova & Krayukhin; Slugs colour figs., P. Bondesen; Persistant rhythmicity in Ostrea virginica and Venus mercenaria, F. A. Brown, Ralph & others; Shell disease in Portuguese oysters, Cole & Waugh; Observations on the benthonic and planktonic

relations in L. Maggiore, Italy, Corbella, Croce & Ravera; Blood pigments, E. Eliassen; Role of terrestrial molluscs in the decomposition of fallen leaves, E. Frömming (3); Shape and form of tracks made by Littorina littorea in different sediments, I. E. Gräf: Conversion of calcite to fluorite with regard to fossils, J. F. Grayson: A survey of the genus *Plecotrema* figs., B. Hubendick (3); Eudemicism in the snails of Jamaica, W. R. Hunter (1); Form and development in relation to mode of life in lamellibranchs, W. Kühnelt: Palaeozoic and Pleistocene fossils of Ohio, U.S.A., La Rocque & Marple; A preliminary note on the Sarmatian fauna of Algeria, O. Mongin; Molluses from a fish-weir excavated at Boston, U.S.A., Clench & Nelson (separate entries); On the Phylogeny of the Umboniinae, K. Oyama (2); Mollusca in metamorphic complexes, Pavlovski & Frolova; Effects of Mollusca and other organisms on newly forming sediments, W. Schäfer; Slugs, figs., R. Schouten: Anatomy Milax sowerbii, A. R. Schouten; Principles for separating the largest systematic groups of Cephalopoda, Shimanski; Revision of some cephalopods, Shimanski (1); The value of Nautiloides for Stratigraphy, Shimanski & Zhuravleva; Biology of Cephalopods in the Far East and the N.W. Pacific, Slentzov; Hearing & instinct in molluses, W. Thorpe: Fauna of shallow running waters in France, Corsica and North Africa, F. Vaillant; Age of Inoceramus of the group Retrorsus keys, Vorontzev & Laptinskaya; Habitats & spat mortality of Pecten yessoensis, G. Yamamoto; Pigment in Venus mer-cenaria, S. I. Zacks; A fish covered with nacre after being trapped in the shell of Meleagrina margaritifera, A. Zänkert.

Monographs.—Monograph of the Strophocheilidae, J. C. Bequaert; Archibenthal Mollusca of New Zealand, R. K. Dell; Linnaean species of Buccinium and Strombus, H. Dodge; Revision of species described in "Monographie des Ammonites" by P. Reynès, D. Donovan; Cardiidae from the Russian Pliocene, A. Eberzin; Pulmonata and Prosobranchia of New Caledonia, A. Franc (2); Biology of mid-European Pulmonata and Prosobranchia, E.

Fromming (4); Roumanian Opisthobranchia and Prosobranchia, A. V. Grossu (3); Post-Triassic nautiloid genera, B. Kummel (1); Rissoinidae and Rissoidae from Northern Australia, Christmas Island and New Guinea, C. F. Laseron; Cylichna (Gastropoda: Tectibranchia), H. Lemche; Pliocene molluses of Adelaide Plains, Australia, N. H. Ludbrook; Revision of 15 species of Rissoa, P. Mars (2); New Zealand species of Baryspira (Olividae), O. P. Olson; Revision of the genus Olivella, A. Olsson; Mitridae from Indian waters, H. Ray; Nudibranchia of Viet-Nam, J. Risbec; Unrolled ammonites of the lower Cretaceous of S.E. France, S. Sarkar.

History.—Work of Coste in the development of oyster culture, M. Boury; Review of Linnaean species of Buccinium and Strombus, H. Dodge; History of New South Wales shells. I. Captain Cook and his associates, T. Iredale; History of Japanese Conchology, T. Kanamaru; History of Japanese Conchology figs., T. Kanamaru (1); History of Japanese Conchology (36), T. Kanamaru (2); History of Japanese conchology. HI Palaeontology, K. Oyama (3).

Biography.—Calvin Goodrich (notice of death), Anon (2); Henry Edward Crampton 1875–1956, H. B. Baker (4); Raymond C. Rush, M. D. 1875–1954, R. W. Dexter; L. de Folin (1817–1896), B. S. Kisch; Henry Edward Crampton 1875–1956, Y. Kondo (1); Robert Jameson centenary, Portraits, J. Ritchie; Edward Forbes centenary portraits, J. Ritchie; Friedrich Mahler, L. Schüller.

Bibliography.—Rejection for nomenclatorial purposes of S. A. Renier's "Tavole per servire alle classificazione e connescenza degli Animali", Opinion 427; "Synopsis of the species of airbreathing Mollusks of North America" W. G. Binney, 1863, rejected as an unpublished proof, Opinion 432; "Museum Boltenianum" published anonymously in 1798, to be attributed to Peter F. Röding, Direction 48; Works relative to fossil Ammonites of the Pacific area, R. S. Allan; Paleontological, early work on Rudistes by Lapeirouse, G. Astre; Mollusca of Portuguese

overseas territories, Azevedo & Medeicas; Dates of the Nautilus, H. B. Baker (2); Relating to Spanish Cretaceous Mollusca, J. Bataller (1); Works relating to Cretaceous Belemnites from Greenland, T. Birkelund; On C. T. Simpson's paper containing the figures of Uniouidae of B. H. & S. H. Wright, W. J. Clench (3); Literature relative to New Zealand deep-water molluscs, R. K. Dell: Works relative to classification of Strombus and Buccinium with reference to Linnaean species, H. Dodge; Revision of species described in "Monographic des Ammonites" by P. Reynès, D. Donovan; Works relating to Prosobranchs and Pulmonates of New Caledonia, A. Franc (2); Works relating to mid-European Pulmonata and Prosobranchia, E. Fromming (4); Conchological works of L. de Folin, B. S. Kisch; List of papers by T. Iredale with index of his new scientific names, McMichael & Whitley; Bibliography relative to cretaceous molluscs, I. G. Mahmoud; Bibliography relative to Miocene Pelecypoda from Russia, Merklin & Nevesskaya; Works relative to Cretaceous American ammonites. E. Packard; A review of "Sites of Infection" by A. Mozley, H. E. Quick; Born's "Index Rerum naturalium Musei Caesarei Vindobonensis" 1778, R. F. Rutsch; Bibliography relating to unrolled ammonites of the Lower Cretaceous of S. E. France, S. Sarkar; Bibliography relative to the fauna of the Upper Triassic of the Western United States, N. V. Sibberling; Works relative to fossil molluscan faunas of E. Asia, K. Suzuki; Relating to cretaceous Pacific marine moll., described by C. B. Adams, R. D. Turner (1); "Introducing Sea Shells" a review, G. L. Wilkins.

Taxonomy.—Fissurellidae, Anon (5;)
Trochidae, Anon (6); Bursidae, Cymatidae, Strombidae, Fasciolariidae, Anon (7); Epitoniidae, Anon (8) (9); On Taxonomy, Anon (11); Use of chromosome numbers in classification. Australorbis glabratus olivaceus, Planobarius corneus and P. metidjensi, Azevedo & Conçalves; Classification p. 44 of certain Bulumoid and Lymaeid snails, Azevedo & Medeiros; Addendum to Families of Pulmonata, H. B. Baker (1); Systematic position of Orthoceras, Z. Balashov; Preliminary revision of the

Loricata, J. Bergenhayn; Morphology of the Loricata, J. Bergenhayn; Study of the systematics of Amphineura, C. Boettger (1); Two new genera of the family Pharciceratidae, B. Bogoslovski (1); New family of Ammonoidae, B. Bogoslovski (2); Taxonomic characters in Mollusca, J. Q. Burch; The genus in systematics, J. Q. Burch (1); Taxonomic characters, J. Q. Burch (2); Characters to study in Classification, J. Q. Burch (4); More random notes, J. Q. Burch (5); Class interpretation of the genus Ptychomya Agassiz figs., A. Chavan; Key to common genera of plantonic Gastropoda p. 265, C. Davis: Key to recent New Zealand species of Cuspidaria, R. K. Dell p. 37; New Zealand species of Limnaea, Myxas and Physastra, R. K. Dell (1); Types of mollusks described by F. C. Baker, D. S. Franzen; Systematic note on Iberus, Garcia San Nicolas Alvarado; Roumanian Opisthobranchia and Prosobranchia, A. V. Grossu (3); Problem of oyster taxonomy, Gunter; Systematic positions of Usatricolia compta (Gld.), Rictaxis punctocoelatus (Cpr.), Nodilittorina pyramidalis Q. and G.: figs., T. Habe (9); Revision of Javanese freshwater gastropods, E. v. B. Jutting; Anatomic key to the slugs of Luxembourg, M. Lamesch; Review of Mytilidae from N.S.W. Australia, C. Laseron (1); Note on taxonomy of *Ianthina*, **D. Laursen**; Key to Czechoslovakian Mollusca, V. Lozek (2); New Orders, families and subfamilies in the Class Coniconchia, Lyashenko; Revision of 15 species of Rissoa, P. Mars (2); Neosuccinea new genus for some species of Succinea, Matekin, P. V.; Systematics of land mollusca of Tyan-Shan, P. Matekin (2); The classification of Paleozoic ammonoids, Miller & Furnish; Taxonomic significance of gaps in pelecypod morphology, D. Nicol (2); Key to subgenera of Olivella on shell characters, p. 166, A. Olsson; Genetical approach to the systematics of the Brazilian Planorbidae, W. L. Paraense; Key to the Brazilian species of Australorbis p. 285, Paraense & Deslandes (3); Type species in the Whitfield collection at the University of California, Peck MacFarland; Review of Phyllidiadae, figs., A. Pruvot-Fol: Problems of classification of Macrotritopus, W. J. Rees (1); Systematics of the ammonite

families Pronoritidae and Medlicottidae, V. Ruzhentzev (3); Systematic position of Euparyha arietina Rossm. determined as E. pisana ssp. arietina, C. Sacchi (3): Systematics of upper-Jurassic Ammonites, Sazonov; Key to identification of slugs, A. R. Schouten; New families of the Triassic Nautiloidea, Shimanski & Erlanger: Subgenera of Calliostoma, J. L. Staadt: Systematics of Belemnites cornuclianus, V. Stehepinsky; Validity of the species of Cephalopyge, J. Steinberg; Key to identification of Genera of land molluscs (with figures) of the Netherlands, S. H. Stock: The Superspecies, Sylvester-Bradley; Review of Primigulella Pilsbry, Verdcourt & Venmans; Taxonomy of the ammonite family Arietitidae, O. Walliser.

Nomenclature.—Opinions; New name Caspicardium for Cerdium, Astafyeva; Land operculate names, H. B. Baker; Family nomenclature of Pulmonata. H. B. Baker (3); Nomenclature of some Scandinavian Unionacea, T. Brauder; Jacobella and Hoplites, M. Breistroffer; Proposed addition to the Official List, of the generic name Chama Linn, 1758, L. R. Cox (3); Morphology of Teredo shells with regard to classification, P. Deschamps; Problems of naming Carboniferous non-marine lamellibranchs, R. Eager: New names for some genera of fossil Cardiids, Eberzin (2); Ammonites mammillatus Schlotheim proposed as neotype Douvilleiceras de Grossouvre, F. Hemming; Nomenclature of certain species of Chitones, P. Kaas; Margaritifera problem of nomenclature, D. McMichael (1); Index of Iredale's new scientific names, McMichael & Whitley; Ovatella myosotis (Drap.), K. O. Meyer; Nomenclatural review of genera and subgenera of Cuccullaridae, D. Nicol (1); On the names Pomacea and Ampullarius, T. Pain (1); Biospecific characterization and generic status in Australorbis centimetralis, Paraense & Deslandes; Paludestrina note on nomen-clature and type species, J. Parodiz (5); Kalinga ornata and Polycera (Greilada) elegans, two Nudibranchs attributed to the family Polyceradae, Pruvot-Fol (4); Nautiloid nomenclature Coelonautilus and Cyrtoceras, J. S. Turner; Notes on pelecypod nomenclature, H. E. Vokes (1); On the use of the

generic names Biomphalaria and Planorbis, C. A. Wright (2).

Economics.-Oyster and mussel culture, Anon; The lattice method of oyster culture in the Philippines, G. Ablan: Oyster culture in northern Spain, Andreu & Arté; Shellfish poisoning from Callista brevisiphonata, Asano, Takayamagi & Kitamura; Efficiency and mode of action of molluscocides, Azevedo various Gomes (1): Underwater observations on Pecten maximus (escallop) beds off County Cork, Baird & Gibson; Cultivation of oysters in the Philippines, Blanco, Villaluz & Montalban; On the biochemistry of the purple obtained from Murex and Purpura, Bouchilloux & Roche (1); Pearl fishing off Tuticorin, Gulf of Manaar, India, P. Chacko (1); Natural control of Lunatia heros by Nassarius trivittatus, A. H. Clarke, jun. (1); Population variations in some Californian littoral molluses, W. Coe; Control of pests and diseases of oysters, Cole & Hancock; Oysters and mussels parasitized by Odostomia, Cole & Hancock (1): Recipe for cooking Octopus, P. Colman; Infections of African freshwater snails. C. C. Cridlands: Conditions affecting wood-boring molluses, P. Deschamps; Quantities of scallops off Newfoundland and Nova Scotia, are found insufficient for fishery development, Dickie & Chiasson; Sudan control of snails by CuSO<sub>4</sub>, S. El Din & H. El Nagar; Molluscocides in Brazil, Dobrovolny & Dobbin; Resistance of woods to marine borers, C. Edmondson: Terrestrial molluses as carriers of human illness, E. Frömming; Note On biology of Martesia striata and Teredo navalis, Ganapati & Nagabhushanam; Pecten maximus in Irish waters, F. Gibson; Nutritional value of Gryphaea angulata, Graça, Almeida & Costa; General biology and culture of oysters, mussels, cockles and other economic molluses, H. A. Cole in M. Graham; Bacteriological control relative to Ostrea, A. Grijns; Mussel culture in Holland, B. Havinga; Ostrea edulis parasitized by Mytilicola intestinalis, B. Hepper; Odostomia impressa parasitic on oysters in South Carolina, U.S.A., S. Hopkins; Artificial breeding of oysters in tanks, Imai, Hatanaka et al.: Bacterial control of oysters, Kelly & Arcisz; Enteric organisms in shellfish,

Kelly & Arcisz (1); Quantitative estimation of a molluscicide effect, J. Klock; An automatic molluscoeide dispenser, J. Klock (1); The problem of assessing the potentialities of new oyster grounds, P. Korringa; Oyster culture in S. Africa, P. Korringa (1): Sodium-pentachlorophenate as a molluscicide, R. Kuntz; Oyster rearing in special trays, Le Dantec; Vectors of human bilharziasis in Africa, P. R. Le Roux; Protection of wood against boring molluses, R. Letaconnoux; Review of oyster mortality, its cause, prevention and cure, R. Logie; Utilization of salt water ponds for oyster cultivation. V. Loosanoff; Setting intensity of Crassostrea virginica in Long Island Sound, U.S.A., Loosanoff & Nomejko; Snail control p. 626, Mackie, Hunter & Worth; On the protection of predatory mollusks, A. R. Mead (1); Mussel culture in Germany, Meyer-Waarden; Molluse vectors of bilharzia in Zambézia, T. d. Morais; Development of Martesia striata, R. Nagabhushanam; Molluscoidal activity of some chemical compounds, Nolan & Bond; Studies on the cultured pearl in Japan. Watabe & Tsujii; Marine Okada. molluses injurious to submerged timber, V. Palekar & D. Bal; Control of Teredinidae, L. Popescu; Fishing-industry in Spain and Spanish territories, L. Rey; Economic importance and control of slugs and snails, M. Ritter; Teredo navalis in Maryland, U.S.A. coastal waters, Scheltema & Truitt; Molluscan products, H. Schlesch; Toxicity of antibiotics to snails, Seneca & Bergendahl; Life history and control of Teredo and other wood-boring molluses, F. G. W. Smith; Damage by boring animals to shell of Pinctada maxima, Takemura & Okutani; Occurence of Teredo round N. China coast, Tchang, Tsi & Li; Possible molluscoeide properties of Neorautenenia pseudopachyrhizus, a common coastal plant in Kenya, C. Teesdale; Method for collection of seed-oysters, M. Thiéblemont; Margaritifera vulgaris chemical investigations on pearl formation, Venkataraman & Chari; Oyster culture in Portugal, H. Vilela (1); Crepidula fornicata in Essex rivers, P. Walne (1); Natural resistance of some American woods to marine molluscan borers, F. Wangaard; Effect on oyster spatfall of controlling barnacle settlement with

DDT, Waugh & Ansell; Stream pollution and mollusca tolerant to

such, C. B. Wartz.

Ethnology.—Neolithic shell-mound at Sonno-machi, Chiba-shi, Japan, A. Kan-o; Occurence of cysters in a fishweir excavated at Boston, U.S.A., T. Nelson; Molluscs found in the Indian camp site at Randolph County, Illinois, P. Parmalee.

## STRUCTURE including HISTOLOGY

General Anatomy.—Pandora, figs., J. A. Allen; Structure of Pandora inaequivalvis and P. pinna, figs., J. A. Allen; Anatomy of some freshwater molluses of Portuguese overseas provinces relative to Schistosomiasis. Azevedo & Medeiros: Structure of shell in Thysanolytoceras, Beznosov; Structure of the hermaphrodite canal of Limnaea stagnalis and possibilities of germinal regeneration in pulmonate gastropods, Bounoure & Aubry; Organs associated with optic tract of cephalopods, Boycott & Young (1); Histology of tentacular regeneration in Arion, M. Chétail: Possible functions of rudist pillars and oscules, L. Chubb (1); Note on structure of the horse clam, J. A. Cunningham; Anatomy: The Osphradium and tentacular chemoreceptor organ, figs., J. Demal; Helix pomatia structure, A. Fabricius; Anatomy of Circulus striatus, V. Fretter; Morphology of Melanopsis dufouri, E. Fromming (1); Japanese land snails, T. Habe (2); Anatomy of Japanese Succineidae, T. Habe (10); Anatomy of Haplotrema concava and kendeighi, L. Hubricht (1); Anatomy of Lymnaea truncatula from Japan figs., H. Itagaki; Morphology of Stiliferidae as connected with their mode of life, Ivanov; General description of 2 Japanese squids, E. Iwai; Structure and diversity of form in the Lamellibranchiata, H. Janus; Anatomy of Lympanotonus fuscatus (L.), J. Johansson; Anatomy of Alvania relative to the systematic position of some prosobranchian families, J. Johansson; Revision of Javanese freshwater gastropods, E. v. B. Jutting; Morphology of some bivalve mollusca, Karpevich; Anatomy and biology of the Ellobiidae, Knipper & Meyer; Structural abnormalities in Belemnitella, S. Koltuipin; Anatomy: molluscan

heart, Kriigsman & Divaris: Northern Atlantic species of Cylichna, H. Lemche: Pomacea lineata general anatomy, H. de S. Lopes (1); Anatomy of some Pyrenearia species from Spain, A. O. Lopez; Gastropods, p. 629, Mackie, Hunter & Worth; Alderia uda and Stiliger talis, general anatomy, E. & E. Marcus (2); Anat. of Ellobiidae, J. E. Morton; Anatomy of Otina otis, J. Morton (1); Structure of midgut-gland of molluses, M. Nakazima; Anatomy relative to spermatogenesis of Anadenus altivagus (Thbld.) figs., Nath & Chopra; Genitalia of Drepanotrema anatinum and D. melleum, Paraense & Deslandes (1); Australorbis inflexus n.sp., general anatomy, Paraense & Deslandes (2); Acteon note on anatomy, A. Pruvot-Fol. (3): Morphology of Petricola pholadiformis, R. Purchon; Brechites penis, R. Purchon (1); Anatomy of Martesia striata, R. D. Purchon (2); Anatomy of Gyraulus costulatus (Krauss) figs., G. Ranson (2); Anatomical studies of Naticidae of New Caledonia figs., J. Risbec (1); Morphology of Pomatiopsis cincinnationsis, Schalie & Dundee; Anatomy of the prostrate gland of Biomphalaria, E. Schwarz; Chromdimorphism in Purpura lapillus, H. Staiger; General structure of Cephalopuge trematoides, J. Steinberg: Japanese species of Epitoniidae, general anatomy, I. Taki; Anomia cytaeum anatomy, J. Tanaka; Species of Bulinus (Planorbidae) from Senegambia, figs., C. A. Wright (3).

Radula.—Mastustenthis cardiformis Chun, Octopus vulgaris Lmk., Octopus aegina Gray, Octopus elegans Brock, Hapalochlaena lunulata Q. & G., Hapalochlaena maculosa Hogle, Benthoctopus spp., figs., W. Adam (1); Loliolus investigations Goodrich Loligo etheridgei Berry Uroteuthis bartschi Reider, W. Adam (1); Doryteuthis sibogae Adam, Doryteuthis pickfordi Adam and species of Cephalopods of the Siboga expedition, W. Adam (1); Gibbula pennanti Phil., fig., Fischer-Piette & Gaillard; Cellana depsta, Patinigera fuegiensis, P. kerguilansis, Purpura dumasi, Siphonaria lateralis, Margarella expansa, figs., J. M. Gaillard (1); Pedicularia (Pediculariona) stylasteris Hedley var., T. Habe (5); Usatricolia compta (Gould), Rictaxis puntocoelatus (Carpenter) Nodilittorina pyramidalis Q. & G. figs., T. Habe (9); Novisuccinea horticola (Ruhdt.) Oxyloma hirasei (Pilsbry) O. pfeifferi gamatoi Habe figs., T. Habe (10); Tornatellides boeningi (Schum. and Bttge.), Diseus pauper (Gould), Punctum rota Pils. and Hirase, Parazoogenetes orcula Bens., Pyramidula conica P. & H., Boysidia plicidens (Bens.), Videna gouldiana (Pils.), Vallonia costata tenera (Reiuhdt.) Eastrobilops nipponica (Pils.), T. Habe (11); Oncomelania nosophora radulan formula is

$$\frac{\frac{1+1+1+(1)}{3-3}}{C} - \frac{3+1+3}{L} - \frac{9}{JM} - \frac{6}{OM}$$

H. Itagaki; Helicodiscus singleyanus inermis H. B. Baker figs., J. G. N. Kuiper; Euphaedusa cumingiana recondita fig. of radula, F. Loosjes (2); Function of the radula in some pulmonates, K. Märkel; Ovatella myosotis (Drap.) fig., K. O. Mayer; Pythia scarabaeus fig. Carychium, Leucophytia, Melampus, Marinula, Ellobium, J. E. Morton; Radula of Burnupena figs., V. Orr; Drepanotrema anatinum and D. melleum radula, Paraense & Deslandes (1); Epitonium scalale, Amaea magnifica and Papyriscala latifasciata fig. of radulas, I. Taki; Alycaeus conformis Fulton; A. thieroti de Morgan; A. kapayanensis de Morgan; A. eydouxi Venmans; figs., L. A. W. C. Venmans; Bathypolypus sponsalis P. & H. Fischer fig., K. Wirz.

External anatomy.—Girdle of Chaetopleura spp., E. Leloup; Anatomy of larval oyster, R. H. Millar.

Alimentary system.—Digestive system of the British Ellobiidae, figs., **J. E. Morton** (2); Digestive system of Akera bullata, Morton & Holme: Anatomy of stomach and digestive diverticula of the Lamellibranchia, G. Owen; Anatomy of the stomach and digestive diverticula in the Nuculidae, G. Owen (1); Stomach types of Protobranchia and Septibranchia, R. Pur-chon (3); Microanatomy and histology of the digestive systems of Berghia coerulescens, Idulia coronata, Polycera quadrilineata, Limapontia sp., Thuridilla splendida, Aplysia depilans and Philine aperta, F. Starmühlner.

Muscular system.—Heart contractile and pacemaker mechanism, Krijgsman & Divaris. Vascular system.—Of Oncomelania nosophora (Rolson), H. Itagaki; Struthiolaria vascular system, J. E. Morton (4).

Reproductive system.—Limnaea stagnalis hermaphrodite gland, R. Aubry; Limnaea stagnalis hermophrodite gland, R. Aubry (1); Hermophrodism and the gland of Limnaea stagnalis, R. Aubry (2); Sexual activity in the hermaphrodite gland of Limnaea stagnalis, R. Aubry (4); Genital system of Oncomelania nosophora, H. Itagaki; Actaeon tornatilus anatomy of pallial gonoduct and its phylogenetic significance, J. Johansson; Garnault's duct and the phylogeny of the genital system of Valvata, J. Johansson; Genital organs of Cerithium vulgatum, Triphora perversa and Melanella intermedia, J. Johansson; Genitalia of two Alvania species, J. Johansson; Periodic changes in ovaries and testes of Oncomelaniid snails, Kuo, J.-c. & Pei, H.-k.; Reproductive system of Limax cinereoniger, figs., M. Lamesch; Ianthina note on reproduction, D. Laursen; Anatomy of genital tract of Limax maximus, fig., P. Laviolette (2); Euphaedusa cumingiana recondita genital apparatus, F. Loosjès (2); Male genitalia of Pomacea canaliculata, H. d. S. Lopes: Male genitalia of Planorbidae from Kenya, W. McClelland (1); Berthella agassizii description of reproductive system, E. & E. Marcus; Receptacle seminalis in some Prosobranchia, M. Martoja-Pierson; Reproductive system of the British Ellobiidae, figs., J. E. Morton (2); Genitalian anatomy of Albinaria, figs., K. L. Pfeiffer (1); Genitalia of Biomphalaria smithi, G. Ranson (4); Genital anatomy of Planorbis planorbis, Ranson & Medeiros: Sexual differentiation in gonads of Cavolinia tridentata, Vitagliano Tadini; Structure and shape of the prostrate gland of Biomphalaria alexandrina, E. Schwarz.

Excretory and secretory organs.—Secretions, Moll. pp. 42-45, M. Gabe (1); Nacreous structure and pearls, Grégoire Duchateau & Florkin; Excretory system of Oncomelania nosophora, H. Itagaki; Nerve-secretions Chlamys varia L. and Mytilus edule L., P. Lubet.

Nervous system.—Innervation of Lamellibranch muscle, Bowden & Lowy;

Golgi elements of the neurones of Helix, Gresson, Threadgold & Stinson; of Oncomelania nosophora, H. Itagaki; Nerve structure of Helix pomatia, Jullien & Ripplinger: Structures of visceral nerve of Helix pomatia, Jullien & Ripplinger; Structure of muscles in the heart of Anodonta, Kirpichnikova; Sense organs of Chaetopleura spp., E. Leloup; Octopus vulgaris and Sepiola rondeleti, comparison of brachial innervation, Martoja & May (1); Stru-thiolaria nervous system, J. E. Morton (4): Origin of Golgi dictyosomes from the mistochondric and acrosome formation in Anadenus altivagus, figs., & Chopra: Morphology of Golgi bodies of Anadenus, R. Rishi; Nerves of Octopus arm. Rossi & Graziadei.

Respiratory system.—Anatomy of the gills of Anodonta, R. K. lias.; of Oncomelania nosophora, H. Itagaki.

Shell.—Structure of some lamellibranch shells, G. Athanassopoulos; Dentition and hinge-structure of Trigonioides, T. Kobayashi (3); Shell structure in the middle Cambrian gastropod Scenella and Stenothecoides, F. Rasetti; Shell structure of Pinctada martensii, Watabe & Wada.

Structure of eyes.—Structure and development of cephalopod eyes, G. da F. Sacarrão (1).

#### PHYSIOLOGY

General.-Mechanical properties of Pinna adductor muscle, Abbott Lowy; Bioelectric interactions in Aplysia fasciata, Arvanitaki & Chalazonitis; Biology of Cypraea tigris L., A. C. Boullaire: Regulation mechanism of persistant rhythmicity in Ostrea virginica and Venus mercenaria, F. A. Brown, Ralph & others; Blood pigments, composition, physological importance, E. Eliasseu; Note on biology of Pecten maximus, L. Faure; Osmotic behaviour and bleeding in Crassostrea virginica, Fingerman & Fairbanks: Biology and morphology of Melanopsis dufouri, E. Frömming (1); Metabolic pattern in some bivalves, S. Ishida; Uricase activity in Venerupis philippinarum and purine metabolism and uricase activity in Meretrix meretrix lusoria, Ishida & Tsuzuki; Distribution

of uricase acitivity in Mactra sulcataria, Anadara inflata, Mutilus edulis and Ostrea gigas, Ishida, Tsuzuki & Nitahara; Pila virens physiology of hibernation, V. Meenakshi (2); Caratenoid in Pila virens, V. Meenakshi (3);
Mytilus edulis adaptation to salinity fluctuations, M. Motwani: Water filtration in Martesia striata, R. Nagabhus-hanam (1); Structure and function of midgut-gland of molluses, M. Nakazima; Rate of urine production in Anodonta cygnea, W. T. W. Potts; Osmotic and ionic regulation in Mytilus edulis and Anodonta cygnea, W. T. Potts (1); Cephalopod chromatophores, G. Sacarrao: Genesis of chromatophores in Tremoctopus, G. Sacarrão (2); Acmaea limatula adaptive differences in water-holding capacity of high and low level intertidal forms, E. Segal: Effect of intertidal height on heart rate, gonal size and spawning readiness in Acmaea limatula, E. Segal (1); Tethys grandis osmoregulation, P. B. v. Weel.

Circulatory system.—Regulation of circulation in Octopus, J. de Wilde; Haemoglobin of Anadara inflata, Yagi, Mishima et al.

Alimentation.—Of certain freshwater mollusca of Portuguese overseas provinces p. 26, Azevedo & Medeiros; Lasaea rubra digestion, Ballantine & Morton; Physiology of molluse nutrition, W. v. Buddenbrock pp. 240-283; Digestion and production of H<sub>o</sub>SO<sub>4</sub> by Molluses, G. R. Fish; Martesia striata digestive enzymes, Ganapati & Nagabhushanam (1); Intracellular digestion of proteins in Lamellidens corrianus, D. Kamat; Phosphatases in the digestive diverticula of Lamellidens corrianus, D. Kamat (1); Digestive gland amylase in some gastropods, V. Meenakshi (5); Food movement in gut of larval oyster, R. H. Millar; Digestive rhythm of Lasaea, J. Morton (3); Midgut-gland of molluses, M. Nakazima; Physiology of digestion in the Nuculidae, G. Owen (1).

Nervous Systems.—Neuronic biopotentials in Aplysia, Arvanitaki & Chalazonitis (2); Reactions to shape and duration of memories in Octopus vulgaris, Boycott & Young; Studies on cardiac nerves of Dolabella auricula, A. Ebara; Cation transport in giant axons of Sepia and Loligo, Hodgkin & Keynes; Helix pomatia, Jullien & Ripplinger; Sensitivity of pedal ganglion

of Agriolimax and Arion to osmotic pressures, Kerkut & Taylor; Temperature effect on the isolated pedal ganglion of the slug, Kerkut & Taylor (1); Scent and taste senses in Arion & Limax, R. Kittel; Cytological study of Patella vulgata neurons, Lacy & Horne; Cytoplasmic inclusions of neurones of Patella vulgata, Lacy & Rogers; Neurosecretion in Chlamys varia and Mytilus edulis, P. Lubet (2); Neurohormones of Cyprina and Buccinum, J. Welsh (1).

Reproductive System.—Reproduction of certain freshwater mollusca of Ultramar Portuguese overseas provinces, Azevedo & Medeiros; Hermaphroditism in Patella, J. M. Dodd; Role of gonads in glandular maturation of genital tracts of some gastropods, P. Laviolette; Sexual differentiation in gonads of Cavolinia tridentata, Vitagliano-Tadini.

Respiratory systems and Respiration.

Water circulation in the mantle cavity of Lottia gigantea Gray, figs.,

D. P. Abbott; Alkaline phosphatase activity in gills of Potamida corrugata,

C. Nambudri.

Muscular System.—Resting tension in snail muscle, Abbott & Lowy (1); Innervation of lamellibranch muscle, Bowden & Lowy; Siphon extension mechanism in Mya and Scrobicularia, Chapman & Newell; Muscular mechanisms in Mytilus edulis, Hoyle & Lowy; Pinctada martensi physiology of sensory responses, I. Honjo; Responses of molluscan smooth muscle to chemical stimuli, B. Twarog.

Sense organs.—Studies on the eye of Euhadra amaliae, Hara & Hara; Scent and taste senses in Arion and Limax, R. Kittel.

Chemical contents.—Significance of polysaccharides in gametes of Columbella rustica, B. Battaglia; B. glucuronidase in Helix pomatia, Billet & McGee-Russell; Mytilus edulis oxidation of L-amino acids, Blaschko & Hope; On the 'Purple' of Murex, Bouchilloux & Roche; On the purple pigment of Murex trunculus L., Bouchilloux & Roche (2); Chemical contents, Brunel & Fischer; nature of pigment in Aplysia, A. Christomanos; Polyphenol oxidases in the liver and their probable connecwith melanin formation in Ceratoides

cornvarietis, J. Daugherty; Vitamin A and carotenoids in Mollusca, Fisher, Kon & Thompson; Seasonal variation of chemical contents of Mytilus edulis, F. Fraga: Chemical contents of Lamellibranchs, Glahn, Manchon & Roche; Caratenoids from Flabellina and Archidoris, Goodwin & Fox; Gryphaea angulata chemical composition, Graca, Almeida & Costa; Production of a substance similar to acetylcholine, the heart of Helix pomatia, Jullien, Ripplinger & Cardot; Phosphatases in Lamellidens corrianus, D. Kamat (1); Porphyrin pigments in Akera bullata, Kennedy & Vevers; Cytochemistry of some Pelecypod eggs, H. Kobayashi; Role of the gonad in the glandular maturation of genital tracts, P. Laviolette; Pecten jacobeus chemical content, López-Benito; the Aragonite percentage, H. A. Lowenstam: Pila virens composition of excretory spherioles in the digestive gland, B. Meenakshi; Seasonal variations in glycogen and fat content of Pila virens, V. Meenakshi (1); Pila proteins obtained from free aminoacids in its food, V. Meenakshi (4); Digestive gland amylase in some gastropods, V. Meenakshi (5); Mytilus edulis localization of zinc in spermatozoa, T. Mizuno; Chemical contents of the mantle epithelium of pearl oysters, Okada, Watabe & Tsujii; Helix pomatia blood-sugar level relative to temperature, Pavlović & Bać; Haem compounds in [Mollusca] invertebrates, E. G. Phear; Exchange of phosphorous in shell of Physa acuta, M. Polster; Percentage of copper in the blood of some molluses, P. Rangneker; Purification and properties of the steroid sulphatase of Patella vulgata, A. B. Roy: Pila globosa excretory constituents of the blood, B. Saxena; Pila globosa nutritive constituents of blood, B. Saxena (1); Comparison of redox-pigments of Sepia officinalis with the ommochrome group of insect pigments, I. Schwinck: Nature of pigment cells in Crassostrea virginica, Stein & Mackin; Content and distribution of cobalt in Ostrea gigas, Tsuchiya & Makuta; Ostrea gigas glycolytic intermediates in the gills, Usuki & Okamura; Behaviour of nucleoli and ribonucleo-proteins in growing occytes of Patella coerulea, E. Vannini: Sanguinolaria diphos amino acids of the crystalline style, V. Venugopalan; Carotenoids of some Brazilian species of *Pomacea*, G. Villela; Eggmembrane lysin from acrosomes of *Mytilus*, Wada, Collier & Dan; Dahllite as a constituent of prodissoconch I of *Pinctada martensii*, N. Watabe; *Venus mercenaria*, G. I. Zacks.

Experimental Studies .- Helix pomatia, experiments on the heart, L. Acolat: Experiments on effects of chemicals on the heart of Helix pomatia, L. Acolat (2); Experiments on the heart of Helix pomatia, Acolat & Cornier; Use of radio-isotopes in metabolic studies. with reference to molluscocide action, Azevedo & Gomes; On the foot muscle of Solen siliqua, Bargoni & Atzori; Sepia officinalis visual perception of movement, P. Boulet: Reactions to shape in Octopus vulgaris, Boycott & Young: Temperature sensitivity of unit synapses in Loligo, T. Bullock; Ilyanassa obsoleta development of isolated blastomeres, A. Clement; Experiments on isolated heart of Helix adspersa, M. Corda; Helix adspersa experiments on the heart, M. Corda (1); Experiments on cardiac cycle Helix adspersa before and during stimulation of extrinsic nerves, M. Corda (2); Experiments in Schistosomiasis infection, C. C. Cridland: Spawning and egg production in Crassostrea virginica and Venus mercenaria, Davis & Chanley; Fluorescent experiments on certain Pulmonata, P.-H. Fischer (6); Effect of 3, 4benzopyrene on fresh-water mussels, L. Haranghy; Potassium permeability of giant axons of Sepia officinalis, Hodgkin & Keynes (1); Experiments on pedal ganglion of Agriolimax, Hughes & Kerkut; Experimental breeding of Ostrea edulis in Japanese waters, Imai, Sakai & Okada: Experimental activation of Mactra egg, K. Iwata; Quantitative aspects of filter feeding. C. B. Jorgensen; Helix pomatia survival of isolated heart at a low temperature in different media, Jullien, Cardot Rerat; Experiments on liver of Sepia officinalis, Jullien, Cardot, Ripplinger & Claudey; On the isolated heart of Helix nemoralis, Jullien, Cardot & Vieille-Cessay; Experiments on heart action, Jullien & Ripplinger; Experiments on heart of Helix pomatia, Jullien, Ripplinger & Cardot; Production of a similar substance to acetyl-

choline by the heart of Helix pomatia. Jullien, Ripplinger & Cardot; Helix pomatia experiments on heart movements, Jullien, Ripplinger & Guyon; Experiments on Helix pomatia heart, Jullien, Ripplinger & Joly; Helix pomatia experiments on cardiac inhibition and action of the visceral nerve, Jullien, Ripplinger & Joly; Experiments on Helix pomatia, Jullien, Ripplinger & Mercier: Effect of histamine and antihistamine on behaviour of Cephalopoda, Jullien, Ripplinger & Moreau; Experiments on the contractile and pacemaker mechanisms of the heart, Krijgsman & Divaris; Arion rufus germinal regeneration after castration, P. Laviolette (1): Effects of removal of nervous centres on gamete emission in some lamellibranchs. P. Lubet (2); Experimental infection of Lumnaea humilis with Fasciola hepatica, L. Mazzotti: Relation between zinc and sperm motility in Mytilus edulis, T. Mizuno; Influence of spermatic hormones on spawning and water flow in Crassostrea virginica, Nelson & Allison; Use of radioautography to study pearl formation, Okada, Tsujii, Watabe & Yoshii; Helix adspersa action on the heart of acetylcholine and electrical stimulation at varying hydrostatic pressure, Peruzzi & Corda; Influence on isolated heart of *Helix pomatia* of *Ecbalium elaterium* extract, **Pora &** Rosca; Lengthening and tension reflexes in smooth muscles of some gastropods, N. Postma; Effect of monovalent cations on Limnaea eggs, C. Raven; on Helix pomatia, J. Ripplinger; Helix pomatia experiments on heart, Ripplinger & Joly (1); on Helix pomatia, Ripplinger & Mercier; Artificial activation of eggs of Caecella chinensis, N. Sawada: Behaviour of unstriated muscle, Schmandt & Sleator; Experiments on responses of smooth muscle, B. Twarog; Ostra gigas effects of 1A and fluoride on ciliary activity of gills, I. Usuki; Effect of cyanide and azide on ciliary activity of gills of Ostrea gigas, I. Usuki (1); Ostrea gigas glycolytic intermediates in the gills, Usuki & Oka-mura; Reaction of isolated heart of Helix pomatia to acetylcholine, Velluda, Ticsa & Csutak; Ostrea edulis reared experimentally, P. Walne (3); Tactile discrimination of blind Octopus, M. J. Wells; Effects of acetylcholine on the isolated heart of Venus mercenaria,

J. Welsh; Visual responses by Octopus to crabs, before and after training, J. Z. Young.

## DEVELOPMENT EVOLUTION AND GENETICS

General.—Form and development in relation to mode of life in lamellibranchs. W. Kühnelt; Cytological study of the neurons of Patella vulgata, Lacy & Horne: Alderia uda, early development, E. & E. Marcus (2); Development of Otina otis, J. Morton (1); Development of a pelagic squid, family Cranchiidae, B. Muus; Development of Martesia striata, R. Nagabhushanam: Genetic approach to the systematics of the Brazilian Planorbidae, W. L. Paraense: Development of non-marine fossil mollusc faunas in E. Asia, K. Suzuki; Development of a species of the Lymnaeidae, Trousil-Linhart; Experimental rearing of Ostrea edulis larvae, P. Walne (3); Development of Sepia officinalis, H. Wendt; Phylogeny of ammonites of the Stephanocerataceae and Perisphinctaceae, G. Westermann.

Oogenesis and Oology.—Cytochemical comparison of spermatogenesis and ovogenesis in Columbella rustica, B. Battaglia; Ovocite growth in Aplysia depilans, A. Bolognari; Effect of temperature on development of ovotestis of Cepea nemoralis, J. Bouillon; Nodiolittorina granularis (Gray), fig., T. Habe (4); The floating egg-capsules of Littorina figs., T. Habe (8); Parasitic degeneration of some gastropod larvae, A. Iwanow (1); Egg maturation in Mytilus edulis, K. Iwata; Pallial oviducts of the Cerithiacea, J. Johansson (2); Life history of some molluses from Port Erin, N. Jones: Cytochemistry of some Pelecypoda eggs, H. Kobayashi: Burnupena papyracea early development, V. Orr; Oology Pholas, J. Pasteels; Artificial activation of eggs of Caecella chinensis, N. Sawada; Nucleoli and ribonucleo-proteins in oocytes of Patella coerulea, E. Vannini.

Spermatogenesis.—Helix pomatia spermatocyte structure, J. André; Comparison with ovogenesis of Columbella rustica, B. Battaglia; Spermiogenesis among Mollusca, A. Franzén (12); Eledone cirrhosa, A. Franzén (2); Helix pomatia the spermatozoid, Grassé, Carasso & Favard (1); Structure of mitochondria in spermatozooid of Helix pomatia, Grassé, Carasso & Favard (2); Helix pomatia, Grassé, Carasso & Favard (3); Anadenus altivagus (Tkbld.), figs., Nath & Chopra; Role of the Golg. Bodies in the secretion of the acrosome in spermatogenesis, Nath & Gupta; Eggmembrane lysin from acrosomes of Mytilus, Wada, Collier & Dan.

Sex.—Factors in sexual differentiation of hermaphrodite molluses, G. Bacci.

Embryology and Embryonic stages.—
Ilyanassa obsoleta development of isolated blastomeres, A. Clement; Development of Bithynia tentaculata, O. Hess
(1) and (2); Gonad development in
Patella vulgata, Orton, Southward &
Dodd; Embryology of cephalopod eyes,
G. da F. Sacarrão (1).

Larval stages.—Bacteria in experiments on rearing oyster larvae, P. Walne; Volsella metcalfi and Atrina pectinata japonica early life history, H. Yoshida.

Growth.—Growth of oysters in rivers of Galice, north-west Spain, Andreu & Arté; Growth in Chlamys varia, Audouin & Letaconnoux; Growth of mussels at Boulogne-sur-Mer, H. Brienne; Growthrate in Cardium edule, H. Cole; Growth rates in different populations of Mytilus californianus, P. Dehnel; Various types of growth in Prosobranchiata, P. Drach; Temperature—Shell-growth relations, Epstein & Lowenstan; Environmental influence on Rumina decollata, E. Fromming (2); Seasonal rhythms in development of Bulinus truncatus. Gaud & Dupuy; Growth in molluscan shells, dissection of variation, I. Hamai; Growth of oysters placed in special trays, Le Dantec; Growth of Chlamys varia in northern French waters. Letaconnoux & Audouin; Effect of environment on growth in Venusmercenaria, Pratt & Campbell; Chlamys farreri growth, Tchang-Si, Tsi & Li (1); Age and growth in Chlamys opercularis, E. Ursin; Growth of Crepidula fornicata, P. Walne (1).

Meiosis.—In male Sphaerium corneum, H.-G. Keyl.

Teratology.—Sweden, A sinistral Oxychilus cellarius (Müller), H. Schlesch (1).

Variation—General.—Polymorphism of the shell of Leucochroa candidissima,

J. Altes (1); Variation of Lima (Acesta) goliath from the Japanese Tertiary, S. Aoki; Variation of some mollusca of Portuguese overseas provinces as regards seasons p. 29, Azevedo Medeiros: Variation in successive generations of some genera of Lymnaeidae, C. Boettger: Regularity of direction in growth variation in shells, I. Hamai; Variation in Corbicula due to environment, Y. Hayashi; Local variation in Littorina brevicula, S. Hayasaka; Variation in Katelysia (Nipponomarcia) nakamurai in the Middle Miocene, J. Itoigawa: Environmental variability in molluses, V. Kuznetsov; In Cardium edule, P. Mars; Geographic variation in Cerion from the Bahamas, Mayr & Rosen; Ecologic variation in Burnupena papyracea, V. Orr; Of Murex tribulus in the Great Bitter Lake, Egypt, figs., M. Taylor; Variation in Ceratites from the Muschelkalk, R. Wenger; Species and varieties of Viviparidae represented in the Senckenberg Museum, A. Filch (1).

Variation: ornament and colour.—Leucochroa candidissima polymorphism, J. Altes; On changes of colour in Cephalopods, (colour photos), W. Holmes; Shell banding patterns in two colonies of Cepaea nemoralis from Ontario, W. Judd; Banding and colour variation in a colony of Cepaea nemoralis, M. A. Landman; Polymorphic colouration in Donax, D. Nolte; Variation in size of some White Sea molluses, Savilov; Microgeographic variation as thermal acclimation in Acmaea limatula, E. Segal (1); Polymorphism of Tapes semidecussata, figs., R. F. Shaw; Colour variation in Agriclimax reticulatus, B. Wood.

Variation of form.—In the Lamellibranchiata, H. Janus; Structural abnormalities in Belemnitella, S. Koltuipin; Variation of carinae in Valvata tricarinata, A. La Roque; Shell variation in Pomacea canaliculata, H. d. S. Lopes; Two form variations of Lymnaea auricularia (L.) in one pond, N. F. MacMillan (4); Abnormal Jurassic ammonites, P. Maubeuge; Shell forms, figs., J. H. Stock; Charopa coma variation in shell form, H. E. Whitten.

Sex and Sex Variation.—Genetic stability in populations of Cepaca nemoralis, C. Goodhart; Influence of Mytilicola intestinalis on the gonad development of Mytilus edulis, H.

Mann; Seasonal gonad changes in Ostrea madrasensis from Ennur, Madras, India, K. V. Rao; Sexuality in Mytilus galloprovincialis Lmk., E. Ricci.

Evolution.—Diversity in Partula taeniata, D. Bailey; Evolution of the Ellobidae, J. E. Morton; Evolution of the ammonite families Pronoritidae and Medlicottidae, V. Ruzhentzev (3). Evolution of cephalopod eyes, G. da F. Sacarrão (1); Evolutional lineage of Suchium giganteum, T. Shuto.

#### **ECOLOGY**

## HABITAT, HABITS, FIELD NOTES AND TECHNIQUE

General.—Cephalopods as food of the sperm-whale, Akimushkin (1); Relation between vegetation and mollusc fauna on Belgian sand dunes, A. Anteunis; General biology of molluses of Portuguese overseas provinces, Azevedo & Medeiros; Ecology of Harpax, Boduilevskii: Marine molluses of the North Devon coast, A. Cutcliffe; Ecological and distributional notes on mollusca of the northern Floridan rivers, W. J. Clench (4); Ecology of wood-boring molluses, P. Deschamps; Note on ecology of intertidal molluscs from islands off the Queensland coast, Australia, Endean, Stephenson & Kenny; Bythinella austriaca from mountain springs in Hungary, Farkas & Vág-volgyi; Ecology of Gibbula umbilicalis Da Costa and G. pennanti Phil., Fischer-Piette & Gaillard; Influence of environment on Rumina decollata, E. Frömming (2); Biology of mid-European Pulmonata and Prosobranchia, E. Frömming (4); Ecology of lamellibranchs from the Thyrrhenian sea, Italy. B. Gianfranco: Ecology of Belemnites, Gustomesov, V. A.: Molluscan communities from waters in the Riese region, Germany, L. Hasslein; Ecology of molluses in a saline desert area of Italian Somaliland, Hemming & Verdcourt: Moravian land snails, L. Jansky; Zonitoides, Opeas and Lamellaxis living in greenhouses, E. Karlin; Mollusca of the Biarritz aquarium filter basin, B. Kisch; Ecology of Pisidium in boggy soil, Kozlovskaya; Environmental variability in molluses, V. Kuznetsov; Biological studies on Teredo, C. Lane; Utilization of salt water ponds for oyster culture, V. Loosanoff; Molluscs of the Potamogeton-zone in East

Holstein lakes, Muller-Liebenau; Ellobiidae evolution. J. E. Morton: Ecology of Macoma baltica, Miloslavskava (1); Growth and decline and distribution of marine Pelecypoda, D. Nicol; Land molluscs of the chalk and an elm forest at Örup, Sweden, A. Nilsson (1): Population density effects on Venerupis semidecussata, S. Ohba; Nodilittorina granularis extruding response, Ohsawa & Tsukuda; Ecological observations on Janthina, figs., T. Okutani; Biology of Valvata piscinalis, M. Origny: Molluscs from different habitats in Hiddensee Island, Germany, H. Plate; Ecology of fresh-water molluses of Ammersee, Germany, P. Salzmann; Ecology of Pila globosa, B. Saxena (2); Biology of Capulus ungaricus, M. Sharman; Ecology of Bulinus in Gambia, S. Smithers; Distribution and habitats of freshwater molluses from Coast Province, Kenya, C. Teesdale: Learning and instinct in mollusca, W. Thorpe; Survey of Bulinus in winter, N. Wajdi; On the synecological occurrence of land gastropods, observations on natural habitats, H. W. Waldén; Biology of Loch Lomond. Note on molluses p. 13, A. Weerekoon; Biology of Ianthina janthina, D. P. Wilson; Ecology of molluscan hosts of African schistsomiasis, C. A. Wright (1). Ecology of Bulinus truncatus, H. Zakaria.

Habitat.-Bivalves in unusual habitat, J. F. Allen; Varying habitats of Australorbis glabratus in Puerto Rico, Harry & Cumbrie; Effect of stream gradient on habitat of Australorbis glabratus, Harry & Cumbie (1); Habitats of some Moravian land snails, L. Jansky; A gastropod domicilary in sea urchin spines, H. A. Pilsbry (1): Notes on small land snails from Kaitaia, North Island, New Zealand, L. Price; Habitat of mussels from Lake Texoma, Oklahoma, U.S.A., Riggs & Webb; Fauna of shallow running waters in France, Corsica and North Africa, F. Vaillant; Habitat & factors influential, H. W. Walden; Bivalves living amongst corals from Queensland and N. S. Wales, Australia, L. Woolacott; Habitat and spat mortality of Pecten yessoensis, G. Yamamoto: Habitats Tridacnidae, C. M. Yonge.

Breeding habits.—Self fecundation in Lymnaea, R. Aubry (3); Certain snails of Portuguese overseas provinces, Azevedo

& Mederios: Spawning habits of some Japanese Opisthobranchia, Baba, Hamatani & Hisai: Observations on the spawning habits of Mytilus edulis v. galloprovincialis, H. Bouxin; Cross- and self-fertilization in Lymnaea stagnalis appressa, G. Cain: Spawning and egg production in Crassostrea virginica and Venus mercenaria, Davis & Chanley; Reproduction in the Limapontiidae, T. Gascoigne; Ancylus fluviatilis, R. Geldiay; Spawning of Cerithidea djadjariensis and C. rhizophorarum, T. Habe (3); Nodilittorina granularis (Gray) T. Habe (4); Capsule structure and hatching process of Urosalpinx cinerea, D. Hancock; The breeding season of Mollusca, F. H. Marshall; Hybridization in Cerion from the Bahamas, Mayr & Rosen; Influence of spermatic hormones on spawning in Crassostrea virginica, Allison; Patella vulgata & Nelson breeding in Britain, Orton, Southward & Dodd: Seasonal gonad changes in Ostrea madrasensis, K. V. Rao; Chlamys farreri reproduction, Tchang-Si, Tsi & Li (1); Ostrea edulis reproduction in western France, P. Trochon; Gryphaea angulata reproduction, H. Vilela; Gryphaea angulata conditions necessary for sexual maturity in forms from the river Tagus, Portugal, H. Vilela (2); Spawning in the Tridacnid clams, S. Wade: Experimental breeding of Ostrea edulis, P. Walne (2).

Feeding habits.—Of Conus, Anon (10); Of certain Bulimid and Lymmaeid snails, Azevedo & Medeiros; Lasaea rubra filtering and feeding, Ballantine & Morton; Feeding habits-' Thrushstones,' etc., P. Brien; The snail's foot as a Langmuir Trough, D. Cheesman; Preference of Littorina obtusata for Fucaceae, A. v. Dongen; Algae as food of ovsters. L. H. Flint; Preference of certain gastropods for particular species of fallen leaves, E. Frömming (3); Food absorption of Pomatias elegans, E. Fromming (5); Cerithiolum reticulatum nutrition, N. Gajevskaja; Feeding in the Limapontiidae, T. Gascoigne; Conus striatus and C. catus feeding on live fishes, A. J. Kohn; Stomach contents of Gryphaea angulata, S. Le Roux; Cassididae species Predatory on Edinoderms D. R. Moore; Agriolimax reticulatus feeding on carrion, G .W. Pitchford; Method of feeding in Petricola pholadiformis, R. Purchon; Grazing rate of Ostrea edulis larvae reared experimentally, P, Walne (3); Unusual method of capturing prey by a cuttlefish, J. Wickstead; Feeding in Tridacnidae, C. M. Yonge.

Habitat general and special.—Mimicry in Anomia ephippium, L. Acolat (1); Memory in Octopus, Boycott & Young (2); Ability of some gastropods to erode littoral rocks, P. Fischer; Seasonal habits and activity of pulmonates, P. H. Fischer (4); Ink ejection in the escape reaction of a squid, D. Hall; Swimming of Akera bullata, Morton & Holme; Nassarius festivus nocturnal activity, S. Ohba; Non-homing Limpets, R. Ohguski; Behaviour of Pila globosa, B. Saxena (2).

Boring habits.—Ability of some gastropods to erode littoral rocks, P. Fischer; Control of Teredo navalis, L. Popescu; Boring adaptation in Botula and Lithophaga, C. M. Yonge (1).

Movement.—Of Sepia officinalis L., P. Boulet (1).

Enemies.—Control of oyster enemies, Cole & Hancock; Penetration of bivalves by sea stars, M. Lavoie; The green crab as a molluscan shellfish enemy in Eastern U.S.A. and Canada, Medcoî & Dickie; Ocinebra erinacea (L.) on the increase on the East Coast of England, Mistakidis & Hancock.

Parasitism and Commensalism .-Chemical control of molluscan Schistosoma hosts, Adlung & Kauth; Parasitic castration of Pandora albida by Cercaria melanocystea, Arvy & Gaillard; Susceptibility of different species Planorbis to Schistosoma mansoni infection, Azevedo & Faro; Bilharziasis survey in S.W. Asia, Azim & Gismann; Turveria gen. nov. ectoparasitic on echinoid, S. Berry; Melanella comatulicola parasitic on Anteda mediterranea, J. Changeux: Control of Australorbis glabratus by a snail and a leech, Chernin, Michelson & Augustine; Odostomia parasitic on mussels and ovsters. Cole & Hancock (1); Commensalism Capulus hemiastericolas sp. nov. and Hemeaster battensis, A. Devries; Association Velutina velutina and the between ascidian Styela, M. Diehl; Land gastropods as carriers of human illness, E. Frömming; Effects of parasites on Philine aperta and Dentalium vulgare, J. M. Gaillard (3): Habitats of Austra-

lorbis glabratus, vector of schistosomiasis, in Puerto Rico, Harry & Cumbrie: Possible commensalism between Spirorbis and some lamellibranchs of the Carboniferous in the Low countries, S. v. d. Heide; Ostrea edulis parasitized by Mytilicola intestinalis, B. Hepper; Odostomia impressa parasitic on oysters. S. Hopkins: Occurrence of Oncomelania quadrasi, host of Bilharzia, in the Philippines, B. Hubendick (1); Means of penetration into host by parastic gastropods, A. Iwanow: Parasitic degeneration of some gastropod larvae, A. Iwanow (1); Bilharziasis in Angola, Janz & Carvalho; Control of bilharziasis by molluscocides, J. Jove: Glochidium sp. parasitic on fish in Czechoslovakian streams, V. Kăsták; Melanoides tuberculatus, effect of parasitism on the digestive gland, Lal & Premvati: On bilharziasis, L. Lamy; Africa. Vectors of human bilharziasis, P. R. Le Roux; Ecology of snail vectors of schistosomiasis in Kenya, W. McClelland (1); Observations on the snail vectors of Schistosomiasis in the Gold Coast, F. McCullough (1); Transmission of schistosomiasis by Bulinus spp. in the Gold Coast, F. S. McCullough (2); Mytilus edulis attacked by Mytilicola intestinalis, Meyer-Waarden & Maun; Parasitology on Echinoderms by a Gastropod, H. A. Pilsbry (1); Protozoan parasites of Bithynia tentaculata and Dreissensia polymorpha from Poland, Z. Raabe; Schistosomiasis in the Congo, J. Schwetz; Classification problems of snail vectors of schistosomiasis, J. Schwetz (1): Methods of controlling gastropod vectors of schistosomiasis, J. Schwetz (2); Studies on Schistosoma infection, Schwetz, Fort & Baumann; On Schistosomiasis problems, Schwetz, Fort & Baumann (1); Bulinus vectors of schistosomiasis in Gambia, Smithers; Relation of numbers pigment cells in Crassostrea virginica to the fungus disease Dermocystidium marinum, Stein & Mackin; Cephalopyge trematoides in New South Wales, Australia, J. Steinberg; Parasitic mollusca of fish in Ribinsk water-reservoir, Stolyarov: Damage to Pinctada maxima shells by boring animals, Takemura & Okutani; Habitats of snail vectors of schistosomiasis in Fukien, China, Tang, Chow, Wang, Sieh & Chow; Quantative survey of Bulinus in winter, N. Wajdi; Bulinus forskalii as vector of Sehistosoma haematobium, C. A. Wright; Ecology of molluscan hosts of schistosomiasis, C. A. Wright (1); Ecology of Bulinus truncatus, H. Zakaria; Physopsis globosa intermediate host of Schistosoma found in Italian Somaliland, E. Zavattari.

Effects of Environment.—On the growth and breeding of Ancylus fluviatilis from the Isle of Skye, W. R. & M. R. Hunter (1); Indoplanorbis exustus environmental conditions, Kathuria, Rao & Hiregaudar; Effects of environment on Tapes semidecussata, figs., R. F. Shaw.

Effects of Temperature.—Effect of temperature on development of ovotestis of Cepea nemoralis, J. Bouillon; Temperature sensitivity of unit synopses in Loligo opalescens, T. Bullock; Temperature effect on the isolated pedal ganglion of the slug, Kerkut & Taylor (1); Helix pomatia blood-sugar relative to temperature, Pavlović & Bać; Thermal effects on an intertidal gastropod, Acmaea limatula, E. Segal (1); Temperature effects on Crepidula fornicata, P. Walne (1).

Luminescence and Light and Heat.— Pholas dactylus luminescence, J. & J. Gaillard; Effects of ultra-violet radiation on Oncomelania, Wong & Wagner.

Effect of toxins.—Chemical control of molluscan Schistosoma hosts, Adlung & Kauth; Effect of narcotics on Aplysia, Arvanitaki & Chalazonitis (1); Effect of molluscocides on Physa acuta and Helisoma duryi normale, Azevedo and Gomes (1); Effect of 3, 4-benzopyrene on fresh-water mussels, L. Haranghy; Control of bilharziasis by molluscocides, J. Jove; Atropine-calcium antagonism in Helix pomatia, Jullien & Ripplinger; Physa rivularis action of sodic acid on the developing egg, V. Mancuso; Poisoning due to cone shell 'bite', L. Petrauskas; Effect of monovalent cations on Limnaea eggs, C. Raven; Effect of Potassium and Calcium ions on the heart of Helix pomatia, Ripplinger & Joly: Effect of salt, vinegar and heat on the coliforms in oysters, E. Salafranca; Ostrea gigus effect of 1A and fluoride on ciliary activity of gills, I. Usuki; Effect of cyanide and azide on cilia of Ostrea gigas gills, I. Usuki (1); Effect of malonate and 2, 4-dinitrophenol on ciliary action of Ostrea gigas gills, I. Usuki (2); Effect

of pentachlorophenol on metabolism of Australorbis glabratus, Weinbach & Nolan.

Association and effects.—Modiolaria and Ascidians, A. Bourdillon; Green algae epiphytic on the shells of freshwater molluses, M. Campion; Association of Velutina velutina and an ascidian, M. Diehl; An association between Mythus, Purpura and Balanus, M. J. R. de. Figueiredo; Bryozoa on living Gasterpod shells, Y. Gautier; The pelecypod Vulsella living in a sponge, K. v. Haffner; Lymnaea stagnalis and associated species in Llywenan 5 m. N. of Llyn Maelog, Anglesey, N. F. McMillan (5); Capulus ungaricus on shells of living lamellibranchs, M. Sharman.

Defences and offences.—On the stinging operation of Conus, G. F. Kline.

Population.—Colony formation in Acmaea dorsursa, N. Abe; Population dynamics of Mya arenaria, J. C. Ayers; Population variations in some littoral molluscs, W. R. Coe; Multiplication of some mollusca in Baikal sea, Russia, Gavrilov; Mollusc population of the Javornik Mts., Czechoslovakia, V. Hudec; Distribution and growth and decline of population of marine Pelecypoda, D. Nicol; Population balance between limpets and seaweeds on wavebeaten shores, A. Southward.

Technique.—Use of radioisotopes in metabolic studies, with reference to molluscocide action, Azevedo & Gomes; Method for weighing mollusc larva, St. Beldescu: On method and uses of glucosamine for experiment on the isolated heart of Helix pomatia, Binet, Jullien, Ripplinger & Cardot; High tide collecting, E. P. Chase; Use of snail stomach cytase in plant cytology, T. C. Chambers; Use of exographs to reveal shell structure, R. Friede; Use of sodium taurocholate as a lysin, J. George: Microscopical investigation of some Belemnites, G. Göke; On the preparation of Nudibranchiata, G. D. Hanna; A method of sampling, Johnson, Southwood & Entwistle; Method of molluscocide application in Venezuela, J. Jove; An automatic molluscocide dispenser, J. Klock (1); Weathering and limestone treatment, K. Komárek; Spectographic examination of fossils, McKerrow, Taylor, Blackburn & Ahrens; Use of deep-sea photography to learn more about squids, N. Mackintosh; Use of radioautography to study pearl formation, Okada, Tsujii, Watake & Yoshii; Estimation of aquatic snail populations, Olivier & Schneiderman; In Oology, J. Pasteels; Technique in showing certain phases of living structures in Helix, A. L. Roque; Separation of small bivalves from sediments by difference in specific gravity when immersed in ZnCl<sub>2</sub> solution, G. Sellmer; Polymorphism of the shell of Leucochroa candidissima, J. Altes (1); Use of exographs to reveal shell structure, R. Friede;

Conchometry. Conchometry and relative growth in molluscan shells, direction and formulae, J. Howai.

Duration of Life.—Freshwater mollusca, Portuguese overseas provinces, Azevedo & Medeiros; Lowestoft mortality in Mactra and Sepia, H. E. Jenner.

Adaptation of Pandora inaequivalvis and P. pinna, J. A. Allen; Adaptation of Freshwater gastropods, W. D. R. Hunter; Venus mercenaria acclimatized in Brittany, L. Marteil; Adaptation to rock boring in Botula and Lithophaga, C. M. Yonge (1).

Regeneration. — Regeneration of drilling mechanism in *Thais haemastoma*, **Demoran & Gunter**; Regeneration in *Arion rufus* L., P. Laviolette (3);

#### ZOOGEOGRAPHY

Dispersal and Distribution.—Zoogeography of Yoldia, W. K. Ockelmann; Distribution of land snails in plant association, J. B. Burch; On the distribution of freshwater shells, H. Kajiyama; Molluscas from the Pleistocene of North Greenland (Pearyland), p. 22, D. Laursen (1); Distribution of Pelecypods in Murman in relation to temperature changes, Miloslavskaya.

General Lit. and Catalogues.—Locomotion of pelagic molluses, Anon (1); Sonic scattering layer caused by swarming of Firoloida desmaresti, M. Blackburn; Catalogue of Polyplacophora from southern S. American waters, Z. Castellanos; Tortonian fossils from the Modry Kamin district, Slovakia, Čechovič & Genaš; Occurrence of clines in molluscan populations, W. J. Clench (1); Tertiary Pectioracea from East Africa, Persia and the Mediterranean region,

Eames & Cox; List of molluses from the Red Sea, A. Franc (1): List of molluses from Bass Strait, south east of King Island, Victoria, Australia, C. Gabriel; Lists of pearl forming mollusca. Grégoire, Duchateau & Florkin; Golgi elements of Helix neurones, Gresson, Threadgold & Stinson; Species included in the genus Homolsomites, R. W. Imlay (1); Ranges of Homolsomites Craspeditidae and species ascribed to the genus. R. W. Imlay (1); On larval molluses, molluse shells and pearls, H. L. Jakl; Neolithic shell mound at Sonno-machi, Chiba-shi, Japan, A. Kan-o; Iredale's writings and new scientific names, McMichael & Whitley; Breeding season in molluses, F. H. Marshall; Checklist of Miocene Pelecypoda from Turkmenstan & Kazahkstan, Merklin & Nevesskaya; Genera and subgenera of Cuccullacidae and types, D. Nicol (1); Nomenclatural review of Cuccullacidae, D. Nicol (1): Mollusca from the "Zone N. of Cannalots," J. Paris; Species observed at Fontainbleau during 1955, Y. Quideau (1); Morphology and biology of fertilization, Lord Rothschild; Chromatophores and colour change in cephalopods, G. Sacarrão: Lists of Molluscan species of the French Mediterranean coastal regions, C. F. Sacchi; C. B. Adams species described from the Eastern Pacific, figs., R. D. Turner (1); Atlantic species described by C. B. Adams, figs., R. D. Turner (2); List of some mollusca tolerant to certain degree of freshwater pollution, G. B. Wurtz.

British Isles.—Species from the Kallaways Beds at Kidlington, Oxfordshire, J. H. Callomon (1); Wiltshire fault Ammonites, R. Casey; Type and figured specimens from the Tortworth Inlier, Gloucestershire, M. L. Curtis: Interglacial mollusca of Bobbitshole, Ipswich, A. G. Davis (1); Lancashire, Mactra corallina cinerea in vast numbers after a storm, L. W. Grensted; List of molluses from the Orkney Islands, Scotland, R. Rendall: Non-marine mollusca of the interglacial of Hoxne, Suffolk, B. W. Sparks: Hertfordshire Mollusca, a catalogue with references and notes, L. W. Stratton; Malham mollusca, lists and maps, L. W. Stratton (2); Mollusca of Malham, Yorkshire, L. W. Stratton (4); Mollusca of Strangford Lough and adjacent coasts, Ireland, pp. 87-108, G. Williams.

Europe.-Mollusca in Donetz basin, Namur, Russia, Aïzenberg & Brazhnikova; Revision of the Liogryphaea of the Luxemberg Museum, Charles & Maubeuge; Mollusca of the River Siagne, France, A. Comfort; Achatininae in the Muséum de Paris, M. Denis; Tertiary bivalves from Zeeland, Holland, H. v. Halm; Mollusca of Pags du Herre, J. Leclercq; Molluscs from Ostend, Belgium, pp. 62-63, 85-88, Lefevere, Leloup & Meel; Species of the Interglacial of Elge, G. Lüttig; Gyraulus laevis, figs., G. Lüttig; Notes on certain species from various localities in the Mediterranean, P. Mars (1); Fauna of Paunou, Vienna, A. Papp; Pleistocene fossils of the Netherlands, pp. 25-31, Vlerk & Florschütz; Land gastropods of the vicinity of Stockholm, H. W. Walden.

Asia.—Pliocene mollusca of the Cnide Peninsular, Turkey, G. Chakut; Anatomical studies in land snails, T. Habe (2); Land shells of Yaku Island, Japan, T. Kiroda; Pleurotomaria of Japan, T. Kuroda (2).

Africa.—List of molluses not previously recorded from the Gold Coast, Buchanan & Anderson; East African Tertiary Pectinaeea, Eames & Cox.

North America.—Some land and freshwater mollusca of the arctic slope of Alaska, figs., G. D. Hanna (1); Virginia, land snails of Shenandoah Park, L. Hubricht; Barrimian Ammonites genera occurring in Trinidad, figs., R. W. Imlay; American Pennsylvanian Cephalopods, A. G. Unklesbay; Ordovician Nautiloids from Central Texas, figs., A. G. Unklesbay (1); Checklist of cephalopods from Florida, G. Voss.

South America. Species of Jurassic mollusca from the Nasca area, Peru, L. R. Cox (4).

Japan.—Illustrated catalogue of the Pholadidae, Taki & Habe.

New Zealand.—Checklist of New Zea. land archibenthal Mollusca, R. K. Dell-

Oceanic Islands.—Mollusca from Kerguelen Island, J. M. Gaillard (1).

Museums, Collections, and Expeditions.—The Beatrice and Kimball Valentine collection now in the Acad. Nat. Sci., Philad., R. R. Abbot; Luxembourg Museum, revision of the Liogryphaea species therein, Charles &

Maubeuge; Muséum de Paris Achatininae, M. Denis; William Hill collection, [A. E. Ellis] (2); Eustoma in the Ecole de mines, Paris, P. H. Fischer (5); Mollusca of the "Calypso" voyage in the Red Sea, 1951-52, A. Franc (1); Kerguelen—gastropods reviewed from MM. Angot, A de la Rüe, Brown and Paulian, J. M. Gaillard (1); Mollusca Terre Adèlie (1950) "Com. Charcot" Exped., J. M. Gaillard (2); Fresh water molluses from Prof. Gisten's Australian Expedition 1951-52, B. Hubendick; Molluses of J. Klapperich's Afghanistan Expedition 1952-53, S. Jaeckel (1); Naiades types in the Museum of Comparative Zoology, Harvard, R. Johnson; On a collection of Pomacea from Colombia, T. Pain; Muséum Demidoff (Museum of the University of Moscow), K. v. W. Palmer; Whitfield collection at the University of California, Peck and MacFarland: Strophocheilidae of the Museum Hist. Nat., Paris, E. Salmon; Some molluses of the Austrian Tyrrhenia Expedition, 1952, F. Starmühlner (1); Viviparidae types in the Senckenberg Museum, figs., A. Zilch (1); Senckenberg Museum, types of Cyclophoridae (4), figs., A. Zilch (2).

[1956]

#### DISTRIBUTION

(A) Geographical

(a) FOSSIL AND LAND AND FRESHWATER

PALEARCTIC REGION

Palaearctic.—Vitrina, L. Forcart.

General.—Benthic distribution of molluses occuring below 100 fathoms in New Zealand waters, R. K. Dell, p. 179; Distribution of Succinea elegans, Matekin, P. V. (1); Distribution of Helix asparsa, W. J. Rees.

Greenland.—New Cretaceous Belemnites from West Greenland, T. Birkelund; Permian Pelecypoda of East Greenland, pp. 16-17, N. D. Newell; New Jurassic Velata from Greenland, A. Rosenkrantz.

Scandinavia.—New Loricata from the Silurian and Cretaceous of Sweden, J. Bergenhayn; Norway lobsters and oysters, A. Dannevig; Pleistocene fossils Vestspitsbergen Feyling-Hanssen; Marine mollusca of the late Pleistocene from Westspitsbergen, pp. 20-24, 29, 176-177, Feyling & Hanssen (1); Illustrated handbook of Swedish Prosobranchia and Pulmonata, B. Hubendick (2); Land molluses of Örup, Sweden, A. Nilsson (1); Distribution of marine mollusea in Northern Norway, new records, T. Soot-Ryen; Land mollusea from the vicinity of Stockholm, H. W. Walden.

British Isles.—Bookham Common, the freshwater mollusca of the bombcraters and gun pits, C. F. J. Bensley: New Loricata from the Ordovician of Scotland, J. Bergenhayn; Vertical ranges of certain snails in the North of England, Blezard; London clay fauna additions, Brown & Castell; Corallian fossils from Beckley, Oxfordshire, J. A. Callomon; Mollusca from the Isle of Cumbrae, Firth of Clyde, Clark & Milne; New Scottish Carboniferous Goniatites, E. Currie; Type and figured specimens from the Tortworth Inlier, Gloucestershire, M. L. Curtis; Swanscombe, Kent, the occurrence of Pisidium obtusale lapphonicum Clessin, S. P. Dance: A new Sussex locality for Pisidium pseudosphaerium Favre, S. P. Dance (1): Suffolk interglacial species of Bobbitshole, Ipswich, A. G. Davis (1); Hygromia subvirescens between Beer Head and Branscombe, S. Devon, A. G. Davis (2); Hygromia subvirescens at Harlyn Bay near Padstow, Cornwall, A. G. Davis (3); Dorset, Anodonta minima in the Corfe River, A. G. Davis (4); Dorset, occurrence of Anodonta minima Millet, A. G. Davis (5); Ammonites from Blue Lias, Somerset, D. Donovan (1); Mollusca from the Bure Valley, Norfolk, England, E. A. Ellis; Leicestershire, some species of the Marlstone rock-beds, A. Hallam; List of Scotland distribution records of nonmarine molluses, W. R. Hunter (2); Mollusca of the Scottish Mountains, W. & M. Russell Hunter; Scotland, new Lamellibranchs from the Pautland Hills. A. Lamont; Charmouth Lias and Ammonites, W. D. Lang; Ena montana Drap. in Hertfordshire, L. Lloyd-Evans; Acroloxus lacustris (L.) in Merionethshire, L. Lloyd-Evans (1); Mollusca of Groomsport Co. Down, MacDonald & MacWilliams; Local non-marine mollusca of Lancashire and Cheshire, N. F. McMilhan: Notes on marine lamellibranes washed ashore in Liverpool Bay and on terrestrial molluscs from Liverpool and Cheshire, N. F. McMillan

(6); Mollusca of Cheshire and North Wales, McMillan & Millott; Molluses & Perkins; Potamopyyus jenkinsi in Lake Windermere, H. P. Mason; Fauna of a sand-bank in the Dee estuary, Scotland, E. Perkins; Mollusca from the river Lugg at Shelwick, Hereford, E. J. Perkins (1): Lower Carboniferous North of England, D. H. Rayner; The distribution of certain inter tidal species around the Irish coast, Southward & Crisp: Interglacial non marine mollusca of the Quaternary deposits at Hoxne, Suffolk, D. W. Sparks; The Molluscs of Hertfordshire, L. W. Stratton; Note on some molluses in Hertfordshire, L. W. Stratton (1): Mollusca of the Malham area, L. W. Stratton (2); Yorkshire, Mollusca of the Malham area, L. W. Stratton (4); Note on molluses found in Loch Lomond, Scotland, p. 13, A. Weerekoon: Lists of Mollusca from Strangford Lough and neighbouring coasts, Ireland, pp. 87-108, G. Williams; Utaturiceras new Cretaceous ammonite from Devon, England, C. W. Wright.

France. — Cretaceous Radiolitidae from the North Pyrenees, G. Astre (1); List of Tertiary molluses from St. Gervais, France, P. Balavoine; Creniceras crenatum from the French Jurassic, A. Caire; France, new Liogryphara species from the Jurassic, Charles & Maubeuge (1): Mollusca of the R. Siagne, France, A. Comfort; A new Neocomian Belemnite from S. France, M. Delattre; Provençale Mollusca, J. P. Vanden Eeckhoudt; Mollusca of Banyuls, P.-H. Fischer (2); Saint Malo Bay fauna, A. Franc; Ammonites from the Jurassic of Deux-Sèvres, France, J. Gabilly; Fauna of the 'Trummer Oolite' of Upper Malm in Württemberg, Lower Dogger, Ortunian, O. F. Geyer (1); Pliocene molluses from Mirabel, Ardèche, France, Grangeon & Jodot: Distribution of Monodonta and Gibbula in an area of the Gulf of Marseille, P. Huvé; France, faunule of Eocene of Sainte-Madeleine, P. Jodot; Mollusca of the filter basin of Biarritz aquarium, B. Kisch; List of Mollusca of lakes near Liège, France, pp. 99-100, Leloup, Meel & Jacquemart; Quaternary Mollusca from Bas-Languedoc, France, Mars & Ottmann; New Jurassic ammonites from France, P. Maubeuge (1); Mollusca of the Delta of the Rhone, M. Paulus: Mollusca of Fontainebleau

and Lower Valley of Loing, Y. Quideau; An ecological study of the French Mediterranian and the land mollusca, C. F. Sacchi; Lower Cretaceous ammonites from S.E. France, S. Sarkar; Verieties and district mollusca, L. Sarlet; Terrestrial molluscs of Biarritz, R. Soyer; Upper cretaceous, gastropods, G. Sermier; Barthonian fossils of Côte d'Or, Tintant & Joly; List of Jurassic ammonites from Dijon, France, Tintant & Larcher; Burdigalian fossils of Cestas, Giroude, Vigneaux & Magne.

Holland.—A new Upper Pliocene Pyrene, C. O. v. R. Altena (1); Fossil shells of the coast, Altena, Bloklander & Pouderoyen; A new fossil Natica from the Post-Pleistocene, Pouderoyen; Netherland land molluscs. A key with figs., J. H. Stock; Netherlands Pleistocene, Vlerk & Florschütz; Miocene molluscs of South Jutland, L. B. Rasmussen.

Belgium.—Relation between vegetation and molluse fauna on Belgian sand dunes, A. Anteunis; Lists of Miocene species of Turridae from Belgium, pp. 61-70, M. Glibert; Molluses of Pags du Herne, J. Leclercq; Molluses from the brackish waters at the mouth of Escant river, Belgium, Leloup & Konietzko, p. 82-86.

Switzerland.—Gastropods from Swiss caves, Aellen & Strinati.

Spain.—New Cretaceous Arcomytilus from Spain, J. Bataller; Spanish Cretaceous Mollusca, J. Bataller (1); Jurassic Mollusca from Cardo, Spain, J. Bataller (2); New Jurassic molluscs from Spain, Castell, Concha & Revilla; Fishing-industry in Spain and Spanish territories, L. Rey; Terrestrial and littoral gastropods of Santander, Spain, J. A. Sanchez.

Portugal.—Miocene molluses from Lisbon, J. Berkeley-Cotter; Nassarius and Ancilla new species from the Portugese Pliocene, L. Cox; Gryphaea angulata from the Tagus river, Portugal, state of maturity, H. Vilela (2).

Italy.—Fossil Limax found in a cave near Como, Italy, Cadeo & Senna; Observations on the benthonic and planktonic populations in Lake Maggiore, Italy, Corbella, Croce & Ravera, Lamellaptychus, Punctaptychus and Laevilamellaptychus from the Italian Jurassic, G. Cuzzi; Gambetta miocene and

species, David & Judot; Mollusca from the Dolomite Mountains, Italy, G. Marcuzzi; Eocene Gastropoda from Udine, Italy, B. Martinis; Limax duplex from Tamborin grotto, Como, Italy, C. Maviglia; New Quarternary Parastrophia from Italy, M. Moncharmont Zei; Molluscs of the Cape of Sorrento, Italy, F. Starmühlner (1); Land and freshwater mollusca of the Dolomites, Italy, L. A. W. C. Venmans (2).

Luxembourg.—The Limacidae and Arionidae, figs., M. Lamesch; New Jurassic ammonites from Luxembourg, P. Maubeuge (2); New Jurassic ammonites from Luxembourg, P. Maubeuge (3).

Germany.—Mollusc fauna of the Görlitz area, Germany, H. D. Boeters; Mollusca from flowing waters in North Württemberg, H. Buck; New Triassic gastropods from Germany, H. Eckert; Württemberg notes on and occurrence of Diceras speciosum (Münsten), O. F. Geyer; S. Bayern Belemnitella zone, H. Hagn; Mollusca and molluscan communities from waters in Riese region, Germany, L. Hasslein; Two prehistoric specimens of Margaritana in Saxony, R. Hertel; Württemberg, and occurrence of Oxytoma scanica in the Planorbis-zone, H. Hölden; Mollusc fauna of Feldberg, Germany, S. Jaeckel: Freshwater Mollusca from the late and post Glacial deposits of Schleswig-Holstein, S. G. A. Jaeckel; Aquatic molluscs of Nuthe-Niederung, Elbe and Warthe, Germany, S. Jaeckel; Lartetia rhenana in Germany, F. Kiefer; Mussel culture in the German Waddensea, Meyer-Waarden; New Devonian ammonites from Germany, K. Müller; Molluscs of the East Holstein Lakes, Germany, I. Muller-Liebenau; Carboniferous fossils of Frankenberg, E. Paproth; Molluses of Hiddensee Island, northern Germany, H. Plate; List of molluses from differing environments, near Görlitz, Germany, L. Ritter; Fresh water molluscs of Ammersee, Germany, P. Solzmann; Note. on molluses of Burgenland, Germany, F. Sauerzopf; Mollusca from waters in the area of Lübbeck, Germany, E. Schermer; Mollusca of inland waters in West Germany, E. Schermer (1); Theba carthusiana in Mecklenburg, Germany, H. A. Schmidt (1); Ceratites species from the Miocene of the Saar, Germany, E. Schneider; New

Jurassic ammonites from Germany, A. Zeiss; Type specimens in the Senckenberg Museum of Cycloporidae, A. Zilch.

Europe.—Comparison of British and Mediterranean Pliocene and Pleistocene molluscs. D. Baden-Powell: Distribution of Viviparidae in Europe, C. R. Boettger (3); Cyrenaica Clausiliidae, R. A. Brandt: Biology of mid-European Pulmonata and Prosobranchia, E. Frömming (4): Molluscs from the basin of the lower Danube, Grossu & Paladian; Helicodiscus singleyanus inermis H. B. Baker new to fauna, J. G. J. Kniper: Ammonites from the Clymenia limestone of Lower Silesia, S. Lewowicki; Elze interglacial fauna, G. Lüttig; Mollusca of Western Bohemia, V. Ložek (4): Mollusca of central Bohemia, V. Ložek (5); Holocene species of the Rhone, right and left estuaries, G. Mazenot; Fossil land snails from South East Europe, A. Papp (1); Senonian fossils from Leštar, E. Servia, O. Pejovic: Cretaceous molluses of N.W. Europe, L. v. d. Poel; Inoceramus from the Tertiary of Trans-Carpathia, Vyalov & Pasternak; Jurassic species of Sphaeroceras and Chondroceras in Europe G. Westermann (1); Chlamys latissima from the Neogene of Moravia, V. Zázvorka.

Austria.—Mollusca from Austria, F. Mahler; Mollusca from lakes near Lunz, Austria, Mahler & Sperling; Pannon, Vienna Becken, A. Papp; Bohemian Holocene molluscs, some new forms, J. Petrbok (1); Mnataia new Pelecypod genus from Bohemia, Prantl & Ružička; Tertiary Lamellibranchia from Vienna, Austria, R. Sieber (1).

Hungary.—New Miocene molluses from Hungary, I. Csepreghyné-Meznerics; Miocene molluses from Hungary, I. Csepreghyné-Meznerics (1); Mollusca from springs of Mt. Dunazug, Hungary, Farkas & Vágvölgyi; Species of the Upper Oligocene of Hungary, L. Strausz (1); Some Hungarian molluses, J. Vágvölgyi.

Poland.—Planorbidae from the Warsaw area, S. Feliksiak; List of species from the Miocene of Andrychow, Poland, W. Krach & W. Nowak; Mollusca of sand, clay and brown coal deposits of Rava Russka, Poland, Kudrin (1); Molluscs from the Silurian of Lwów, Poland, Sandler & Glushko.

Roumania.—Cretaceous Matheronia sp. from Roumania, Atanasin Semska; First record of Potamopyrgus jenkinsi from Roumania, A. V. Grossu: Gastropods of Minis-Banat, Roumania, A. Grossu (1); New gastropod from Roumania, A. V. Grossu (2); Roumanian Opisthobranchia & Prosobranchia, A. V. Grossu (3): Unio wetzleriformis new Miocene species from Rumania, N. Macarovici: Valenciennius annulatus in the Pliocene of Roumania, P. Nita; Unio & Melania spp. from the Roumanian Pliocene, M. Paucă: Anthracosia from the Roumanian Permian, G. Răileanu: Aptychus fauna from the Roumanian Jurassic, Răileanu, Bădălută & Pelin; New species of Anodonta from the Roumanian Pliocene, Rosculescu-Soverth.

Czechoslovakia.—Molluscs of the Javornik Mts., Czechoslovakia, V. Hudec; Occurence of Cerithium and Melanopsis martiniana near Olomone, Czechoslovakia, V. Kalabis; Molluscs from Moravice, Czechoslovakia, O. Kapler; Mollusca of Central Lake region, Czechoslovakia, J. Lellák; Note on Czechoslovakian molluscs, V. Ložek; Mollusca of the Erze mountains, Czechoslovakia, V. Ložek (1); Czechoslovakian Mollusca, V. Ložek (2); Catinella, Pupilla and Pisidium from Czechoslovakia, V. Ložek (3); Naiads from Czechoslovakia, J. Petrbok.

Slavakia.—Tortonian fossils, Čechovič & Seneš; New Pupilla from the Pleistocene of Slovakia, J. Petrbok (2); Some helvetian fossils of Southern Slovakia, J. Seneš (2); Helvetian fossils of S. Slovakia, J. Seneš (2); Oligocene of Slovakia, J. Seneš (4); Miocene molluscs from E. Slovakia, J. Svagrovský.

Yugoslavia.—Cretaceous molluses of Yugoslavia, M. Andelkovic; Gastropods from Jugoslavia, Albania and Macedonia, Jaeckel & Meise; Chondrodonta from the Yugoslavian Cretaceous, M. Pleničar; Two new Pyrgohydrobia species from Ohrid, Jugoslavia, P. Radoman; New molluses from the Miocene of Yugoslavia, P. Stevanović; Triassic cephalopods from Peca, Yugoslavia, L. Žlebnik.

Latvia.—Soil mollusca of Latvian Republic, Eglitis.

Mediterranean Islands.—New Clausilidae from Crete, F. Loosies; Pliocene lamellibranchs from Sicily, A. Malatesta; Pliocene and Quaternary molluses from Milazzo, Sicily, P. Mars (3); The Albinaria of the Dodecanese, K. L. Pfeiffer; Land molluses from Pantelleria Island, Mediterranean, C. Sacchi (1).

Russia.—New Eocene molluses from the Aral and Ferghana, Russia, A. Alexeiev: Mollusca of Sakeranl horizon in Abkhasia, Russia, Archvadze; Doricranites and other Triassic molluses from Turkmenia, Russia, T. Astakhova: New Carboniferous Imitoceras from Russia, E. Balashova; Mollusca in the layers of upper Crimea, Russia Triassic-Jurassic, Barkhatov; Mollusca in Fergan depression, Paleogen, Russia, Belskaya; Mollusca in Vetlyan deposits, Russia, Blom; Mollusca in Apt deposits, Kobra river, Russia, G. I. Blom (1); New Cretaceous Cardium from Caucasus region, Russia, N. Bobkova: Lower Jurassic mollusca in the river Okuma basin. Abkhazia, Russia, Bukiya; Mollusca in Datsk stratum of Megrelia upper, Paleocene, Russia, S. Bukiya (1); Mollusca in North-Western part of Abkhazia, Lias, Russia, Bukiya (2); Mollusca in upper Saratov deposits, Russia, Chibrikova; New Cretaceous ammonites from Crimea and Northern Caucasus, V. Drushchitz: Upper Miocene mollusca in Western Grazia. Dzevelya; Pliocene Cardiidae from Russia, A. Eberzin; Mollusca in the Vizlisk deposits, Russia, Einor; Mollusca in Western Primugodzav deposits, Russian Cretaceous, A. Ghlazunova; New Pectunculus from the Miocene of S.W. Russia, V. Ghoretzkii; Known and new mollusca for the Lower Kama river basin, Quarternary, Russia, Goretzki, G. I.; Mollusca in Black Sea deposits, G. Goretzki (1); Fossil molluscs in the valley of Kuban river, Russia, Kakhadze; Caspian molluscs in riverbed deposits of lower Dneper and Ingultz rivers, N. Karlov (1); Mollusca in the Artinsk layers of Central Ural, Khabakov; Bithynia leachi in the lakes of Krasnoyarsk Territory, U.S.S.R., N. Khristenko; New and known mollusca of the Paleogene in Middle Asia, Russia, Korobkov; Mesozoic mollusca from Preverkhoyansk deposits, Russia, Koshelkina & Tuchkov; Absence of Prosobranchia in the deposits of the lakes situated in the Tobol and Ishim basin, Russia, Kozlovokaya (1); Venus

sobieskii from the Miocene of the Russian Platform, L. Kudrin: Mollusca in the Tortonian (Oligocene) at the S.W. ridge of Russian platform, Kudrin (2); Mollusca in coal deposits of Sovetska Vyatke river, Russia, Larionova & Safonova: Molluses in Kimmeriïsk deposits, Russia, Lituinenko; First record of Pulchellia from the Cretaceous of the Caucasus, N. Luppov; New Cretaceous Cicatrites from the Caucasus, N. Luppov (1): New Permian ammonites from the Yurezan river, Russia, S. Maksimora: Mollusca of Konsk horizon, Kara-Bogaz-Gol, Russia, R. Merklin (1); Mesozoic mollusca of N. Taimyr, Russia, Miroshnikov, L. D.; Mollusca from the limestones of Primose Land, Russia, Moiseev; Paleogene molluses of Russia, Osipova & Liverovskaya; Mollusca of the lower-Permian of Dagoria, Russia, Permyakov: Prilukiella new pelecypod genus from Russian Triassic. M. Plotnikov: New Triassic lamellibranchs from Russia, M. Plotnikov (1); Tertary Teredo, Kuïbrshev, Russia, Y. N. Popov; Apsheron deposits mollusca of Ala-Dag, Russia, G. I. Popov (2): Namurian Goniatites of Middle Ural, Pronin; Spectoniceras versicolor from the Russian Cretaceous, Reyment, R. (1); New and known Carboniferous and Permian ammonites from S. Urals, Russia, V. Ruzhentzev (3); New ammonites from the Sakmara deposit, Kazakh, Russia, V. Ruzhentzev (4); Mollusca of Kara-Bogaz-Gol, Russia, Rikhtez; Mollusca in the chalk deposits of Oskolsk layer, Russia, Ryabikh; New Cretaceous and Triassic molluscs of Dzhungar depression, Russia, M. Saidov; Mollusca in Megozviena of Sir-Darva river plain, Samodurov; Molluses in the Aptian deposits of the Central Russian platform, Sazanova; Namurian mollusca, Russia, Semikhatova; Fossil mollusca of Karabogaz coast line, Russia, Sergeev; Mollusca from phosphoritic deposits of Karagadinsk Basin, Russia, Shatski; Mollusca in Sarisuïsk cupolas, Central Kazakhstan, Russia, Smekhov; Jurassic mollusca, Black Sea area, Russia, Sukharevich. P. M.: Devonian mollusca in Pripyatsk, Russia, **Tikhomizov**; Fossil Pelecypods at Dzhungaric border Cretaceous, Russia., Tuaev New Triassic molluscs from Okhotsk Sea area, Russia, I. Tuchkov; The species of Pseudomonotis in the Noriisk layer

upper Cambrian, Russia, Tuchkov (1); Mollusca of Apsheron deposits between Bolshi Uzen and Urala rivers, Russia, Vostryakov.

Turkey.—Pliocene mollusca of the Cuide peninsular, G. Chaput; On the Pliocene of environs of Antalya, Chaput & Darkot.

Syria.—Pila ovata from the Palestine Quarternary, N. Avnimelech; Two Triassic ammonites from Macedonia, W. Medwenitsch.

Irak.—New Cretaceous rudist from Persia and the Antilles, L. Chubb (1); Persia Tertiary Pectinacea, Eames & Cox.

Afghanistan.—Molluses from Afghanistan, S. Jaeckel (1).

Asia.—New Tibetan Ammonites of the Bajocian Jurassic, W. J. Arkell (2); Guinea, Angola, Mozambique S. Thomas and Principe, India and Timor, Azevedo & Medeiros: Mollusca in Schwagerine horizon, Donetz, Russia, Bankovski & Redichkin; New Triassic Unio species from Eastern Ural Mts., B. Chernuishev; Afghanistan a Jurassic marine fornula, R. Faron; Asiatic snails and liverflukes, S. B. Kendall; Mollusca in Triassic deposits of Eastern Trans-Baikal. Marinov: Fossil mollusca in some lake basins in Asia, G. Martinson (3); Mollusca in the continental deposits of Transbaical, Russia, G. Martinson (4); New Pelecypoda from the Miocene of Turkmenstan and W. Kazakhstan, Merklin & Nevesskaya: Carboniferous ammonites from Ferghana, Turkestan, Miklukho-Maklai, & Ruzhentzev; Known and new molluses from the Quaternary of Turkmenia, L. Nevesskaya; Mollusca in the volcanic muds at Elton lake and Transvolgian territory, Russia, Pravo-slavlev; Mollusca of Priverkhoyansk deposits, Pushcharovski; Size of White and Okhotsk sea mussel, Savilov (1); Molluscs of the Tarkhansk horizon of Asserbaidzhan, Sultanov; Development of non-marine fossil mollusc faunas in E. Asia, K. Suzuki; Distribution of Mollusca in bays of Okhotsk and Behring seas, Vinogradova; Mollusca in Apt-Senoman layer of Transcaspian, Vistelius; Mollusca in Kinsk deposits, Zhgenti.

Siberia.—Tertiary mollusca of Kazakhestan and Western Siberia, Lavrov, V. V.; New Cretaceous pelecypods from Sakhalin Island, Siberia, E. Liverovskaya; Probaicalia new gastropod genus from the Cretaceous of Siberia, G. Martinson; Notes on some Siberian slugs, H. W. Walden (3).

North Africa. — New Cretaceous species of Nerinea from Sinai, G. H. Awad; Praecardiomya new Cretaceous genus from Egypt, G. H. Awad (1); Cretaceous ammonites of the family Oosterellidae, found in N. Africa, Basse & Delga; South Tunis, Cretaceous of Granos Chott, Castany, Coque & Domerque; Algeria, a new ammonite fauna, David & Gornay; New Cretaceous Pycnodonta from Algeria, Freneix & Magné; N. African lower Cretaceous Nerinea, G. & J. Glacon; New Cretaceous molluscs from Sinai, Egypt, I. G. Mahmoud; Algerian 'Jurmatic' fauna, a preliminary note, D. Mongin; Tertiary molluscs from Basse-Kabylie, Algeria, P. Muraour.

## ORIENTAL REGION AND EAST INDIES

India.—New sub-genus and species of Astarte from the Jurassic of Cutch, India, S. Agrawal; Kutch, India, Jurassic species, G. K. Agrawal (1); Fossil Trigonia from Kutch, India, Agrawal (2); Utaturiceras new Cretaceous ammonite from India, C. Wright.

Ceylon,—A new Corilla from Ceylon, figs., G. A. S. Barnacle; New Miocene gastropods from Ceylon, P. Deraniyagala.

Indo-China.—Marine Mollusca, C. Dawydoff.

Java.—Revision of freshwater gastropods from Java, E. v. B. Jutting.

Japanese Subregion.-New Cretaceous molluses from southwest Japan, M. Amano; Species found in a deposit in Ofenata Bay, Northern Japan, T. Habe; Enidae from Japan, T. Habe (6); Ostrea konbo from the Japanese Cretaceous, I. & S. Hayasaka; New Tertiary molluses from Japan, K. Hirayama; Endemicism in Jamaican snails, W. R. Hunter (1); New Japanese Triassic pelecypods, K. Ichikawa; Trigonia from Mesozoic of Hokkaido, Japan, S. Imanishi: Miocene molluses from Kyoto Prefecture, Japan, J. Itoigawa; A new Miocene Japalion from the Jôban coalfield, Y. Kamada; Jurassic molluses from Japan, T. Kimura; A new form of

Apollon pusillus from Kii, Japan, T. Kira: Jurassic Trigonians from Japan, T. Kobayashi (2); New Cretaceous Steinmanella from Japan, Kobayashi & Amano; New Triassic Pectinidae from Japan, Kobayashi & Ichikawa; New Triassic pelecypods from Japan, Kobayashi & Ichikawa (1); New Triassic Entomonotis from Japan, Kobayashi & Ichikawa (2); Yaku Island land shells, T. Kuroda; New Miocene shells from the Noto Peninsula, Japan, K. Masuda: Miocene Pectinidae from Sendai, Japan, K. Masuda (3): New Cretaceous ammonites from Japan, T. Matsumoto (1); Cretaceous molluses from Hokkaido, Japan, Matsumoto & Obata; Jurassie molluses from Hashuira district, Japan, K. Mori; New and known Triassic Pelecypoda from Kyoto Prefecture, Japan, K. Nakazawa: New Japanese Pliocene gastropods, Y. Otuka; List of Miocene molluscs from Sendai, Japan, K. Oyama; Mollusca of Aogasima, Japan, K. Oyama (1): New species of Cymatoceras from the Japanese Cretaceous, Ozaki & Katto; New ammonite, Damesites, from Japanese Cretaceous, Saito & Matsumoto; Stephanoceras from the Jurassic of Japan, T. Sato (1); New Tertiary Trochidae from Japan, T. Shuto; Anatomy of sane Japanese Epitoniidae, I. Taki; New Miocene Platyodon from Japan, Uozumi & Fujie; Studies in Japanese slugs, Yamaguchi & Habe.

Korea.—Note on four Korean snails, T. Habe (7).

## AUSTRALIAN AND POLYNESIAN REGION

Australia and Tasmania.—Molluscs from the Cainozoic of northern Tasmania, Gill & Banks; Fresh water molluscs from Prof. Gisten's Australian Expedition 1951–1952, B. Hubendick; Fluviatile and lacustrine Pelecypoda from Victoria, Australia, R. C. Kershaw; Rissoinidae and Rissoidae from the Solanderian and Dampierian zoogeographical provinces, C. F. Laseron; Pliocene molluscs of Adelaide Plains, Australian, N. H. Ludbrook; New Devonian molluscs from Victoria, Australia, J. A. Talent; Siluro-Devonian molluscs from Victoria, Australia, Talent & Philip; New Australian gastropods, L. Woolacott (1).

New Zealand.—Taxonomy of the New Zealand species of Limnaea, Myxas and Physastra, R. K. Dell (1); Pleistocene molluscs from Whakatane, New Zealand, C. Fleming; Mode of occurrence in New Zealand (diagrams), J. E. Morton; New Tertiary Olividae from New Zealand, O. P. Olson; Small land snails from Kaitaia, North Island, New Zealand, L. Price; List of molluscs from Stewart Island, New Zealand, E. K. Smith; Molluscs from Leigh, New Zealand, P. Warren.

Pacific Islands.—New Pyrene from New Guinea, A. C. v. Bruggen; Pulmonata and Prosobranchia of New Caledonia, A. Franc (2); Occurence of Oncomelania quadrasi in the Philippines, B. Hubendick (1); Hawaii, the first Helix aspersa, Y. Kondo; Helix aspersa in Hawaii, Y. Kondo (2); Clausiliidae from New Guinea, F. Loosjes (1); New Guinea freshwater mussels, figs., D. F. McMichael; Granocardium pomeyroli from the Cretaceous of New Caledonia, R. Pomeyrol.

## ETHIOPIAN REGION

Ethiopia.—Ethiopian land and freshwater mollusks, Verdcourt.

East Africa.—Mediterranean land snails in Tropical E. Africa. C. R. Boettger (4); Dahoney, a new lower Eocene Turritella, Ph. Brébion; First record of Eocene molluscs from Italian Somaliland, E. Giannini; New Gastropoda from Tanganyika, E. Africa, H. Knipper; Anatomy of Tanganyika, Ellobiidae, Knipper & Meyer; Differentiation of East African species of Bulinus, W. F. J. McClelland; East African mollusca, additional notes, T. Pain (2); Lists of freshwater molluscs from the Coast province of Kenya, C. Teesdale; Belgian Congo, a new subspecies of Pila africana (Masteres), L. A. W. C. Venmans (1); Somalian land and freshwater mollusks, Verdcourt; Physopsis globosa found in Italian Somaliland, E. Zavattari.

West Africa.—List of molluses not previously recorded from the Gold Coast, Buchanan & Anderson; Short list of molluses from the Cameroons, W. Africa, S. Jaeckel (2); Bilharziasis in Angola, Janz & Carvalho; New terrestrial Gastropoda from Fernando Póo Island, W. Africa, López &

Rocandio; Gold Coast, resistance of a snail to infection, F. S. McCollough; Transmission of schistosomiasis by Bulinus spp. in the Gold Coast, F. McCullough (2); Quaternary species from Senegal, M. Paulus (1); New Cretaceous ammonites from Nigeria, R. Reyment.

Central Africa.—Atlas—Western Sahara, occurrence of Ermoceras, Arkell & Lucas; New Cretaceous Plicatounio from French Sudan, J. Goumard; Molluscs of Italian Somaliland, Hemming & Verdcourt; Sahara fossils, A. L. de Lapparent; Cretaceous and Tertiary nautiloids from Angola, Miller & Carpenter; Mollusc vectors of bilharziasis in Zambezia, T. d. Morais; Tanganyika, a new Gulella, B. Verdcourt (1); Tanganyika district, species of Primigulella, Verdcourt & Venmans.

South Africa.—Commensal Oligochaeta on freshwater snails in S. Africa, F. A. H. Bayer; Rhodesian Permian Hydrobia, figs., L. R. Cox (1); Oyster culture at Knysna, S. Africa, P. Korringa (1).

Atlantic Islands.—Variation in Cerion from the Bahamas, Mayr & Rosen.

#### NEARCTIC REGION

Greenland.—Pleistocene fossils from Pearyland, North Greenland, D. Laursen (1).

Alaska.—Some land and freshwater mollusks of the Artic slope, figs., G. D. Hanna (1).

Canada.—British Columbia and range to other countries of *Homolsomites*, R. W. Imlay (1).

North America.—North American Lioplax and Campeloma, Clench & Turner (1); Distribution of American Pennsylvanian Cephalopods, A. G. Unklesbav.

United States as a whole.—McDonald Cirque, Montana and Discus brunsoni, Berry, R. B. Brunson; On some American Planorbids, G. Ranson (3); New Cretaceous bivalve genus from Mississippi, L. W. Stephenson.

North Eastern United States.—Pennsylvanian faunas of the Amsden formation, Wyoming, C. A. Burk; Molluses from a fish-weir constructed in the recent past, and excavated at Boston, U.S.A., W. Clench; Pleistocene molluses from Fairfield County, Ohio, La

Rocque & Conley; New Cretaceous lamellibranchs and an ammonite from Alabama, U.S.A., L. Stephenson (1).

South Eastern United States .- Virginia, distribution of land snails and plant association, J. B. Burch; A survey of freshwater mollusca of the rivers of northern Florida, W. J. Clench (4); Fresh water molluses of Alabama. Georgia and Florida, Clench & Turner: Land snails of Shenandoah National Park, Blue Ridge Mountains, northern Virginia, L. Hubricht; Land snails collected in Louisiana, L. Hubricht (2); Florida, a living example of Panope bitruncata Conrad from St. Augustine, fig., M. C. Johnson; Texas lower Turomian ammonites, Kummel Decker: Lower Turonian Ammonites from Texas and Mexico, Kummel & Decker: New species of Olivella from Florida, recent and tertiary, A. A. Olsson; Texas, Beach drift Polygyridae, Pilsbry & Hubricht; Ventridens inter-textus eutropis in Tennessee living examples, M. C. Teskey; Nucinellidae from the Eocene of Oregon, H. E. Vokes; Checklist of cephalopods of Florida, G. Voss.

Western United States.—Inyo Co., California, Panamint Lake recently extinct molluscan fauna, J. L. Baily inr.; Wyoming fauna from the Amsden formation of the Pennsylvanian, C. A. Burk; New Nassarius species from Pliocene of California, G. Kanakoff; Jurassic Nautiloids from Western N. America, B. Kummel; Habitat of mussels of Lake Texoma, Oklahoma, U.S.A., Riggs & Webb; Nonmarine mollusca from the Saratoga Valley, Wyoming, Miocene, L. S. Russell; Western United States and the upper Triassic fauna, N. J. Silberling.

#### CENTRAL AMERICAN REGION

Mexico.—Pleistocene molluses from Punta China, Mexico, W. Emerson; Molluses from Cozumel Island, Quintana Roo, Mexico, E. Hidalgo; List and plates of Cretaceous molluses from Mexico, L. Kellum; Mexican Lower Turomian ammonites, Kummel & Decker; New molluses from Mexico, H. A. Pilsbry (3). New species of Cyclophoridae from Mexico, A. Solem.

Barbados.—Land shells of Barbuda, W. J. Clench (6).

Bahamas.—New Cerion from the Bahamas, W. Clench (5).

Cuba.—Cretaceous Rudists from the Antilles, L. Chubb; Cuban malacological notes, M. H. Jacobson; Puerto Rico—a new Macroceramus, figs., M. K. Jacobson; Trigonia from the Cayetano formation of Cuba, K. Krommelbein.

Jamaica.—New Cretaceous rudists from Jamaica, B.W.I., L. Chubb; New Cretaceous rudist from Jamaica and Persia, L. Chubb (1); Jamaican snails and endemicism, W. R. Hunter (1).

Trinidad.—Fossils from the Lower Cretaceous, L. R. Cox (2); Barremian Ammonites, figs., R. W. Imlay.

## SOUTH AMERICAN REGION

Bolivia.—A new locality for Systrophia stenogyra zischkai, W. Blume; A new species of Clausiliidae, W. Weyrauch (1).

Colombia.—Strophocheilidae from S. America, J. Bequaert; Cretaceous cephalopods from Muzo mines, Boyacá, Colombia, S.A., H. Bürgl; Two new varieties of Dufreynoya texana from the Cretaceous, Colombia, H. Bürgl (1); Cretaceous ammonites of Colombia, H. Bürgl (2); A new species of Synapterpes, T. E. Crowley; On Pomacea from Colombia, T. Pain; La Nueva, Rio Truando Choco, N. Western Colombia, a southern record for Pomacea cumingi (King), H. A. Pilsbry (2).

British Guiana and Surinam.—Melanidae, T. Pain (3); Surinam Melanidae, T. Pain (3).

Venezuela.—Mollusca from the Cogollo and La Luna Cretaceous formations, Rutsch & Salvador.

Brazil.—Molluses of Brazilian shellmounds, J. Bigarella; Molluses of shellmounds of Santa Catarina, Brazil, Bigarella, Tiburtius & Sobanski; Some snails from the Amazon region and new species, F. Haas; Molluses from the island of Fernando de Noronha, Brazil, Lopes & Alvarenga; Two new slugs from Brazil, E. & E. Marcus (2); New species of Australorbis from Brazil, Paraense & Deslandes (2); Brazilian species of Australorbis, Paraense & Deslandes (3); On a collection of South Brazilian Artemon, A. Solem (1). Non-marine mollusea from Salabra, Matto Grosso, Brazil, A. Solem (1).

Peru.—Jurassic mollusca from Peru, L. R. Cox (4); Peru, a new Bostrya, figs., H. A. Pilsbry; Naesiotus and other Bulimulidae from Peru, W. Weyrauch; Peru, Paleocene and Eocene Venericardia, figs., P. Verástegui; Peru, a new species of Clausiliidae, W. Weyrauch (1); New land shells of Peru, W. K. Weyrauch (2).

Chile,—New Cretaceous molluses from Chile, Leanza & Castellaro; Araucania, new genus of Gastropoda from Chile, J. Parodiz; Key to species of Pacificor from the Paleocene and Eocene of Western South America, P. Verástegui.

Argentine.—Occurrence of Nenia, Anon (3); New ammonite family from the Argentine Jurassic, Maubeuge & Lambert; New Bulimulus species from Argentina, J. Parodiz (1); New Bulimulus species from Argentina, J. Parodiz (2); Two new Epiphragmophora from Argentina, J. Parodiz (3); New species of Triphora, first record of genus from Uruguayan area, J. Parodiz (4); Tentaculites from the Ordovician of Argentina, C. Rusconi; Cambrian gastropods from Mendoza, Argentina, C. Rusconi (1); List of fossil molluses from Mendoza, Argentina, p. 22–26, 53, 54, C. Rusconi (2); Ulpia new genus of Odontostominae, from Argentina, M. I. H. Scott (1).

Patagonia.—Leopoldia paynensis from the Cretaceous of Patagonia, A. Cecioni; Kossmaticeratidae in Patagonia, G. Cecioni (1).

Paraguay.—New Aylacostoma species from Paraguay, M. I. H. Scott.

# (β) Marine ARCTIC

Arctic Seas.—A new record for Iceland, the occurrence of Sipho verkrüzeni, Kobelt, I. Óskarsson (2); Registration of Cephalopods, new for the fauna of Far-East seas of Russia, Akimushkin (2); Iceland bivalves, I. Óskarsson; Lamellisbranchs in Iceland, I. Óskarsson; Cardium edule in Icelandic waters, I. Óskarsson (1).

#### NORTH TEMPORATE

North Atlantic.—Distribution of a north Atlantic pelagic squid, B. Muus; Four Atlantic species of Macrotritopus, W. J. Rees (1); Chlamys opercularis in Danish and Faroese waters, E. Ursin.

Eastern North Atlantic .-- R. Fal, Cornwall and Callochiton fulvus, R. H. Baird; Molluscan fauna of the Schelde Estuary, Belgium, W. G. Beiftinck; New subspecies of Neptunea decemcostata, A. H. Clarke, jnr.; Marine molluscs from the North Devon coast, A. Cutcliffe: Intertidal mollusca of Spanish Northern coasts, E. Fischer-Piette; Short list of Nudibranchs of Saint Malo Bay, A. Franc; Pecten maximus in Irish waters, F. Gibson; List of molluscs collected off Port Erin, Isle of Man, N. Jones; North Atlantic species of Cylichna, H. Lemche; Chlamys varia from Northern France, Letaconnoux & Audouin; Opisthobranchs of Lancashire and Cheshire, N. F. McMillan (1); English Channel (Fal R.) Callochiton fulvus (Wood), N. F. McMillan (3); Akera bullata first record from Plymouth area, Morton & Holme; Two mollusca of the rockpools of the Eastern Pyrenees, Ohm & Remmert; North Sea plankton, C. B. Rees; Northern Norway, new records of marine Mollusca, T. Soot-Rven: List of intertidal species on the Irish Coasts and their distribution, Southward & Crisp.

Western North Atlantic.—Chesapeake Bay, notes on Amygdalum papyria Conrad, J. F. Allen (1); Additional marine mollusca at Cape Ann, Massachusetts, R. W. Dexter (1); C. B. Adams' species described from the Atlantic, figs., R. D. Turner (2).

Gulf of Mexico.—New ectoparasitic molluses from Mexico, S. Berry; Caribbean Pteropoda, A. D. Howard; Bermuda, a new Glycymerid, D. Nicol (3).

Baltic Sea.—Molluses from the Baltic, B. Forsman.

Mediterranean.—Mitra (Episcomitra) zonata in the Bay of the Ciotat, S. France, J. Devidts; Mollusca in Black Sea sediments, Fedorov, P. V.; Lamellibranchs from the Thyrrhenian Sea, off the Tuscan coast, B. Gianfranco; N.W. African Neritina, F. Llabador; Observations on certain species, P. Mars (1); Gastropoda from the Leghorn coast, Italy, Sordi & Majidi.

Caspian Sea.—Mollusca of Apsheron, Caspian Sea, Klenova.

Eastern Pacific.—Some problems in West Coast of America, Naticidae, J. Q. Burch (3); New species of Dentalium off

west coat of Mexico, W. K. Emerson (1); Eastern Pacific species described by C. B. Adams, figs., R. D. Turner (1).

Western Pacific.—Gastrochaena rostrata and Barnea manilenses from the Philippines, G. Ablan (1); Geographical distribution of Cephalopods in Far-Eastern seas, Akimuskin; Tosa Bay, a new locality for Cypraea guttata Gmel., M. Azuma; Euselenops luniceps from Japan, K. Baba; Note on Philippine species of Ostrea, Blanco, Villaluz & Montalban; Notes on New Zealand Rissoina spp., A. K. Hipkins; New Conidae from Japan, T. Kuroda (1); A new Lischkeia from Japan, Kuroda & Kawamura; Okhotok Sea: a new Volutharpa, Kuroda & Kinoskita: New Rissoinidae and Rissoidae from Northern Australia, New Guinea and Christmas Island, C. F. Laseron; Mytilidae from the Peronian zoogeographical Province, N.S.W. Australia, C. Laseron (1); New genus of the Nassariidae from N.S.W. Australia, C. F. & J. Laseron.

### TROPICAL AND SUBTROPICAL

Western Atlantic. — Polyplacophora from southern South America waters, Z. Castellanos; Aplysia, Berthella and Pleurobranchaea from Brazil, E. & E. Marcus; Two Atlantic Onchidella species, from Brazil and Tenerife, E. & E. Marcus (1).

Indian Ocean.—Lambis spp. from Indonesia, L. J. Butot; First record of Cassis cornuta from Bombay, India, V. Chari; Mollusca of the Red Sea, A. Franc (1); Kerguelen and Heard Islands Lamellibranchs, J. M. Gaillard; New Teredo sp. from South India, Nair & Gurumani; Mitridae from Indian waters, H. Ray; Seychelles Pelecypoda, Rost & Soot-Ryen.

## SOUTH TEMPORATE AND SUBANTARCTIC

Indo-Pacific Waters.—Malaya Cypraea, R. J. Griffiths; New Northern Australian volute, T. Iredale.

Western Pacific, North.—Two squids from Japan, E. Iwai; Galiteuthis armata from Japan, T. Hikita.

Western Pacific, South. — Amoria grossi first record from N.S.W., Australia, J. Allan; List of intertidal molluses, p. 331, 342, on islands off the Queens-

land coast, Australia, Endean, Stephenson & Kenny; Archibenthal Mollusca of New Zealand, R. K. Dell; New Caledonian Naticidae anatomical studies, figs., J. Risbec (1); Giant Clams of Australian Seas, C. M. Yonge.

Eastern Pacific, North.—New species of Olivella from Costa Rica and Panama, A. A. Olsson; Armina digueti new species from the Gulf of California, A. Pruvot-Fol (1).

Eastern Pacific, South.—New species of Olivella from Ecuador and Venezuela, A. A. Olsson.

South Pacifc.—New Aeolidian from Kaukura Island, South Pacific, A. Pruvot-Fol (2).

Southern Ocean.—New molluses from Bass Strait, South east of King Island, Victoria, Australia, C. J. Gabriel.

#### (B) Geological

Quaternary.—Pila ovata from Palestine, N. Avnimelech; Quaternary mollusca of Lake Massacinecoli, Blanc, Settepasci & Tongiargi; Holocene, Belgian Congo, Braucourt; Quarternary Tongiargi; Holocene, mollusca of the Caspian Sea, P. Fedorov (1); Molluses from aboriginal sites in Tasmania, Gill & Banks; Known and new Quaternary molluscs from Kama river basin, Russia, G. Goretzki; Slug granules from post glacial deposits in the Lea Valley, J. F. Hayward; Fresh water molluses from the late and Post Glacial of Schleswig-Holstein, S. G. A. Jaeckel; List of Holocene molluses from Wateringbury, Kent, England, M. Kerney; Molluses from Milazzo, Sicily, P. Mars (3); Molluses from Bas-Languedoc, France, Mars & Ottmann: New Parastrophia from Italy, M. Moncharmont Zei; Known and new mollusca from Turkmenia Quartenary, Nevesskaya, L. A.; Quaternary of Senegal, 'List of species, M. Paulus (1); Bohemian Holocene molluses, with some new forms, J. Petrbok (1); Black Sea and Caspian, G. I. Popov; New and known mollusca in Black Sea and Caspian Quaternery deposits, G. I. Popov (1); A Post-Pleistocene Natica from Netherlands, Pouderayen: Quaternary fauna of Austria, R. Sieber: Development of non-marine fossil molluscan faunas in Eastern Asia with list of Quaternary species, p. 130, 132, K. Suzuki.

Pleistocene.—Comparison of Pleistocene molluscs from Britain and the Mediterranean, D. Baden-Powell; Pleistocene, Belgian Congo, Braucourt; Molluscs from the Loess in Donau, Germany, M. & K. Brunnacker; Plio-Pleistocene molluses from Suffolk, England, P. Cambridge; Pleistocene mollusca at Stutton Ness, Suffolk, A. G. Davis; List of molluses, p. 337, from Punta China, Mexico, W. Emerson; Pleistocene mollusca at Kapp Wijk, Westspitsbergen, Feyling-Hanssen: Late Pleistocene fosfrom Vestspitsbergen, Feyling-Hanssen (1); Molluscs from Whakatane, New Zealand, C. Fleming; Note on Pleistocene molluses from Ohio, La Rocque & Conley; Fossils of Ohio, U.S.A., La Rocque & Marple; Pleisto-cene of Pearyland, D. Laursen (1); New Pupilla from Slovakia, J. Petrbok (2); Pleistocene in the Netherlands, Vlerk & Florschütz: Pleistocene Picidium vincentianum and associated species from Ehringsdorfer, H. Zeissler.

Tertiary.—Tertiary mollusca of Sakhalin, Alekseïchik, Kuzina & Ratnovski; Lima species from Japan, S. Aoki; List of molluscs from the Tertiary of St. Gervais, France, P. Balavoine; Note on two Terteriary species of Arion described by Wenz, 1911, H. Bräm: Mediterranean Region Tertiary Pectinacea, Eames & Cox: Bivalves from Zeeland, Holland, H. v. Halm; New Tertiary molluses from Hikoshima, Japan, K. Hirayama; Glycymera from Japan, S. Kanno; Tertiary Mollusca, Zaporozhe, Russia, Karlov & Gryaznov; Molluses of Kezakhstan and Western Siberia, V. Lavrov; Tertiary molluses of Sintz-yan, Mongolia, G. Martinson (2): Tertiary molluses in north-western Siberia, Merklin; Tertiary nautiloids from Angola, Miller Carpenter; Molluscs from the Ofanto valley, Colabria, Italy, M. Moncharmont Zei (1); Tertiary molluses from Basse-Kabylie, Algeria, P. Muraour; New Tertiary species of Baryspira (Olividae) from New Zealand, O. P. Olson; New species of Olivella from the Pliocene and Miocene of Florida, Costa Rica and S. America, A. A. Olsson; New species of Chlamys from the Japanese Tertiary, M. Shibata; New Pliocene-Miocene Trochidae from Japan, T. Shuto; Tertiary Lamellibranchia from the Vienna area, Austria, R. Sieber (1); Non-marine fossil molluscan faunas in E. Asia with lists of Tertiary species pp. 120-129, 131, K. Suzuki; Inoceramus from Trans-Carpatha, Vyalov & Pasternak.

Cenozoic.—Cenozoic Belgian Congo, Braucourt; Caenozoic molluscs of Caenozoic clays in mid Pre-Ob basin, Feniksova, V. V. List of molluscs, p. 19, from the Mowbray Swamp, Tasmania, Gill & Banks.

Pliocene.—A new upper Pliocene Pyrene from the Netherlands, C. O. v. R. Altena (1); Comparison of British & Mediterranean Pliocene molluscs, D. Baden-Powell; Mio-Pliocene Mollusca of Buton, C. Beets: Pliocene mollusca of Lake Massacinccoli, Blanc, Settipasci & Tongiargi; Pliocene molluscs of Cuide Peninsula, Turque, G. Chaput; Nassarius and Ancilla new species from Portugal, L. Cox; Suffolk-Pleistocene mollusca at Stutton Ness, Suffolk, A. G. Davis; Cardiidae from Russia. A. Eberzin: Pliocene molluscs of Evskin and Caspia and their relationships Russia, Eberzin (1); Pliocene mollusc, from Mirabel, Ardèche, France, Grangeon & Jodot; New Nassarius species from California, G. Kanakoff; Upper Pliocene mollusca in Guria deposits. Russia, Kvaliashviti; Pliocene molluscs of Adelaide Plains, Australia, N. H. Ludbrook; Lamellibranchs from Sicily, A. Malatesta; Pliocene molluscs from Milazzo, Sicily, P. Mars (3); Valenciennius annulatus in the Pliocene of Roumania, P. Niță; New Pliocene gastropods from Japan, Y. Otuka; Unio wetzleri flabellatiformis and Melania escheri from the Roumanian Pliocene. M. Paucă; New species of Anodonta from the Roumanian Pliocene, Rosculescu-Soverth; Mollusca from the Pliocene of the Black Sea basin, Sukhova; Mollusca Aktchagil in Moldavia middle Pliocene, Viktorova & Kovalevski.

Neogene.—Neogene molluses of Minbulak, Russia, Gramm.

Miocene.—Miocene molluscs from Lisbon, J. Berkeley-Cotter; Plio-Miocene, Belgian Congo, Braucourt; New molluscs from Hungary, I. Csepreghyne-Meznerics; Molluscs from Hungary, I. Csepreghyné-Meznerics (1); Miocene, West Africa, Dauteville & Roger; New Miocene species from the environs of

Gambotta, David & Judot: New gastropods from Ceylon, P. Deraniyagala; Molluscs in Guriïsk layers of Western Grazia, Dzvelaya & Maglaperidze; Miocene Pectinacea from East Africa. Persia and the Mediterranean region, Eames & Cox; Pectunculus from the frontiers of the South West Russian Platform, V. Ghoretzkii; Turridae of Belgium and the basin of the Loire. M. Glibert: New and known molluses from Kyoto Prefecture, Japan, J. Itoigawa; A new Japelion from Japan, Y. Kamada: Miocene of Poland around Andrychow, W. Krach & W. Nowak; Venus sobieskii from the Russian Platform, L. Kudrin; Unio wetzleriformis new species, from the Brad-Baia Basin, Rumania, N. Macarovici; Molluscs from the Miocene of Virginia, U.S.A., J. McLean: Japan, new Miocene species from the Nota Peninsula, K. Masuda; Miocene Pectinidae from Sendai, Japan, K. Masuda (1); New Pelecypoda from Turkmenstan and W. Kazakhstan, Merklin & Nevasskaya; Lists of molluses from Sendai, Japan, K. Oyama; Anomia fossil byssus from Germany, A. Papp (2); Miocene molluses of South Jutland, L. B. Rasmussen; New Miocene species from Wyoming, L. B. Russell; Ceratites species from the Trochitenkalk of the Saar, Germany, E. Schneider; Miocene molluscs from Porto Santo Island, N.E. of Madeira, G. H. da Silva; Miocene molluscs from Prince Island, Gulf of Guinea, G. H. da Silva, (1); New molluscs from Yugoslavia, P. Stevanović; Molluscan fauna of E. Slovakia, J. Svagrovský; New Platyodon from Japan, Vozumi & Fujie.

Oligocene.—Mollusca in Mandrikovsk oligocene, Karlov; Oligocene fossils of South Slovakia, J. Seneš (4); Potamides from the Oligocene of Hungary, L. Strausz; Upper Oligocene of Hungary, L. Strausz (1).

Eocene.—New molluses from the Aral and Ferghana, Russia, A. Alexeiev; Lower Eocene Turritella from Dakomey, Pb. Bribion; A new Eocene genus from Pakistan (Indovoluta), F. E. Eames; First record of Eocene molluses from Italian Somaliland, E. Giannini; An Eocene Aturia from Cyrenaica, figs., O. Haas (1); Eocene (Lutetian) fauna, P. Jodot; New Aturia from the Japanese Eocene, T. Kobayashi (1); Gastropoda from Lutetian of Rocca Bernarda, Udine,

Italy, B. Martinis; Nucinellidae from the Eocene of Oregon, H. E. Vokes.

Cretaceous .-- New Cretaceous molluses from southwest Japan, M. Amano: New varieties of molluscs from Yugoslavia, M. Andelkovic; Radiolitidae from the North Pyrenees, G. Astre (1); Matheronia semirugata from Roumania, Atanasiu & Semaka; New species of Nerinea from Sinai, G. H. Awad: Praecardiomya siniatica new genus from the Egyptian Cretaceous, G. Awad (1); Cretaceous ammonites of the family Oosterellidae, in North Africa, Basse & Delga; New Arcomytilus from Spain, J. Bataller; Spanish Mollusca, J. Bataller (1); New Cretaceous Loricata from Sweden, J. Bergenhayn; New Belemnites from West Greenland, T. Birkelund; New species of Cardium from Caucasia, N. Bobkova: Cephalopods from Muso mines, Boyacá, Colombia, S.Am., H. Bürgl; Two new varieties of Dufreynoya texana from the Aptian, Colombia, H. Bürgl (1); Cretaceous Pulcheliidae from Colombia, H. Bürgl (2): Ammonites from the Wiltshire Gault, R. Casey; Albian fossils in a new deposit in the Corbieres orientales, Casteras & Villatte; Tortonian fossils, Slovakia, V. Čechovič & J. Senes; Leopoldia paynensis from Patagonia, G. Cecioni; Cretaceous Rudist assemblages from the Antilles, L. Chubb; New Cretaceous rudists from Jamaica B.W.I.. L. Chubb; Lower Cretaceous fossils from Pointe-à-Pierre, Trinidad, L. R. Cox (2); Some Senonian fossils of Souk Ahras (S. Constantinois), L. David; Coniacian of Algeria, a new Ammonite fauna, David & Sornay; France, Cretaceous Belemnite, M. Delattre; New ammonites from the Crimea & Northern Caucasus, V. Drushchitz: Lower Cretaceous mollusca of Crimea, Eristavi; New Pycnodonta from Algeria, Freneix & Magné; New varieties of Lepthoplites from Kopet-Dagha, Russia, A. Ghlazunova; Aptian Nerinea of N. Africa, G. & J. Glacon; New Plicatounio from French Sudan, J. Goumard; Tortonian fossils of Kosihoviec, Slovakia, V. Hans; Ostrea konbo from Hokkaido, Japan, I. & S. Hayasaka; Ammonite from Marin County, California, U.S.A., L. G. Hertlein; Barremian ammonites (Early Creta-ceous) from Trinidad, figs., R. W. Imlay; List of molluscs, from the Aurora limestone, Mexico, L. Kellum; New

Steinmanella from Japan, Kobayashi & Amano: New molluscs from Chile. Leanza & Castellaro; New pelecypods Sakhalin Island, Siberia, E. Liverovskaya: First record of Pulchellia from the Caucasus, N. Luppov; New Cicatrites from the Caucasus, N. Luppov (1); Cretaceous molluses from Moghara Massif, Sinai, I. G. Mahmoud; New and known molluses from Sinai, Egypt, I. G. Mahmoud: Probaicalia new gastropod from Siberia, G. Martinson; New Cretaceous ammonites from Japan. T. Matsumoto (1); Cretaceous molluscs from Hokkaido, Japan, Matsumoto & Obata: List of Ammonites and lamellibranchs from the Gault at Sevenoaks, Kent, England, R. Milbourne; Cretaceous nautiloids from Angola, Miller & Carpenter; New Cymatoceras from Japan, Ozaki & Katto; New Engonoceras from Oregon, U.S.A., E. Packard; Senonian Beograd. M. Pasic: East Serbia. Senonian fauna, D. Pejović; Chondrodonta munsoni from the Yugoslavian Cretaceous, M. Pleničar; Campanian molluses of N.W. Europe, L. v. d. Poel; Granocardium pomeyroli from New Caledonia, R. Pomeyrol; New am-monites from Nigeria, R. Reyment; Spectoniceras versicolor from Russia, R. Reyment (1); Mollusca from the Cogollo and La Luna formations of the Cretaceous of Western Venezuela. Rutsch & Salvador; New and known molluses of Dzhungar depression, Russia, M. N. Saidov; Uncoiled ammonites from the Lower Cretaceous of S. E. France, S. Sarkar; San Francisco, Scklosken, Boriella & Imlay; Molluses from the Corbières, France, P. Sénesse; New Cretaceous pelecypod from Mississippi, L. W. Stephenson; New lamellibranchs and an ammonite from the Cretaceous of Habama, U.S.A., L. Stephenson (1); Upper Cretaceous gastropods in the S. of France, G. Termier; Côte d'Or Bathonian fossils, Tintant & Joly; Molluses from the Planenschen Grundes near Dresden, K. A. Tröger; Mollusca in lower Cretaceous (Alba), Mongolian Republic, Turishchev; New Cretaceous ammonite Utaturiceras, C. W. Wright.

Jurassic.—New sub-genus and species of Astarte from Cutch, India, S. Agrawal; New Jurassic mollusca from Kutch, India, S. K. Agrawal (1); New species of Trigonia from Kimmeridgian,

Kutch, India, S. Agrawal (2); New genera of Jurassic ammonites, W. J. Arkell (1); Bajocian ammonites from Tibet, W. J. Arkell (2); Jurassic mol-lusca of Tungussian-Vilyuïsk deposits, Russia, Arsenyev; Upper Jurassic molluscs in the mid-course of rivers Agstev and Gasan-Tu, Armenia, Russia, Atabekyan: Lists of Mollusca from Cardó. Spain, J. Bataller (2); Jurassic Mollusca of the Northern Caucasus, Bryun; Creniceras crenatum from the French Jura, A. Caire: Creniceras crenatum from Turcenay, France, A. Caire (1); Oxford, the occurrence of Creniceras crenatum Brug, A. Caire (3); Corallian fossils from Beckley, Oxfordshire, T. H. Callomon; New molluses from Spain, Castell, Concha & Revilla; New Jurassic Liogryphaea species, France, Charles & Maubeuges (1); Peruvian mollusca from the Jurassic, L. R. Cox (4); Lamellaptychus, Punctaptychus and Laevilanellaptychus from Italy, Cuzzi: List of Jurassic ammonites from Mexico, H. Erben (1); Ammonites found for the first time at Deux-Sèvres, France, J. Gabilly: Ammonites from the Dogger of the upper Rhine, H. Illies; Mollusca of the middle Jurassic. Khain. Shardanov, Kasimova; List of molluscan fossils from the Torinosu limestone, Japan, T. Kimura; Trigonians from Japan, T. Kobayashi (2); Jurassic cephalopods of Transcaucasus, Krimgoltz; Trigonia from the Cayetano Formation, West Cuba, K. Kröm-melbein; Western North America, three new Jurassic Nautiloids, B. Kummel; Upper Jurassic Ammonites of Donetz, Russia, Makridin & Zinov-yev; Yebisites new Jurassic ammonite from Japan, T. Matsumoto; New and deformed ammonites from France, P. Maubeuge: New ammonites from France, P. Maubeuge (1); New ammonites from Luxembourg, P. Maubeuge (2); New ammonites from Luxembourg, P. Maubeuge (3); Upper Oxford in the Paris basin, P. L. Maubeuge (4); New ammonite family from the Argentine Jurassic, Maubeuge & Lambert; Molluses from Hashiura district, Japan, K. Mori; New species of Chlamys from Yorkshire, England, J. Neale; Aptychus fauna from the Roumanian Jurassic, Răileanu, Bădălută & Pelin; New Velata from Greenland, A. Rosenkrantz; Ammonites from Switzerland, W. Rüefli; Correlation of Japanese Jurassic by

means of ammonites, T. Sato; Stephanoceras sp. from Japanese Jurassic, T. Sato (1); Strenoceras in ammonites of the 'Dogger'—sections, figs., O. H. Schindewolf; Molluses from the Black Sea area, Russia, P. Sukharevich; A middle Jurassic superspecies of oyster, Sylvester-Bradley; List of Oxfordian ammonites from Dijon, France, Tintant & Larcher; Mollusca of lower Jurassic, Anabar, Russia, Voronetz & Laptinskaya (1); Stephanocerataceae and Perisphinctaceae, phylogeny of ammonites from the Dogger, G. Westermann; Jurassic species of Sphaeroceras and Chondroceras in Europe, G. Westermann (1); New ammonites from Blumberg, Germany, A. Zeiss.

Lias.—Revision of Liassic ammonites described in the monograph by P. Reynès, 1879, D. Donovan; Ammonite succession in the Blue Lias of Keynsham, Somerset, England, D. Donovan (1); Liassic ammonites from Isle of Raasay, Inner Hebrides, M. Howarth; Mid Lias (Pliensbachian), new ammonite find in the Domerian, E. Lörcher; Ammonites from the Lias of S.W. Germany, O. Walliser.

Post-Triassic.—Nautiloid genera, B. Kummel (1).

Trias.—Doricranites and other molluses from Turkmenia, Russia, T. Astakhova; New Unio species from the Eastern Urals, B. Chernuishev; New Triassic gastropods from Germany, H. Eckert; New Triassic Japanese pelecypods, K. Ichikawa; Mollusca from Russia, Kiparisova, L. D. & Popov, Y. N.; New Upper Triassic Pectinidae from Japan, Kobayashi & Ichikawa; New Entomonotis spp. from Japan, Kobayashi & Ichikawa (2); New Triassic Pelecypoda from Japan, Kobayashi & Ichikawa (1); Two ammonites from the Debar-zone of the Pelagoniden. Macedonia, W. Medwenitsch; Prilukiella new pelecypod genus from Russia, M. Plotnikov; New lamellibranchs from the Tatarian of Russia, M. Plotnikov (1); A symmetrical splitting in the (blade?) line in the shell in Triassic ammonites, T. N. Popov (1); New and known molluscs of Dzhungar depression, Russia, M. Saidov; Upper Trias and the Trachyceras zone in the Western United States, N. J. Silberling; New molluscs from the Okhotsk Sea area, Russia, I.

Tuchkov; Triassic cephalopods from Peca, Yugoslavia, L. Žlebnik.

Mesozoic.—Mesozoic of Belgian Congo, Beaucourt; Trigonia from sandstone of Cretaceous? of Hokkaido, Japan, S. Imanishi; Mesozoic mollusca in continental deposits of Trans-Baical, Russia, G. Martinson (1); Mollusca from Taimyr, Russia, L. Miroshnikov; Mesozoic molluscs from Kara-Tan, Russian Turkestan, E. Rammelzhmeier; Nonmarine fossil molluscan faunas in Eastern Asia with lists of Mesozoic species pp. 117–120, K. Suzuki; Mesozoic Psiloceras from central Switzerland, R. Trümpy; Mesozoic Mollusca of the S.E. Caucusus, Vassoevich.

Permian.—Upper Permian of Southern Rhodesia, L. R. Cox (1); Permian Pelecypods from Western Australia, J. M. Dickins; Western Australia, Permian Pelecypods, J. M. Dickins; Mollusca in Permian of Verkhoyan, Russia, Kulikov: Mollusca in the Donetz layer. Permian, Russia, Lapkin; New ammonites from the Yurezan river, S. Urals, Russia, S. Maksimora; Paleozoic ammonoids, their classification, Miller & Furnish; East Greenland, Permian Pelecypods, N. D. Newell; Anthracosia spp. from Roumania, G. Răileanu; List of Peruvian Mollusca in Asselsk stratum, Russia. Ruzhentzev: New Paleozoic Ammonoids, Ruzhentzev (1); Permian Ammonites, Ruzhentzev (2); New ammonites from the S. Urals, Russia, V. Ruzhentzev (3); New ammonites from the Sakmara deposit, Kazakh, Russia, V. Ruzhentzev (4); Mollusca of Kolvu-Vishersk Land, Russia, upper Paleozoic, Sultanaev.

Carboniferous.—New Imitoceras from Russia, E. Balashova; New Scottish Goniatites, E. Currie; Naming nonmarine lamellibranchs, R. Eager; Carboniferous of the Pennine District, R. M. C. Eager (1) Ammonites from Kara-Chatir, Ferghana, Turkestan, Miklukho-Maklai & Ruzhentzev; Mollusca in lower Carboniferous of Karagand basin, Monakhova; Lower Cretaceous Valanginian fossils of Grand Canon, Verdon, Mongin & Trouvé; Mollusca of the Ostrog suite of Kuzbass, Carboniferous, Muromtzev; Frankenburg Carboniferous fossils, E. Paproth; Northern England a review of Lower Carboniferous fossils, D. H.

Raquer; New and known ammonites from the Southern Urals, Russia, V. Ruzhentzev (3); Mollusca of Carboniferous in Galizia—Volhymian depression, Poland, Shulga.

Devonian. - New Devonian ammonoids, Bogoslovski; Devonian Goniatitacea, H. K. Erben; New ammonites from Germany, K. Müller; Devonian Mollusca of Kinovsk layers, Russia, Muravyev & Ivanov: Pseudoscolar pele; evpods from the Urals, B. Nalivkinz Devonian Mollusca in Dnieper-Donetdepression, Novik; Mnataia new Peleeypod genus from Bohemia, Prantl & Ružička: Devonian Mollusca in Pripet river deposits, Poland, Stefanenko & Makhnach; New Devonian molluses from Victoria, Australia, J. A. Talent; Siluro-Devonian molluses from Marble Creek, Victoria, Australia, Talent & Philin: Mollusca in mid-Devonian deposits of Tuva, Russia, Varentzov.

Silurian.—New Silurian Loricata from Sweden, J. Bergenhayn; Type and figured specimens from the Tortworth Inlier, Gloucestershire, M. L. Curtis; Silurian, new species of the Upper Gala-Tarannon horizon, A. Lamont; Siluro-Devonian molluses from Marble Creek, Thomson river, Victoria, Australia, Talent & Philip.

Ordovician.—New Loricata from the Ordovician of Scotland, J. Bergenhayn; Type and figured specimens from the Tortworth Inlier, Gloucestershire, M. L. Curtis; Spitzbergen Ordovician Cephalopods, H. Major; Tentaculites from Argentina, C. Rusconi; List of Ordovician molluses from Mendoza, Argentina, pp. 53, 54, C. Rusconi (2); Nautiloids from the Tanyard formation of central Texas, A. G. Unklesbay (1).

Cambrian.—Scenella and Stenothecoides, on internal shell structure, F. Rosetti; Two new gastropods from Mendoza, Argentina, C. Rusconi (1); List of Cambrian molluses from Mendoza, Argentina, pp. 22,-26, C. Rusconi (2); New mollusea for Cambrian of northern Tyan-Shan, Russia, Volozdin.

Palaeozoic.—Fossils of Ohio, U.S.A., La Rocque & Marple; Paleozoic molluscs of Tyan-Shan, Sinikin; Norway on stranded specimens of giant squids, E. Sivertsen.

### III.—SYSTEMATIC INDEX

N.B.—FULL REFERENCES CAN BE FOUND WHEN THEY ARE NOT GIVEN IN THIS INDEX BY REFERENCE TO 'TITLES' UNDER THE NAME OF THE AUTHOR GIVEN IN THE SYSTEMATIC INDEX AND THE NUMBER OF THAT AUTHOR'S PAPER WHERE MORE THAN ONE IS RECORDED. WHERE FULL REFERENCE IS GIVEN IN THE SYSTEMATIC INDEX THE VOLUME NUMBER IS GIVEN IN small clarendon.

Note.—The arrangement here used follows in general that of Thiele's 'Handbuch'. The most important differences are that Epitonacea and Pyramidellacea are placed immediately preceding Muricacea in order of grouping and the families Triviidae and Eratoidae are placed in Lamellariacea.

Where more than one page is given in a reference, the one in roman type is where the new name first occurs and the one in *italic* is where the gen. sp. or var. is actually described.

## Polyplacophora CHITONACEA

Amphineura study of the systematics C. Boettger (1).

Catalogue of species from Argentinian and adjacent waters, Z. Castellanos.

Description of spermiogenesis in some species, A. Franzén (1).

Acanthopleura (Maugeria) granulata description and first record from Cozumel Is., Quintana Roo, off east coast of Mexico, E. Hidalgo p. 4 pl. 3, 4.

Callistochiton (Lophochiton) carpenterianus nom. nov. for Callistochiton carpenteri Nierstrasz (Banda Indonesia), P. Kaas Basteria 20 4/5 Dec. 1956 p. 106.

Callochiton falvus (Wood) in the River Fal, Cornwall, R. H. Baird.

Chaetopleura papilio and C. apiculata Structure of girdle and sense organs, E. Leloup.

†Chelodes variegatus p. 13 pl. 1 f. 4; pl. 2 f. 3, Gotlandian, Silurian, Sweden; C. longissimus pl. 14 pl. 1 f. 5, Bala, Ordovician, Scotland spp. nov., J. Bergenhayn.

Chiton marmoratus description and first record from Cozumel Is., Quintana Roo, off east coast of Mexico, E. Hidalgo p. 2 pl. 1, 2.

†Haeggochiton gen. nov. p. 22 Scanochitonidae Genotype; Haeggochiton haeggi sp. nov. p. 22, pl. 1 f. 12; pl. 2 f. 11, Senon, Cretaceous, Sweden, J. Bergenhayn.

†Gotlandochiton gen. nov. p. 15 Gotlandochitonidae fam. nov. Genotype: Gotlandochiton interplicatus p. 15 pl. 1 f. 6; pl. 2 f. 4; G. laterodepressus p. 17 pl. 1 f. 8; pl. 2 f. 5; G. birhombivalis p. 18 pl. 1 f. 7; pl. 2 f. 6; G. troedssoni p. 19 pl. 1 f. 9; pl. 2 f. 7 spp. nov. Gotlandian, Silurian, Sweden, J. Bergenhayn.

†Gotlandochitonidae fam. nov. Type genus: Gotlandochiton gen. nov. Silurian, Sweden, J. Bergenhayn p. 15.

†Ivoechiton gen. nov. p. 22 Scanochitonidae. Genotype: I. levis sp. nov. p. 23 pl. 1 f. 16; pl. 2 f. 10, Senonian. Schonen, Ivö Island, Sweden, J. Bergenhayn.

Lioplax of North America, Clench & Turner (1).

†Llandeilochiton gen, nov. Llandeilochitonidae Genotype: L. ashbyi sp. nov. Llandeilo, Ordovician, southern Scotland, J. Bergenhayn p. 26 pl. 1 f. 14; pl. 2 f. 12.

†Llandeilochitonidae fam. nov. Type genus: Llandeilochiton gen. nov. J. Bergenhayn p. 26.

Paricoplax profundior sp. nov. 130 fathoms, off the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 157 f. 213-219.

†Septemchiton gen. nov. Septemchitonidae Genotype: Septemchiton vermiformis sp. nov. Scotland, Bala, Ordovician, J. Bergenhayn p. 24 pl. 1 f. 15; pl. 2 f. 13, 13a-b.

†Septemchitonidae fam. nov. Type genus: Septemchiton Ordovician, Scotland, J. Bergenhayn p. 24.

Septemehitonina new order, Class Loricata, J. Bergenhayn p. 24.

#### Gastropoda

Pelagic swimmers and drifters, Anon (1); Spanish Cretaceous Gastropoda, J. Bataller (1); Physiology of nutrition, W. v. Buddenbrock pp. 255-279; Key to common genera of planktonic Gastropoda p. 265, C. Davis; Ability of some gastropods to erode littoral rocks,

P. Fischer; Spermiogenesis in some species, A. Franzén (1); Observations on micro-secretion in numerous species, M. Gabe; Gastropoda secretions, M. Gabe (1); Adaptation of Freshwater species, W. D. R. Hunter; Means of penetration of host by parasitic Gastropoda, A. Iwanow; Lengthening reaction (Bingham Curve) of smooth muscles in some gastropods, N. Postma; Gastropoda from Strangford Lough and neighbouring coasts, Ireland, G. Williams.

#### STREPTONEURA

## Aspidobranchiata

## Rhipidoglossa BELLEROPHONTACEA

†Bellerophon sitiricus, B. Khayanlakhensis spp. nov. Verkhoyansk ridge, Russia, Permian p. 1103, Kulikov, M. V. C.R. Acad. Sci. Moscow 101 6 1055

†Tremanotus cyclocostatus sp. nov. Siluro-Devonian. Marble Creek, Thomson River, Victoria, Australia, Talent & Philip p. 61 pl. 6 f. 11, 15; pl. 7 f. 16, 17 text-f. 4.

†Worthenia kobelsbergia p. 102 pl. 1 f. 7a-c; W. goederti p. 103 pl. 1 f. 8a, b; W. schmidti p. 103 pl. 1 f. 9a, b spp. nov. Trochitenkalk, Triassic, Wiesloch, Germany, H. Eckert.

#### HALIOTIDACEA

†Emarginula didactica p. 8 pl. 2 f. 2; E. dilatoria p. 9 pl. 2 f. 3 sp. nov. Pliocene, Australia N. H. Ludbrook [nom nud. op. cit. 77 1954: p. 58.]; E. rochefortensis p. 323 f. 1 2; E. charmantensis p. 324 f. 3, 4; E. arnaudi p. 325 f. 7, 8; spp. nov. Upper Cretaceous S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954.

Entennotrochus rumphii Schepman a specimen from near Taiwan, but destroyed pl. 9 f. 3, 4, T. Kuroda (2).

Fissurellidae keys and list, Anon (5).

Haliotis—Notes on an ecological study, R. Talmadge; H. varia notes on the species, R. Talmadge (1).

†Kebinia gen. nov. Pleurotomariidae Type: K. pulchra sp. nov. p. 355 f. 2 River Karagaïlyu-Bulak, Chu-Iliisk mountains, Kirgizia, Russia. A. G. Vologdin. †Orthonychia marblecreekensis p. 65 pl. 7 f. 22; O. pentalvea p. 65 pl. 7 f. 28, 29 spp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip.

†Platyceras decorum p. 64, pl. 7 f. 21; P. trirotundolobatum p. 64, pl. 7 f. 26, 27 spp. nov. Siluro-Devonian, Marble Creek, Thomson river, Victoria, Australia, Talent & Philip.

†Platyostoma triangularis sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 64, pl. 6 f. 5, 6.

Pleurotomaria teramachii sp. nov. off Tosa south of Bungo Channel, Pacific side of Shikoku Island, Japan T. Kuroda (2) Venus 18 4 1955 p. 213 pl. 8 f. 1, 2.; P. africana Tomlin comparative table with P. teramachii sp. nov., T. Kuroda (2).

†Temnotropis palatina sp. nov. Trochitenkalk, Triassic, Wiesloch, Germany, H. Eckert p. 105 pl. 2 f. 17a, b, 18, 19.

## Prosobranchiata

#### Docoglossa

#### PROSOBRANCHIATA

Various types of growth in prosobranchs, P. Drach; New Caledonian prosobranchs, A. Franc (2); Roumanian Prosobranchia, A. V. Grossu (3); Illustrated handbook of Swedish forms, B. Hubendick (2).

#### PATELLACEA

Acmaea dorsursa colony formation, N. Abe; A. limatula adaptive differences in water-holding capacities of high and low level intertidal forms, E. Segal; A. limatula adaptations to microhabitat temperature, E. Segal (1).

Lottia gigantea Gray, Water circulation in the mantle cavity, figs., D. P. Abbott.

Maoricrater gen. nov. Lepetidae. Type: Notoacmea explorata Dell, 15-330 fathoms in New Zealand waters, R. K. Dell p. 57 p. 58 f. 249-252.

Patella hermaphroditism in three British species, J. M. Dodd; P. vulgata cytoplasmic inclusions of neurones, Lacy & Rogers; P. vulgata observations on breeding, Orton, Southward & Dodd; P. vulgata purification and properties of its steroid sulphatase, A. B. Roy;

P. vulgata population balance with seaweeds on wave-beaten shores, A. Southward; Patella note on work done by J. H. Orton, Southward & Dodd; P. coerulea behaviour of nucleoli and ribonucleo-protein in oocytes, E. Vannini.

Patellidae, observations on non-homing species, R. Ohgushi.

Scenella on the internal shell structure, F. Rasetti.

#### TROCHACEA

Alertalex gen. nov. Calliostomatidae. Type: A. blacki sp. nov. 220–280 fathoms, Chatham Rise, Chatham Islands, New Zealand, R. K. Dell p. 46 f. 61, 120, 260.

†Astraea (Bellastraea) hesperus sp. nov. Pliocene, Australia, N. H. Ludbrook p. 23 pl. 2 f. 8 [nom. nud. loc. cit. 77 1954: p. 58.]

Brookula benthicola sp. nov. 300 fathoms, off east Otago coast, New Zealand, R. K. Dell p. 49 f. 55.

Calliostoma subgeneric classification, J. L. Staadt.

Cantharidus kingensis sp. nov. south east of King Island, Victoria, Australia, C. J. Gabriel p. 5 f. 3.

Circulus striatus general anatomy, V. Fretter.

Cirsonella propelaxa sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 54 f. 54.

†Collonia omissa sp. nov. Pliocene, Australia, N. H. Ludbrook p. 23 pl. 2 f. 7 [nom. nud. op. cit. 77 1954: p. 58.]

Gibbula umbilicalis da Costa G. pennanti Phil. Ecological notes, Fischer-Piette & Gaillard; Gibbula distribution of species in part of the Gulf of Marseille, P. Huyé.

Liotella mackenae sp. nov. 130 fathoms east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 50 f. 72.

Lischkeia undosa sp. nov. deep sea off Tosa Bay, Japan, Kuroda & Kawamura Venus, 19 1 1956 p. 16 text-fig. 1 2.

Lissotesta otagoensis p. 52 f. 64, Portobello Alert Station 54-17, off east Otago coast, 260-350 fathoms, L. ambigua p. 52 f. 69, 120-125 fathoms, N.E. of Kaingaroa, Chatham Islands spp. nov. New Zealand, R. K. Dell. Micrelenchus caelatus archibenthicola subsp. nov. 300 fathoms, off east Otago coast, New Zealand, R. K. Dell p. 45 f. 16.

Monodonta distribution of species in a creek of the Gulf of Marseille, P. Huvé.

†Pellax jejuna sp. nov. Pliocene, Australia, N. H. Ludbrook p. 24 pl. 2 f. 9 [nom. nud. op. cit. 77 1954: p. 58.]

†Spectamen planicarinatum p. 17 pl. 2 f. 4; S. praecursor p. 18 pl. 2 f. 5 sp. nov. Pliocene, Australia, N. H. Ludbrook [nom. nud. op. cit. 77 1954 p. 58.]

†Suchium (Suchium) koyuense p. 49 pl. 4 f. 1 text-f. 1, Yamaji; S. (Protorotella) maxima p. 53 pl. 4 f. 7 text-f. 3, Kakoi; S. (P.) hyuguensis p. 54 pl. 1 f. 3, 4 text-f. 4, 5, Yamaji spp. nov. Miyazaki Group, Pliocene-Miocene, Koyu Gun, Miyazaki Prefecture, Japan, T. Shuto; S. (Suchium) giganteum evolutional lineage, T. Shuto p. 61.

†*Teinostoma yabei* sp. nov. Miocene Noto Peninsula, Japan, K. Masuda Tr. Proc. Palaeont. Soc., Japan N.S. No. 20 1955 p. 123 pl. 19 f. 16-18.

Thoristella chathamensis profunda subsp. nov. 260-350 fathoms, off E. Otago coast, New Zealand, R. K. Dell p. 44 f. 57.

Trochidae Keys and lists, Anon (6). Umboniinae, notes on Phylogeny, K. Oyama (2).

Zetela variabilis sp. nov. 155 fathoms, S.E. of Pitt Island, Chatham Islands, New Zealand, R. K. Dell p. 48 f. 70,74.

#### NERITACEA

†Hologyra frentzeni sp. nov. Trochitenkalk, Triassic, Wiesloch, Germany, H. Eckert p. 107 pl. 2 f. 22a, b, 23, a, b, 24.

†Naticella spinosa sp. nov. Trochitenkalk, Triassic, Wiesloch, Germany, H. Eckert p. 109 pl. 3 f. 28a-d, 29, 30.

Neritidae, biology of middle European species, E. Frömming (4) p. 273-280.

Neritidae from Java, E. v. B. Jutting. Neritina Lamarck, 1816, gender feminine, Direction 42.

Theodoxia numidica Rech. vars major, minor, subcarinata vars. nov. ex MSS (in coll.) N.W. Africa, F. Llabador J. de Conchyliol. 96 3 Nov. 1956 pp. 87, 96-97.

#### CYCLOPHORACEA

Alycaeus, notes on various species and figures of radulae and jaws, L. A. W. C. Venmans.

Ampellarius and Pomacea, notes on the generic names, T. Pain (1).

†Ampullela peruviana sp. nov. Jurassic Nasca area Peru, L. R. Cox (4) J. Pal. 30 5 1956 p. 1181 pl. 127 f. 2, 3.

Bellamya from Java, E. v. B. Jutting.

Campeloma of North America, Clench & Turner (1).

Cyclophoridae 'type' specimens in the Senckenberg Museum, A. Zilch; Types in the Senckenberg Museum (4) figs., A. Zilch (2) Arch. Mollusk. Frankfurt a.M. 85 4/6 1956 pls. 12, 13.

Mexycyclotus petersi petersi sp. nov. p. 56 pl. 5 f. 9; pl. 6 f. 14-17, La Placita, Michoacan; M. petersi damianensis subsp. nov. p. 57 pl. 5 f. 11, pl. 6 f. 13, 18. North of San Pedro Damian Narajestilla, Michoacan, Mexico, A. Solem.

†Paludina proserpinae—P. mangikiani — P. turrita — P. dresseli hypertrophica — P. lungershauseni — P. elatior spp. nov. Lower Kama river basin, Quaternary, Russia, G. Goretzki p. 33; P. (= Viviparus) vivipara minor f. nov. Holocene, Ražice, Cidlina river, Bohemia, J. Petrbok (1) p. 8.

Pila virens composition and function of spherioles in the digestive gland, V. Meenakshi; P. virens seasonal variations in glycogen and fat content, V. Meenakshi (1); P. virens physiology of hibernation, V. Meenakshi (2); P. virens caratenoid, V. Meenakshi (3); P. proteins obtained from free aminoacids in the food, V. Meenakshi (4); P. Javanese species, E. v. B. Jutting; P. globosa exeretory constituents of the blood, B. Saxena; P. globosa nutritive constituents of the blood, B. Saxena (1); P. globosa ecology and behaviour, B. Saxena (2); P. africana koleensis subsp. nov. river basin of the Aruwimi near Kole, Belgian Congo, L. W. A. C. Venmans (1) J. Conch. Lond. 24 4 July, 1956 p. 141 and plate

† Pila ovata from the Quarternary, Tel Aviv, Palestine, N. Avnimelech.

Pomacea canaliculata, general biology, variation in shell size, genitalia, H. d. S. Lopes; P. lineata general anatomy,

H. de S. Lopes (1); P. cumingi urabaensis subsp. nov. Golpo de Uraba, northern Antioquia, Colombia, T. Pain J. Conch. 24 3 Feb. 1956 p. 75 text-figs.; P. and Ampullarius, notes on the generic names, T. Pain (1); P. cumingi (King) found at Nueva on the Rio Truando, dept. Choco, north western Colombia, H. A. Pilsbry (2); Pomacea carotenoids of some Brazilian species, G. Villela.

Valvata significance of Garnault's duct in the phylogeny of the genital system, J. Johansson; V. tricarinata (Say) variation of carinae, A. La Roque; V. piscinalis biology, M. Onigny.

†Valvata antiqua gigas "sp. nov." [?var. nov.] Quaternary, lower Kama basin, Russia, G. Goretzki p. 33; V. (Atropidima) wenzi sp. nov. Vienna, A. Papp Mitt. Geol. Ges. Wien. 44 1956 p. 110 pl. 4 f. 4, 5.

Valvatidae, biology of middle European species, E. Frömming (4) p. 268–272.

Viviparidae from the rivers of Alabama, Georgia and Florida, U.S.A., Clench & Turner p. 109; Viviparidae, biology of middle European species, E. Frömming (4) p. 247–267; Viviparidae in the Senckenberg Museum. Types, species and varieties, figs., A. Zilch (1).

Viviparus species of Europe and their distribution, C. R. Boettger (3). V. viviparus, radula, W. v. Buddenbrock p. 258 f. 98.

†Viviparus kinelicus sp. nov. Lower Kama basin Quarternary, Russia, Goretzki, G. I. p. 34; V. brevis sp. nov. Triassic, Dzhungen depression Russia, Saidov, M. N. p. 94; V. sinensis sp. nov. Triassic, Dzhungen depression, Russia, Saidov, M. N. p. 95; V. praeglacialis Don river village Krasni Kut, Russia Quarternary, Popov, G. I. (1) p. 34.

### LITTORINACEA

Littorina littorea general biology, H. A. Cole in M. Graham p. 177-178; L. obtusata—preference for Fucaceae for food, A. Dongen; L. littorea shape and form of trails made in different sediments, I. E. Grät; L. brevicula local variation, S. Hayasaka.

†Littorinopsis deserti sp. nov. Cretaceous, Moghara Massif, Sinai Egypt, I. G. Mahmoud p. 141 pl. 16 f. 2, 3.

Nodilittorina granularis (Gray) fig. Breeding, T. Habe (4); N. picta (Phil.); N. granularis Gray; N. pyramidalis Q. & G. egg-capsules, fig., T. Habe (8); N. pyramidalis Q. and G. systematic position; figs. radula, T. Habe (9); N. granularis extruding response, Ohsawa & Tsukuda.

Peasiella roepstorffiana Nevill eggcapsules, figs., T. Habe (8).

Pomatias elegans feeding habits, E. Frömming (5).

#### RISSOACEA

Alvania reticulata, A. subsoluta reproductive system, J. Johansson.

†Alvania montaqui trochiformis subsp. nov. Miocene. Hungary, I. Csepreghy-Meznerics p. 443-444 pl. 2 f. 7-10.

†Amphithalamus (Pisinna) subbicolor sp. nov. Pliocene, Australia, N. H. Ludbrook p. 27 pl. 2 f. 10.

Apataxia gen. nov. p. 401 Rissoinidae, genotype: A. erecta p. 401 f. 35, Queensland coast, Australia; A. eripona p. 402 f. 36, Port Moresby, New Guinea spp. nov. C. F. Laseron.

Assimineidae, biology of middle European species, E. Frömming (4) p. 244–246.

Australaba gen. nov. p. 459 Rissoidae genotype A. bowensis p. 459 f. 195, Bowen, Queensland; A. difformis p. 460 f. 196-198, Darwin, Northern Territory spp. nov. Australia, C. F. Laseron.

Austrosina gen. nov. p. 393 Rissoinidae genotype: Rissoina pulchella Brazier, 1877; Austrosina quinia p. 394 f. 14; A. evanida p. 394 f. 15 spp. nov. Christmas Island, Indian Ocean, C. F. Laseron.

†Baicalia obrutschewi; B. carinataides spp. nov. Triassic, Dzhungar depression, Russia, M. N. Saidov p. 96.

Bythinella austriaca from mountain springs in Hungary, note on ecology, Farkas & Vagvolgyi; B. (Fraunefeldtia) intermedia nov. sp. nom. nud. in list p. 22. Austria, F. Mahler.

†Bithinia tentaculata minor f. nov. Holocene, Kostomlaty on the river Elbe, Bohemia, J. Petrbok (1) p. 8.

Bithynia Leach, 1818, objection to proposed validation, C. R. Boettger (2);

B. tentaculata embryology, O. Hess; and O. Hess (1 and 2); Bithynia leachi distribution in the lakes of the Krasnoyarsk Territory USSR, N. Khristenko; B. tentaculata protozoan parasites of specimens from the Firth of Vistula, Poland, Z. Raabe.

†Bithynia cumulata sp. nov. Triassic, Dzhunger depression. Russia, Saidov, M. N. p. 94; B. paluchella sp. nov. Triassic, Dzhunger depression, Russia, M. N. Saidov p. 96 (description begins as last word on p. 95.)

Bithyniinae, biology of middle European species, E. Frömming (4) p. 230-243.

†Chartroniella saxorum sp. nov. Lower Cretaceous Trinidad, L. R. Cox (2) J. Paleont. 28 5 1954 pp. 624 625 pl. 67 f. 2a b, 3a b.

Chrystella gen. nov. Rissoinidae. Type: C. islandica sp. nov. Christmas Island, Indian Ocean, C. F. Laseron p. 247 f. 111.

Cochliopa lubricella distribution in Hungary, J. Vágvölgyi.

Condylicia gen. nov. Rissoinidae. genotype: C. collaxis sp. nov. Howick Group, Queensland, Australia, C. F. Laseron p. 418 f. 80, 81.

Contraxiala gen. nov. Rissoinidae. genotype: C. obliqua sp. nov. Michaelmas Cay, Queensland, Australia, C. F. Laseron p. 421 f. 88.

Costabieta gen. nov. Rissoinidae. genotype: C. paucina sp. nov. p. 421 f. 87. Michaelmas Cay, Queensland, Australia, C. F. Laseron.

Costalynia gen. nov. p. 394 Rissoinidae, genotype: Rissoina cardinalis Brazier; Costalynia torresiana p. 395, f. 18; C. bilinea p. 397 f. 25, 26, Murray Island; C. decapitata p. 396 f. 20 Michaelmas Cay; C. dux p. 397, f. 22, Green Island; C. birestes p. 397 f. 23, 24 Torres Strait, Queensland; C. truncata p. 396, f. 21 Darwin, Northern Territory, Australia; C. isolata p. 396 f. 19 Christmas Island, Indian Ocean spp. nov., C. F. Laseron.

Cylonidea gen. nov. Rissoidae genotype: C. carina sp. nov. Christmas Island, Indian Ocean, C. F. Laseron p. 453 f. 181.

Dardanula difficilis sp. nov. south east of King Island, Victoria, Australia, C. J. Gabriel p. 7, f. 5.

Dentrissoina gen. nov. p. 415, Rissoinidae, genotype: D. thornleyana sp. nov. p. 415, f. 70, 71 Murray Island, Queensland, Australia, C. F. Laseron.

Dipsotoma gen. nov. p. 416, Rissoinidae, genotype: Rissoa mercurialis Watson, 1886 Wednesday Island, Queensland, Australia, C. F. Laseron.

Diala conspicua p. 455 f. 183; D. fossa p. 455 f. 184 Michaelmas Cay; D. premeris p. 456 f. 185, Thursday Island; D. planalba p. 456 f. 187; D. diffilata p. 457 f. 190, 191 Bowen; D. cacumenata p. 457 f. 188; D. vestigia p. 457 f. 189 Hope Island; D. cento p. 458 f. 192 Keppel Bay; D. tumerosa p. 458 f. 193 20 fm. off Endeavour Reef spp. nov. Queensland, Australia; D. watsoni nom nov. for (?) Barleeia imbricata Watson, 1886, non Diala imbricata A. Adams, 1862, C. F. Laseron.

Discrevinia gen. nov. Rissoidae, genotype: D. balba sp. nov. Christmas Island, Indian Ocean, C. F. Laseron p. 437 f. 140.

Estea rufoapicata latior subsp. nov. p. 63 f. 82; E. sculpturata sp. nov. p. 63 f. 81 300 fathoms off E. Otago coast, New Zealand, R. K. Dell; E. hoggartae sp. nov. south east of King Island, Victoria, Australia, C. J. Gabriel p. 6 f. 4; E. lunata p. 441 f. 147, Bowen; E. perdigna p. 441 f. 148 Mast Head Is., Capricorn Group, Queensland, Australia spp. nov. Also included in this genus, Eulima eurychades Watson and temporarily Mucronalia xanthias Watson, C. F. Laseron.

Eprigrus obesa p. 448 f. 167; E. truncata p. 448 f. 168 spp. nov. Darwin, Northern Territory, Australia, C. F. Laseron.

Fractoralla gen. nov. p. 398 Rissoinidae, genotype: F. praecida p. 398 f. 27; F. transita p. 399 f. 28 Darwin, Northern Territory; F. decipiens p. 399 f. 29 Port Curtis; F. heronensis p. 399 f. 30, 31 Heron Island, Capricorn Group, Queensland, Australia spp. nov., C. F. Laseron.

Haurakia subsuturalis p. 61 f. 73; H. otagoensis p. 62 f. 76 spp. nov. 300 fathoms, off E. Otago coast, New Zealand, R. K. Dell; H. firma p. 438 f. 142, Keppel Bay; H. formosita p. 438 f. 143 Michaelmas Cay, Queensland, Australia; H. isolata p. 439 f. 144 Christmas Island, Indian Ocean spp. nov., C. F. Laseron. Hydrobia (Hydrobia) testulata p. 113 pl. 7 f. 2-4; H. (Odontohydrobia) cobessdonfensis p. 114 pl. 8 f. 15-18 spp. nom., A. Papp Mitt. Geol. Ges. Wien 44 1953.

†Hydrobia gonduranica sp. nov. Karros Bed and (Upper Permian) Southern Rhodesia, L. R. Cox (1) Geol Mag. 90 3 1953 p. 204 text-fig. 1 a-f.

Hydrobiidae, biology of middle European species, E. Frömming (4) p. 206–224.

Isselia polytropa augerea subsp. nov. p. 411 f. 60; I. lateralis p. 412 f. 61; I, undulata p. 412 f. 62, Michaelmas Cay. Queensland, Australia; I. incisa p. 412 f. 63 Christmas Island, Indian Ocean spp. nov., C. F. Laseron.

Lithoglyphinae, biology of middle European species, E. Frömming (4) p. 225–229.

Lucidestea gen. nov. p. 448 Rissoidae genotype: L. vitrea p. 449 f. 169; L. obesa p. 449, f. 171, Michaelmas Cay; L. maculosa p. 450 f. 172, Lindeman Island; L. laterea p. 450 f. 173, Bowen; L. sublacuna p. 451 f. 175 Mast Head Is., Capricorn Group, Queensland; L. perforata p. 449 f. 170, Darwin, Northern Territory spp. nov. Australia, C. F. Laseron.

Lucidinella gen. nov. p. 427 Rissoinidae. genotype: L. conicera p. 427 f. 112, 113, Whitehaven Bay; L. sublevis p. 428 f. 116 Curtis Island, Queensland spp. nov.; L. conicera patruelis p. 428 f. 114, 115 Darwin, Northern Territory subsp. nov. Australia, C. F. Laseron.

Mellitestea gen. nov. p. 452 Rissoidae genotype: M. simplex sp. nov. 5-10 fm. Murray Island, Queensland, Australia, C. F. Laseron p. 453 f. 180.

Merelina queenslandica p. 433 f. 130, Barrier Reef, off Cairns; M. solida p. 434 f. 131, 14 fm., Albany Passage; M. reversa p. 435 f. 134, Bowen; M. corruga p. 436 f. 136, Heron Island, Capricorn Group; M. goliath p. 436 f. 137, 138, Hervey Bay, Queensland; M. sucina p. 434 f. 132, 133; M. humera p. 435, f. 135, Darwin, Northern Territory, Australia; M. hians p. 436, f. 139, Christmas Island, Indian Ocean spp. nov.. C. F. Laseron.

†Merelina (Linemera) varisculpta sp. nov. Pliocene, Australia, N. H. Ludbrook p. 28 pl. 2 f. 11 [nom. nud. op. cit. 77 1954 p. 59.]

Mesodestea gen. nov. Rissoidae genotype M. resima sp. nov. Darwin, Northern Territory, Australia, C. F. Laseron p. 451, f. 176.

Micromelania (Micromelania) goberoderfensis f. 121 pl. 5 f. 5-8; M. (Goniochilus) breviformis p. 123 pl. 6 f. 8 spp. nov.; M. (Goniochilus) banatica fuchsi subsp. nov. p. 122 pl. 6 f. 9, A. Papp Mitt. geol. Ges. Wien 44 1956.

Moerchiella persista p. 414 f. 67, Thursday Island; M. deleta p. 414 f. 68, Port Curtis; M. volvera p. 415 f. 69, Annan River spp. nov. Queensland, Australia, C. F. Laseron.

Nanadoma gen. nov. p. 447 Rissoidae genotype: N. imitoris sp. nov. Darwin, Northern Territory, Australia, C. F. Laseron p. 447, f. 165.

Nobolira orientalis p. 64 f. 75, 130 fathoms, east of the Forty Fours, Chatham Islands; N. merelina p. 65 f. 80, 300 fathoms, off east Otago coast spp. nov. New Zealand, R. K. Dell.

Obtortio columna p. 463 f. 211–214; O. duplex p. 466 f. 226, Darwin, Northern Territory; O. varicosa p. 464 f. 220; O. ambiqua p. 465 f. 221; O. fabrica p. 465 f. 222–224, Bowen; O. crassa p. 466 f. 225, Michaelmas Cay, Queensland, Australia; O. rugosa p. 464 f. 218, 219, Christmas Island, Indian Ocean spp. nov.; O. fulva axialis p. 462 f. 204, 205, Darwin; O perparvula progreda p. 464 f. 217, Darwin and Murray Island subspp. nov. Australia, C. F. Laseron.

Oncomelania nosophora (Robson) Anatomy studies and radular formula, H. Itagarki; O. quadrasi occurrence in the Philippines, B. Hubendick (1); Oncomelania effects of ultraviolet radiation, Wong & Wagner.

†*Oonia trinidadensis* **sp. nov.** Lower Cretaceous Trinidad, **L. R. Cox** (2) J. Paleont **28** 5 1954 pp. 624 626 pl. 65 f. 3, 6.

Lartetia rhenana localities in Germany, F. Kiefer.

Linemera solitaria p. 66 f. 77, 130 fathoms, east of the Forty Fours, Chatham Islands; L. abrupta p. 66 f. 78, off east Otago, in 300 fathoms spp. nov. New Zealand, R. K. Dell.

Palisadia gen. nov. p. 431 Rissoinidae genotype: P. subulata sp. nov. Christmas Island, Indian Ocean, C. F. Laseron p. 432 f. 129. Paludestrina note on nomenclature and type species, J. Parodiz (5).

Pandalosia gen. nov. p. 391, Rissoinidae genotype: P. excelsis p. 391, f. 5, Murray Island, Queensland; P. darwinensis p. 391 f. 6, Darwin, Northern Territory; P. subulata p. 392 f. 7, Heron Island, Capricorn Group, Queensland; P. oceanica p. 392, f. 8, 9; P. delicatula p. 393 f. 12 Christmas Island, Indian Ocean; P. obtusa p. 392 f. 10, 11, Michaelmas Cay, Queensland spp. nov. Australia, C. F. Laseron.

Paradiala gen. nov. Rissoidae genotype: P. supraflecta sp. nov. 17-20 fm., Mast Head Is., Capricorn Group, Queensland, Australia, C. F. Laseron p. 459 f. 194.

Parashiela gen. nov. p. 439 Rissoidae genotype: P. ambulata sp. nov. Michaelmas Cay, Queensland, Australia, C. F. Laseron p. 439 f. 145.

Pellamora reflecta p. 423 f. 94; P. angulata p. 424 f. 95, 96; P. diminuta p. 424 f. 97, 98; P. truncata p. 425 f. 100, 101, Darwin, Northern Territory; P. capitata p. 425 f. 106, 107, Albany Passage; P. spiralis p. 426 f. 108, 109; P. amplexa p. 426 f. 110; Michaelmas Cay, Queensland, Australia; P. minatura p. 424 f. 99, Port Moresby, New Guinea spp. nov., C. F. Laseron.

Peripetella gen. nov. p. 400 Rissoinidae genotype: P. queenslandica p. 400 f. 32, Bowen; P. immersa p. 401 f. 33, Rocky Is., Cape Flattery; P. linearis p. 401 f. 34, Heron Is., Capricorn Group spp. nov. Queensland, Australia, C. F. Laseron.

Phintorene allanae proxima p. 405 f. 45 subsp. nov. Darwin; P. apicina p. 406 f. 46, 47, Facing Is. Port Curtis; P. ultima p. 406 f. 48 Albany Passage; P. caelata p. 406 f. 49, Cape Sidmouth; P. angusta p. 407 f. 50; P. digera p. 408 f. 52, Michaelmas Cay, Queensland; P. emina p. 407 f. 51; P. paenula p. 408 f. 53, Darwin, Northern Territory spp. nov. Australia, C. F. Laseron.

Phosinella cyatha sp. nov. p. 404 f. 41-43 Albany Passage, Queensland, Australia, C. F. Laseron.

Planapexia gen. nov. p. 408 Rissoinidae genotype: P. fractura p. 409 f. 54; P. quadrina p. 410 f. 58; P. elevata p. 410 f. 59, Darwin, Northern Territory; P. decepta p. 409 f. 55, Curtis Island, Queensland spp. nov. Australia, C. F. Laseron.

Pleneconea gen. nov. p. 402, Rissoinidae, genotype: P. angulata p. 402 f. 37 sp. nov. Lindeman Is., Queensland, Australia, C. F. Laseron.

Polamopyrgus jenkinsi, E. A. Smith and its distribution in Lake Windermere, H. P. Moon.; P. jenkinsi from Lake Razelm, Dobroudja, first record from Roumania, A. V. Grossu.

Pomatiopsis cincinnatiensis detailed morphology, Schalie & Dundee.

†Probaicalia gen. nov. p. 79, Micromelaniidae genotype: Probaicalia gerassimovi (Reis, 1910); P. hydrobioides p. 79 pl. 9 f. 4; P. vitimensis p. 80 pl. 9 f. 2; P. prinadae p. 80 pl. 9 f. 3 spp. nov.; P. hydrobioides rammelmeyeri p. 79 pl. 9 f. 6; P. hydrobioides elongata p. 79 pl. 9 f. 5 vars. nov.; P. gerassimovi; (Reis), comb. nov. p. 80 pl. 9 f. 1; near Lake Baikal, East Siberia, Cretaceous, G. Martinson.

†Prororthemia sinensis sp. nov. Triassic, Dzhunger depression, Russia, Saidov, M. N. p. 94.

Pyramidelloides pacifica p. 419 f. 82, Darwin, Northern Territory; P. turris p. 419 f. 83, Albany Passage; P. cylindrica p. 420 f. 84, Michaelmas Cay, Queensland, Australia; P. viticula p. 420 f. 85, Christmas Island, Indian Ocean spp. nov. C. F. Laseron.

Pyrgohydrobia st. naumi p. 25 f. 1-5; P. jablanicensis p. 26 f. 6-10 spp. nov. Ohrid, Jugoslavia, P. Radoman.

Rissoa revision of some of the species, **P. Mars** (2).

Rissoina notes on New Zealand species, A. K. Hipkins; R. polytropa Hedley, 1899, is now removed to Isselia Schmeltz, C. F. Laseron p. 411.

†Rissoina tinela sp. nov. Pliocene, Australia, N. H. Ludbrook p. 31 pl. 2 f. 19 [nom. nud. op. cit. 77 1954 p. 59.]

Saltratricula nom. nov., p. 445, for Saltatrix Laseron, 1950. Genotype: Epigrus protractus Hedley; S. ballerina p. 446 f. 162, Whitehaven Bay; S. stringera p. 446 f. 163; S. improrsa p. 446 f. 164 Noosa Head spp. nov. Queensland, Australia, C. F. Laseron.

Schwartziella lata sp. nov. Christmas Island, Indian Ocean, C. F. Laseron p. 390 f. 4.

Scrobs tropica p. 442 f. 151; S. incipiens p. 443 f. 153, Heron Is., Capricorn Group; S. compressa p. 442 f. 152, Michaelmas Cay; S. fulcira p. 443 f. 154, 155, Lindeman Island; S. oblata p. 444 f. 156, Noosa Head spp. nov. Queensland, Australia, C. F. Laseron.

Stenothyra from Java, E. v. B Jutting.

Synceridae from Java, E. v. B. Jutting.

†Turboella praenovarensis p. 28 pl. 2 f. 14; T. elimattae p. 29 pl. 2 f. 15 sp. nov. Pliocene, Australia, N. H. Ludbrook [nom. nud. loc. cit. 77 1954 p. 59.]

Vitricithna gen. nov. Rissoidae. Type: Cithia marmorata Hedley; V. relativa p. 452, f. 179 sp. nov. Michaelmas Cay, Queensland, Australia, C. F. Laseron.

Zebina heronensis p. 429 f. 117, Heron Is., Capricorn Group; Z. nitens p. 429 f. 119, 120, Michaelmas Cay, Queensland, Australia; Z. hebes p. 430 f. 121, Port Moresby, New Guinea; Z. isolata p. 430 f. 122; Z. acicula p. 430 f. 123, 124; Z. linearis p. 430 f. 125; Z. constricta p. 431 f. 126 Christmas Island, Indian Ocean; Z. inflata p. 431 f. 127, 128, Rocky Is., Cape Flattery, Queensland, Australia spp. nov., C. F. Laseron.

Zymalata gen. nov. p. 416, Rissoinidae. Genotype: Z. concinna p. 417, f. 75 sp. nov. Michaelmas Cay, Queensland, Australia. Genus also includes Rissoina triangularis Watson and R. inermis Brazier, C. F. Laseron.

### CERITHIACEA

Amnipila gen. nov. Potadominae genotype: A. pila (= Pachycheilus pila Pilsbry & Hinkley, 1910). San Luis Potosi, Mexico, H. A. Pilsbry (3) p. 38 Pl. 4 f. 7, 8, 9 text-f. 4.

†Ampullina chalaguarensis sp. nov. Maestrichtian La Chalaguire nr. Fabas S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 335 figs. 22.

Aylacostoma chlorotica p. 45 f. 1; A. stigmatica p. 47 f. 2 spp. nov. upper Paraná river, Paraguay, M. I. H. Scott. †Campanilopsis gen. nov. for Nerinea margaritifera d' Arch. P.-H. Fischer (3) J. de Conchyliol. Paris 96 1 June 1956 p. 3 and p. 10.

Cerithiacea, on the open Pallial oviduets, figs., J. Johansson.

Cerithidea djadjariensis C. rhizophorarum, spawning, T. Habe (3).

†Cerithinopsis arnaudi sp. nov. Turonian, S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 367 f. 61.

Cerithiolum reticulatum food and feeding habits in Black Sea forms Gajevskaja.

Cerithium vulgatum anatomy of genital organs, Johansson, J.; Cerithium occurrence near Olomouc, Czechoslovakia, V. Kalabis.

†Cerithium (Ptychocerithium) procrenatum from the Miocene of Slovakia, J. Svagrovský; C. livernantensis p. 359 f. 50 Campanian; C. (Pyrazus) arnaudi Cenomanian p. 360 f. 51; G. (P.) shoelleri p. 360 f. 52 Cenomanian; C. (P.) belestensis Maestrichtian p. 364 f. 58; C. (P.) chalaguerensis; p. 365 f. 59 Maestrichtian; spp. nov. Upper Cretaceous S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954; C. alulovići sp. nov. Senonian Serbia, O. Pejović (1) Zborn Redova geol. Inst. Beograd 5 1953 p. 81 pl. 1 f. 9.

†Cirsocerithium manzourianum sp. nov. Cretaceous. Moghara Massif, Sinai Egypt, I. G. Mahmoud p. 145 pl. 16 f. 12.

Cleopatra hemmingi sp. nov. Italian Somaliland, 18 miles west of Beles Cogani, in large shallow pools, B. Verdcourt (see Hemming & Verdcourt) p. 60 f. 3, 4c, e.

†Cossmannea nascaensis p. 1181 pl. 127 f. 4-6; C. peruviana p. 1182 pl. 127 f. 7-8; spp. nov. Jurassic Nasca area Peru, L. R. Cox (4) J. Pal. 30 5 1956.

Doryssa geijskesi sp. nov. Makambo Island, Coppename River, Surman, T. Pain (3) Basteria 20 4/5 Dec. 1956 p. 99 [nom et fig.] p. 102.

Melanella intermedia anatomy of genital organs, J. Johansson.

Melania cancellata Say from India not N. America, W. J. Clench (2). †Melania escheri from the Pliocene of the Salaj basin, Roumania, M. Paucă; M. ariegensis sp. nov. Maestrichtian Bilesta S.W. France, G. Termic Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 338 f. 26.

Melanoides tuberculata and Chaetogaster, F. A. H. Bayer; M. tuberculata (Müll) digestion and production of H<sub>2</sub>SO<sub>4</sub>, C. R. Fish; M. tuberculatus and Allopolyploidy, J. Jacob; M. tuberculatus parasitic effects in the digestive gland, Lal & Premyati.

Melanopsis dufouri morphology and biology, E. Frömming (1); M. martiniana occurrence near Olomoue, Czechoslovakia, V. Kalabis.

†Melanopsis impressa pseudonarzolina p. 132 pl. 9, f. 14-18; M. impressa posterior p. 133 pl. 9 f. 19-23; M. fossilis pseudoimpressa p. 135 pl. 11 r. 5-8 subspp. nov. M. tortispina p. 147 pl. 12 f. 18-20; M. scalariformis p. 15 3 pl. 3 f. 27-28; spp. nov., A. Tapp Mitt. Geol. Ges. Wien 44 1953.

Mendax nucleoproducta sp. nov. east of the Forty Fours, Chatham Islands, New Zealand, in 130 fathoms, R. K. Dell p. 68 f. 105.

†Mesalia astrei sp. nov. Maestrichtian Bélesta S.W. France, G. Termier Bull. Soc. Hist nat. Toulouse 39 3-4 1954 p. 346 f. 35.

†Metacerithium (?) tuberculatum sp. nov. p. 144 pl. 16 f. 11 text-f. 76, Cretaceous Moghara Massif, Sinai, Egypt, I. G. Mahmoud.

†Nerinea mandhourensis p. 26 pl. 2 f. 1 Gebel Mandhour; N. mistanensis p. 28 pl. 2 f. 7; N. pre-olissopomensis p. 31 pl. 1 f. 9, El-Mistan, Albian; N. hallalensis p. 29 pl. 1 f. 4, Gebel Hallal; N. dentiloba p. 27 pl. 1 f. 8; N. sinaitica p. 32 pl. 2 f. 3, Cenomonian, Spp. nov. Cretaceous, Sinai, Egypt, G. H. Awad; N. margaritifera d'Arch. the genotype of Campanilopsis [q.v.], P.-H. Fischer (3); †Nerinea cf. byzacenica Paving; N. zumoffeni Delpey; N. coquandi Orbizug; N. cf. salignaci Coquand; N. aptiensis Pict. & Cacufs; N. prefteuriani Delpey, of the Lower Cretaceous N. Africa, G. &. J. Glaçon; N. vagdti (La Moctillet) (emend) sousepice sublimensis sub sp. nov., Mongian Trouvé, Bull Soc. geol. Fr. 6 3 1-3 1953 p. 230 pl. iii f. 2; N. verdonensis

p. 227, pl. iii f. 1; N. granulifera p. 229 pl. iii f. 4 and text fig. 2a spp. nov. Valanginian Grand Casion du Verdon, Mougin & Trouvé Bull. Soc. geol. Fr. 6 3 1/3 1953.

†Nerinella lahfanensis sp. nov. Aptian, Cretaceous. Bir Lahfan, Sinai, Egypt, G. H. Awad p. 24 pl. 1 f. 5; N. caballensis sp. nov. Jurassic Nasca area Peru, L. R. Cox (4) J. Pal. 30 5 1956 p. 1182 pl. 127 f. 9-12.

Pachychilus pasionensis sp. nov. Peten and Alta Vera Paz, Guatemala, H. A. Pilsbry (3) p. 34 pl. 4 f. 4, 5.

Paramendax subapicina sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 69 f. 106.

†Parastrophia garganica sp. nov. Quaternary. Punta delle Pietre Nere, Foggia, Italy, M. Moncharmont Zei p. 120 f. 1-13.

†Planaxis simplex sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 148 pl. 16 f. 23-25.

Plotia Roeding, 1798, suppressed under the Plenary Powers, Opinion 386.

†Potamides tricarinatus occurrence at Saint Owen à Herblay, L. Feugueur; P. (Pirenella) plicatus and P. (Tympanotonus) margaritaceus from the upper Oligocene of Hungary, L. Strausz; P. (Tympanotonus) margaritaceus Br. and P. (Pirenella) plicatus Being very common in the Upper Oligocene of Hungary, figs., L. Strausz (1); P. reticosum maurensensis var. nov. p. 358 f. 47 Maestrichtian; P. occitanicum sp. nov. p. 358 f. 49 Maestrichtian. Auzas S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954.

Potamididae from Java, E. v. B. Jutting.

†Pyrazus aegyptiacus sp. nov. p. 145, pl. 16 f. 13, 14; P. sexangulatus Zekeli, "prémutation costata nov. premut." p. 146 pl. 16 f. 15; Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud.

†Stomatopsis? arnaudi sp. nov. Cenomanian Rochefort, S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 337 f. 25.

Thiaridae from Java, E. v. B. Jutting.

Torquesia Douvillé 1929 proposed addition to the "Official List of Generic

Names in Zoology" and proposed designation of Turritella granulata Sowerby (J. de C.) 1827, as the type species, J. Marwick.

Triphora perversa anatomy of genital organs, J. Johansson; T. medinae sp. nov. Punta Carretas, Montevideo, Uruguay, J. Parodiz (4) p. 59 fig.

†Turritella arnaudi sp. nov. Maestrichtian Bramfort, S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse \$9.3-4.1954 p. 346 f. 34. T. gradata sp. nov. Eocene? Ferghana, Russia, A. K. Alexeiev p. 30 pl. 5 f.4-10; T. (Haustator) slanskyi sp. nov. Lower Eocene Dahomey Ph. Brébion Bull. Mus. Hist. nat. Paris (2) 27 5 1955 p. 414; T. donkantensis Paleocene Dahomey, Ph. Brébion (1) Bull. Mus. Hist. nat. Paris (2) 27 3 1955 p. 244 figs.; T. incisaeformis sp. nov. Miocene, Hungary, I. Csepreghy- Meznerics p. 444 pl. 2 f. 36.

Tympanotonus fuscatus (L.) On the Anatomy; figs., J. Johansson.

Vermetus cristatus biondi in the western Mediterranean and plates form, R. Molinier.

#### AMALTHEACEA

Cymatiidae Keys and Lists, Anon (7).

#### CALYPTRAEACEA

Calyptraea Lamarck, 1799, gender feminine, Direction 42

Capulus ungaricus note on biology, M. Sharman.

†Capulus hemiastericolus sp. nov. as a commensal with Hemiaster batnensis preliminary note, R. Devries, C.R. Soc. geol. Fr. 7 1952, 1115 text-fig.

Crepidula fornicata Quantitative aspects of filter feeding, C. B. Jorgensen; C. fornicata biology and distribution in Essex waters, P. Walne (1).

#### STROMBACEA

†Arrhoges (Monocyphus) aegyptiacum sp. nov. Cretaceous. Monghara Massif, Sinai, Egypt, I. G. Mahmoud p. 150 pl. 17 f. 10-14.

†Dientomochilus armaudi sp. nov. Sentonian, S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 374 f. 68, 69. Lambis violacea and L. sebae, newly recorded from Indonesian seas, L. J. Butot.

†Pterodomta elajans sp. nov. Campanian, S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3—4 1954 p. 376 f. 7a.

Strombidae keys and Lists, Anon. (7). Strombus historical review of Linnaean species, H. Dodge.

†Strombus spinosus minihagali subsp. nov. Miocene, Minihagal Kanda, Ceylon, P. Deraniyagala p. 3 pl. 2 f. 1e, d, 2b.

Struthiolaria papulosa Quantitative aspects of filter feeding, C. B. Jørgensen-Struthiolaria vascular and nervous systems, J. E. Morton (4).

### ATLANTIDACEA

Firoloida desmaresti, sonic scattering layers due to swarming of this species, M. Blackburn.

#### NATICACEA

Falsilunatia powelli p. 74 f. 88, 129, 220 fathoms, Chatham Rise; F. subperforata p. 75 f. 83, 87, 330 fathoms, S.E. of Pitt Island spp. nov. Chatham Islands, New Zealand, R. K. Dell.

Lunatia heros and its natural control by Nassarius heros, A. H. Clarke, jr. (1).

†Natica and its allied group. Note on Japanese Tertiary and other forms, K. Hirayama p. 117; N. bevelandensis sp. nov. shell bed in the Westerschelde near Eidewoursdijk, Post-Pleistocene Holland Pouderoyan Basteria 20 4/5 Dec. 1956 p. 78.

Naticidae of the West Coast of America and some problems, J. Q. Burch (3); Naticidae of New Caledonia Anatomical studies, figs., J. Risbec (1).

Notocochlis luculentus (Iredale) description of operculum from shell previously known as Natica luculenta, L. Woolacott (1) p. 75 f. 2 5.

Uberella alacris sp. nov. 260–300 fathoms, off east Otago coast, New Zealand, R. K. Dell p. 77 f. 84 85.

#### LAMELLARIACEA

†Erato laevis punctata subsp. nov. Miocene Hungary, I. Csepreghyné-Meznerics p. 445 pl. 4 f. 4 5.

Velutina velutina association with the ascidian Styela, M. Diehl.

#### CYPRAEACEA

Cypraea guttata Gmelin from Tosa Bay, Japan, M. Azuma; C. (Erosaria) guttata Gmelin a new locality, M. Azuma; C. tigris L. in Oceania, A. C. Boullaire; C. arabica L. specimens and C. eglantina Dualos from Malaya, figs., R. J. Griffiths; C. zebra description and first record from Cozumel Is., Quintana Roo, off east coast of Mexico, E. Hidalgo, p. 8 pl. 5; C. cicercula L. a note and synonymy, G. Ransom.

Cypraeoidea nom. nov. for superfam. Cypraeacea, E. Hidalgo p. 8.

Erronea caurica brief description of living animal, F. McCamley.

Gratiadusta walkeri continens brief description of living animal, F. McCamley.

Pedicularia (Pediculariona) stylasteris Hedley var. notes on this and allied fauna, T. Habe (5).

Pediculariella probably a name for the younger stages of Pedicularia on Pediculariona, T. Habe (5).

#### TONNACEA

Bursidae Keys and Lists, Anon. (7).

Cassis cornuta first record from Bombay, India, V. Chari; C. madagascariensis spinella Clench and Cassis tuberosa (L.) predatory on Echinoderms, D. R. Moore.

Phalium granulatum predatory on Echinoderms, D. R. Moore.

†Phalium kanmonensis sp. nov. Tertiary, Watase, Hikoshima, Yamaguchi Prefecture, Japan, K. Hirayama p. 121, pl. 8 f. 10, 11.

Xenophalium (Xenogalea) abernethyi sp. nov. Off Castlepoint, east coast Wellington, in 40 fathoms, New Zealand, R. K. Dell p. 87 f. 119, 122

#### PTENOGLOSSA

#### **EPITONIACEA**

Amaea magnifica anatomical study and fig, of radula, I. Taki p. 74 pl. 13 f. 3; pl. 15; pl. 17 f. C.

†Claviscala livernantensis sp. nov. Campanian Livernant S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 350 f. 38. †Confusicala arnaudi sp. nov. Turonian S.W. France, G. Termier Bull. Soc. Hist. nat. Toulouse 89 3-4 1954 p. 351 f. 39.

Epitomiidae Keys and Lists, Anon. (9).

Epitonium scalale anatomical study and fig. of radula, I. Taki p. 71 pl. 13 f. 1; pl. 14; pl. 17 f. A.

Ianthina note on taxonomy and genital structure, D. Laursen; I. janthina biology, D. P. Wilson.

Janthina, ecological observations, figs., T. Okutani.

Melanella comatulicola (Graff) 1874, parasitic on Antedon mediterranea, J. P. Changeux.

Papyriscala latifasciata anatomical study (fig. of radula), I. Taki p. 75 pl. 13 f. 2, 4; pl. 16; pl. 17 f. B.

†Scala minihagali sp. nov. Miocene, Minihagal Kanda, Ceylon, P. Deraniyagala p. 3 pl. 2 f. 2c.

#### AGLOSSA

#### **PYRAMIDELLACEA**

"Balcis" otakauica p. 80 f. 90, 91; "B." alertae p. 81 f. 94, 95; "B." puhana p. 81 f. 93 spp. nov. 260-350 fathoms, off east Otago coast, New Zealand, R. K. Dell.

Enteroxenos parasitic degeneration of larvae, A. Iwanow (1).

Mucronalia nidorum sp. nov. from "Triton" dredgings off Palm Beach, Florida in 25 fathoms domiciled in spines of sea urchin, H. A. Pilsbry (1) Nautilus 69 4 Ap. 1956 p. 110 f. 4-8.

Odostomia corpulentoides sp. nov. New Zealand, Chatham Islands, 120-330 fathoms, R. K. Dell p. 157 f. 177; O. eulimoides parasitic on oysters and mussels, Cole & Hancock (1); O. impressa parasitizing oysters at South Carolina, U.S.A., S. Hopkins.

Planpyrgiscus lawsi sp. nov. 260-350 fathoms, east Otago coast, New Zealand, R. K. Dell p. 156 f. 176.

Pyramidella Lamarck, 1799, validation under the Plenary Powers, Opinion 386.

Stilifer neozelanica sp. nov. in 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 82 f. 107, 108.

Terelimella benthicola sp. nov. 260-350 fathoms, off east Otago, New Zealand, R. K. Dell p. 156 f. 178.

Turveria gen. nov. Stiliferidae, genotype: T. encopendena sp. nov. Cholla Cove, Sonora, Mexico and ectoparasitic on the echinoid Encope grandis Agassiz, S. Berry p. 356 f. 12.

Ulpia gen. nov. p. 65 Odontostominae genotype: U. venusta sp. nov. p. 67 f. 1-4 Lumbrera, Salta Province, Argentina, M. I. H. Scott (1).

#### STENOGLOSSA

### (a) Rachiglossa

#### MURICACEA

Apollon pusillus cuspidataeformis subsp. nov. Islet off Nagashima, Kii peninsula, T. Kira Venus 19 2 1956 122 text-figs.

Chathamidia gen. nov. Muricidae. genotype: C. expeditionis sp. nov. 260 fathoms, Chatham Rose, Chatham Islands, New Zealand, R. K. Dell p. 118 f. 159, 160.

Comptella coronata sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 117 f. 154.

Murex On the 'purple' products, Bouchilloux & Roche; On the purple therefrom and its biochemistry, Bouchilloux & Roche (1); M. trunculus on the purple pigment, Bouchilloux & Roche (2); M. tribulus Variation in the Great Bitter Lake, Egypt, figs., M. Taylor.

Nucella lapillus (L.) glandular histology, M. Pierson.

†Nucella (Purpura) lapillus (L.) in Iceland, S. Thoraninsson.

Poirieria kopua sp. nov. 290 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 114 f. 161, 162, 165.

Purpura. On the biochemistry of the purple colouring matter, Bouchilloux & Roche (1); Purpura lapillus chromosanal dimorphism and structural heterozygosity, H. Staiger.

†Taurasia szobensis sp. nov. Miocene. Hungary, I. Csepreghyne-Meznerics p. 445 pl. 3 f. 38, 41.

Terefundus axirugosus p. 115 f. 155; T. anomalus p. 115 f. 156 spp. nov.; T. quadricinctus unicarinatus p. 116 f. 112 subsp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell.

Thais haemostoma regeneration of drilling mechanism, Demoran & Gunter.

Urosalpinx cinerea capsule structure and hatching process, D. Hancock.

### BUCCINACEA

Aeneator marshalli separabilis p. 98 f. 135, Ohope Beach, Bay of Plenty; A. otagoensis cookianus p. 99 f. 141, 20–30 fathoms, off Kaikoura subspp. nov. New Zealand, R. K. Dell.

Afrocominella elongata fig. of radula, V. Orr p. 60 f. I i.

Anentome helena from Java, W. v. B. Jutting.

Antizafra aoteana sp. nov. 220 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 111 f. 110.

Austronassaria simplex gen. et sp. nov. Nassariidae 6-9 fathoms, Port Jackson, N.S.W. Australia C. F. & J. Laseron p. 71 f. 1, 2. Type in the Australian Museum.

Buccinoidea nom. nov. for superfam. Buccinacea, E. Hidalgo p. 10.

Buccinium historical review of Linnaean species, H. Dodge.

Buccinulum (Evarnula) flexicostatum sp. nov. 94 fathoms, Petre Bay, Chatham Islands; 300 fathoms off Otago coast, New Zealand, R. K. Dell p. 102 f. 138.

Buccinum Linn. 1758, gender neuter Direction 42.

Buccinum undatum general biology and economic value, H. A. Cole in M. Graham p. 179-180; B. undatum nervous and scent organs, figs., J. Demal; B. undatum neurohormones, J. Welsh (1).

Burnupena note on the systematics, studies on the radula, ecologic variation and early development, V. Orr.

Busycom canaliculatum blood pigments, E. Eliassen.

Columbella Lamarck, 1799, gender feminine, Direction 42.

Columbella rustica cytochemical comparison of atypical spermatogenesis and ovogenesis, B. Battaglia.

Columbellidae Keys and Lists, Anon (4).

Eucominia olsoni sp. nov. 61 fathoms, Mernoo Bank, Chatham Islands, New Zealand, R. K. Dell p. 95 f. 124.

Fasciolaria tulipa description and first record from Cozumel Is., Quintana Roo, off east coast of Mexico, E. Hidalgo p. 10 pl. 6.

Fasciolariidae Keys and List, Anon (7),

Fax alertae sp. nov. 220 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 92 f. 133.

Ilyanassa obsoleta development of isolated blastomeres, A. Clement.

†Japelion yabei sp. nov. from this Jôban Coalfield (Miocene) Japan, Y. Kamada Venus 18 3 1955 185 186 pl. 3 f. 1, 2.

†Lathyrus (Dolicolathyrus) hontensis sp. nov. p. 448 pl. 8 f. 7, 8; L. crassus vindobonensis subsp. nov. p. 447 pl. 7 f. 1, 2. Miocene, Hungary, I. Csepreghyne-Meznerics.

Liratilia conquisita chathamensis subsp. nov. p. 110 f. 114; L. isolata sp. nov. p. 111 f. 115, 130 fathoms, off the Forty Fours, Chatham Islands, New Zealand, R. K. Dell.

†Nassa (Hima) franzenaui p. 447 pl. 7 f. 32, 33; N. (Uzita) hontensis p. 446 pl. 7 f. 24, 25; spp. nov. Miocene, Hungary, I. Csepreghy-Meznerics.

Nassarius trivittattes a natural controller of Lunatia heros a predator of Mya, A. H. Clarke, jr. (1); Nassarius festivus nocturnal activity, S. Ohba.

†Nassarius (Uzita) malachiasi sp. nov. Pliocene, Marinha Grande, Portugal, L. Cox p. 7 Pl. 1 f. la, b; N. stocki p. 110, pl. 30 f. A-C; N. hildegardae p. 113 pl. 31 spp. nov. Pico formation, Pliocene, Los Angeles County, California, U.S.A., G. P. Kanakoff.

Neptunea decemcostata clenchi p. 116 pl. 7 f. 1 in 340-350 fathoms at 42° 46′ N. Lat. and 63° 22′ W. Lon., approximately 100 miles E.S.E. of Cape Sable, Nova Scotia; N. d. turnerae p. 117 pl. 7 f. 2, New Brunswick, Canada, Manan Island subspp. nov., A. H. Clarke, jr. Nautilus 69 4 Ap. 1956.

Penion benthicola sp. nov. 220 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 96 f. 125, 130, 131.

†Phos fusiformis sp. nov. Tertiary, Watase, Hikoshima, Yamaguchi Prefecture, Japan, K. Hirayama p. 85, p.125 pl. 8 f. 5-7.

Pleia cryptocarinata sp. nov. 290 fathoms, Chatham Rise, Chatham Islands, New Zealand, R. K. Dell p. 89 f. 137.

†Pollia (Tritonidea) londini Wrigley in the London Clay at Oxshott, Brown & Castell

†Pyrena (Columbellopsis) scaldensis sp. nov. Scaldisian (Upper Pliocene Dorst, province of Noord-Brabant, Netherlands, C. O. v. R. Altona (1) Basteria 20 4/5 Dec. 1956 p. 79; P. (Mitrella) sorongensis sp. nov. 200 m. off shore, Sorong, West New Guinea, A. C. v. Bruggen p. 12 f. 1.

Sipho verkrüzeni Kobelt, new to Iceland, fig., I. Osharsson (2).

†Siphonalia shojii sp. nov. Tertiary, Tomari, Hikoshima, Yamaguchi Prefecture, Japan, K. Hirayama p. 85 p. 122 pl. 8 f. 1-3.

Volutharpa ainov sp. nov. S. coast of Okhotsk Sea 120-130 fathoms, Kuroda & Kinoshita, Venus 19 1 1956, p. 19 text-fig.

Zemitrella benthicola sp. nov. 260-350 fathoms, off east Otago coast, New Zealand, R. K. Dell p. 110 f. 109.

### VOLUTACEA

Agaroniinae subfam. nov. of Olividae, Swainson 1840. Comprises the genus, Agaronia, A. A. Olsson p. 169.

Amoria grossi new record for New South Wales, Australia, J. Allan.

†Ancilla allani sp. nov. lower Eocene, Otaio Gorge, New Zealand, O. P. Olson p. 30 pl. 7 f. 13, 14.

†Ancilla marinhensis sp. nov. Pliocene. Marinha Grande, Portugal, L. Cox p. 9 pl. 1 f. 2a, b.

Aulica quaesita sp. nov. Northern Australia, Mid Arnhem Land, T. Iredale p. 76 f. 1.

Baryspira novaezelandiae benthicola subsp. nov. p. 124 f. 174, 175, 300 fathoms, Chatham Islands; B. bathami sp. nov. p. 124 f. 173, 300 fathoms, off east Otago. New Zealand, R. K. Dell.

†Baryspira (Baryspira) australis epacra subsp. nov. p. 12 pl. 1 f. 8-11 Pleistocene; B. (B.) erica p. 13 pl. 2 f. 6, 7 Pleistocene; B. (B.) gladiolaria p. 13 pl. 2 f. 4, 5 Pliocene-Pleistocene: B. (B.) oraria p. 14 pl. 2 f. 8, 9 Pliocene spp. nov. Baryspira (Gemaspira) subgen. nov. p. 14 Type: Ancilla (Baryspira) robusta Marwick 1924; B. (G.) pristina p. 16 pl. 3 f. 14, 15 Oligocene sp. nov.; B. (Spinaspira) subgen. nov.; p. 17 Type: B. (S.) stortha p. 18 pl. 4 f. 1, 2; B. (S.) cingulata p. 18 pl. 4 f. 5, 6; B. (S.) pakaurangiensis p. 19 pl. 4 f. 11, 12 spp. nov.; B. (S.) stortha irrisa p. 18 pl. 4 f. 3, 4 var. nov. Miocene; B. (Alocospira) cuppedia sp. nov. p. 22 pl. 5 f. 11, 12 Oligocene; B. (Gracilispira) subgen. nov. p. 23, Type: Ancillaria novaezelandiae Sowerby 1859; B. (G.) wairarapaensis p. 26 pl. 6 f. 25, 26 Miocene; B. (G.) firthi p. 26 pl. 6 f. 15-18 Pleistocene; B. (G.) gigartoides p. 27 pl. 6 f. 13, 14 Oligocene; B. (G.) rimuensis p. 27 pl. 6 f. 19, 20 Pliocene spp. nov. New Zealand, O. P. Olson.

Costellaria sculptilis interviewi var. nov. 8 miles west of Interview Island, Andamans, R. Ray p. 12 pl. 1 f. 3.

†Gemaspira subgen. nov. of Baryspira Olividae: Type: Ancilla (Baryspira) robusta Marwick 1924. New Zealand, O. P. Olson p. 14.

Gracilispira subgen. nov. of Baryspira Olividae Type: Ancillaria novaezelandiae Sowerby 1859. Fossil-Recent. New Zealand, O. P. Olson p. 23.

†Fulgoraria (Psephaea) kosibensis p. 304 f. 8; F. (P.) kamakurensis p. 304 [in text] f. 7 spp. nov. Kosiba sandstone, Pliocene, Miura peninsula, Japan, Y. Otuka.

†Indovoluta gen. nov. Volutidae, genotype Voluta humberti d'Archiac and Haima, Lower on Middle Eocene of Western Sind (Pakistan), F. E. Eames Proc. Malac Soc. 32 3 Dec. 1956 109 pl. 22.

Jaspidella gen. nov. p. 212 Olividae. Type: Voluta jaspidae Gmelin; J. miris sp. nov. p. 214 pl. 15 f. 2, 2a. Exact locality not known, maybe Gulf of Mexico? A. A. Olsson.

Leporemax chathamensis sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 120 f. 166, 167.

Macgintiella subgen. nov. of Olivella Swainson, 1831 Olividae. Type: O. watermani McGinty, A. A. Olsson p. 205. Marginella (Kogomea) hardingae p. 126 f. 180, Cook Strait, 75 fathoms; M. (Microvulina) vidae p. 126 f. 182 off White Rocks, Queen Charlotte Sound, 25 fathoms; M. (Glabella) judithae p. 127 f. 179, 130 fathoms, east of the Forty Fours, Chatham Islands; M. (G.) otagoensis p. 127 f. 183; M. (G.) cracens p. 128 f. 181; M. (volvarinella) fusuloides p. 128 f. 186. Off east Otago spp. nov. New Zealand, R. K. Dell.

Minioliva subgen. nov. of Olivella Swainson, 1831 Type: O. perplexa sp. nov. Florida Olividae, A. A. Olsson p. 209.

Mitra (Episcomitra) zonata, in the Bay of the Ciotat, S. France, J. Devidts.

†Mitra yokoyamai nom. nov. for M. ebenus, Yokoyama, 1920 (not of Lamarck), Y. Otuka p. 303.

Mitridae: review of forms from Indian waters, H. Ray.

Niteoliva subgen. nov. of Olivella Swainson, 1831, Olividae. Type: O. minuta (Link), A. A. Olsson p. 189.

Olivella nivea chiriquensis subsp. nov. p. 173 pl. 11 f. 4. North coast of Panama. Bocas del Toro; O. (Olivella) adelae p. 173 pl. 10 f. 8, 8a; pl. 16 f. 9, 9a. Florida; O. (O.) macgintyi p. 175, pl. 12 f. 3. Off Palm Beach, Florida; O. (O.) stegeri p. 176 pl. 14 f. 10. East of Alacran, Mexico, 28 fathoms; O. (O.) bitleri p. 178 pl. 15 f. 4, 4a. Panama Bay; O. (O.) broggii p. 179 pl. 9 f. 5, 5a. Northwest coast of Peru; O. (O.) riverae p. 179 pl. 10 f. 4, 4a. Northwestern Peru to Costa Rica; O. (O.) cocosensis p. 180 pl. 14 f. 9, 9a. Chatham Bay, Cocos Island, Pacific Ocean; O. (O.) drangai p. 180 pl. 15 f. 7. Galapagos Islands; O. (O.) rehderi p. 181 pl. 10 f. 7; fishands, O. (Niteoliva) peterseni p. 191 pl. 11 f. 7, 7a, b. Zorritos, Peru; O. (N.) morrisoni p. 192 pl. 15 f. 3, 3a. San José Island, Pearl Islands, Panama; O. (Callianax) thompsoni p. 197 pl. 14 f. 8, 8a. Off Marathon, Florida Keys, in 100 fathoms; O. (Minioliva) perplexa p. 209 pl. 16 f. 7. Florida; O. (M.) acteocina p. 210 pl. 12 f. 6. Bahamas and Panama spp. nov.; O. (Niteoliva) p. 189; O. (Pachyoliva) p. 201; O. (Zanoetella) p. 204; O. (Macgintiella) p. 205; O. (Minioliva) p. 209 subgen. nov., A. A. Olsson.

†Olivella (Olivella) venezuelensis p. 181 pl. 13 f. 3. Miocene, Venezuela; O. (O.) wilsoni p. 183 pl. 14 f. 4, 4a. Pliocene or Miocene, Acline, Florida; O. (Dactylidia) mansfieldii p. 187, Miocene, Jaskson Bluff, Florida; O. (Dactylidella) paraisa p. 188 pl. 13 f. 1. Miocene, Banana River, Costa Rica; O. (Niteoliva) terryi p. 193 pl. 13 f. 4, 4a. Miocene, N.E. Costa Rica; O. (N.) marksi p. 194 pl. 14 f. 6. Miocene, Ecuador; O. (Callianax) johnsoni p. 198 pl. 12 f. 8, 8a. Miocene, Chipola, Florida; O. (Pachyoliva) locklini p. 203 pl. 14 f. 5. Miocene, Alligator Creek, Florida; Spp. nov. O. (Niteoliva) p. 189; O. (Pachyoliva) p. 201; subgen. nov., A. A. Olsson.

Olivellinae subfam. nov. of Olividae, Swainson 1840. Comprises genus *Olivella* with several subgenera, A. A. Olsson p. 169.

Pachyoliva subgen. nov. of Olivella Swainson, 1831. Olividae. Type: O columellaris (Sowerby), A. A. Olsson p. 201.

†Spinaspira subgen. nov. of Baryspira Olividae: Type: B. (S.) stortha Olson sp. nov. Miocene, New Zealand, O. P. Olson p. 17.

Teremelon knoxi sp. nov. 320 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 121 f. 169, 172.

Turbinella Lamarek 1799, support for proposed preservation, C. Hubbs.

Turbinella Lamarck 1799, comments on proposed preservation of the name, Zilch, Mertens & Kraus.

Zanoetella subgen. nov. of Olivella Swainson, 1831 Type: O. zanoeta (Duclos) Olividae, A. A. Olsson p. 204.

Voluta (Amoria) caroli brief description of living animal, F. McCamley.

STENOGLOSSA

(β) Toxoglossa

#### CONACEA

†Acamptogenotia straeleni sp. nov. Miocene, Belgium, M. Glibert Mem. Inst. Sci. nat. Belg. No. 129 1954 p. 23 pl. iv f. 9.

Antiguraleus abernethyi p. 138 f. 200, east of the Forty Fours, Chatham Islands, 130 fathoms; A. pulcherrimus p. 138 f. 201; A. multistriatus p. 139 f. 202. Chatham Rise, Chatham Islands,

290-300 fathoms; A. fusiformis p. 140 f. 206, off east Otago, 260-350 fathoms spp. nov. New Zealand, R. K. Dell.

110

Antimelatoma canyonensis sp. nov. 260–350 fathoms, off east Otago, New Zealand, R. K. Dell p. 142 f. 196.

Asperdaphne expeditionis sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 137 f. 204, 205.

Asprella (Conasprella?) ichinoseana sp. nov., off Tosa p. 10 pl. 1 f. 5; A. i. prioris subsp. nov., off Tosa p. 11 pl. 1 f. 6; New Cones from Japan, T. Kuroda (1) Venus 19 I 1956.

†Brachytoma takeokensis sp. nov. Pliocene, Tomiya, Japan, Y. Otuka p. 307 f. 18.

Chelyconus (Profundiconus) profundorum p. 5 text-figs. 8, 9, 'S.W. part of off Tosa' deeper than 100 fathoms; C. kinoshitai p. 6 text-fig. 7 akin to fulmen Reeve, Kii, 100 fathoms spp. nov.; C. fulmen kirai subsp. nov. Tamoura, p. 7; New Conidae from Japan, T. Kuroda (1) Venus 19 1 1956.

†Clavatula neogradata nom. nov. for gradata Dollf. and Dautz. 1886 non Defrance in Bellardi p. 18 pl. iii f. 10a b, M. Glibert Mem. Inst. sci. nat. Belg. No. 129 1954.

†Clavus (Crassispira) powelli p. 26 pl. iv f. 12a b; C. (Cr.) staringi p. 30 pl. iv f. 15; spp. nov. Miocene, Belgium, M. Glibert Mem. Inst. sci. nat. Belg. No. 129 1954.

Comitas onokeana vivens subsp. nov. 260-350 fathoms, Chatham Islands and east Otago, New Zealand, R. K. Dell p. 131 f. 171.

Conus feeding habits, Anon. (10).

Conus, Observations on the stinging operation of various species, G. F. Kline.

†Conus (Leptoconus) asphaltodes sp. nov. Mio-Pliocene Waisioe Buton Malay Arch., C. Beets Leid. geol. Meded. 17 1953 p. 251 pl. 1, f. 1–2.

Conus striatus and C. catus feeding habits, A. J. Kohn.

†Cymatosyrinx hosonaga p. 305 f. 10 Plicene, Tomiya, Japan; C. hotei p. 305 [in text] f. 6, living form from the sea of Japan, spp. nov., Y. Otuka. Embrikena stupa p. 1 pl. 1 f. 1 off Tosa about 100 fathoms, Shikohu Id.; E. stupella p. 3, pl. 1 f. 3 off Tosa 60-70 fathoms, spp. nov. Japan, T. Kuroda (1) Venus 19 1 1956.

Fenestrosyrinx benthicola sp. nov. off east Otago, in 300 fathoms, New Zealand, R. K. Dell p. 141 f. 197.

Liracraca odhneri benthicola subsp. nov. 300 fathoms, off east Otago, New Zealand, R. K. Dell p. 140 f. 209.

†Leucosyrinx (Aforia) circinata minatoensis subsp. nov. Tomiya tuffaceous sandstone, Pliocene, Tomiya, Japan, Y. Otuka p. 305 f. 11.

†Loranko? akahida p. 308 f. 15; L.? tomiyaensis p. 308 f. 17 spp. nov. Pliocene, Tomiya, Japan, Y. Otuka.

†Mangelia turonensis decemcostata forma nov. p. 51 pl. vi f. 108; M. t. simplicior Peycot, M. Glibert Mem. Inst. sci. nat. Belg. No. 129 1954.

Micantapex parengonius sp. nov. 260 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 129 f. 168.

†Mitromorpha (Mitrithara) dollfusi payroti forma nov. p. 45 pl. vi f. 3; M. (Mitrithara) cancellata dautzenbergi forma nov. p. 46 pl. vi f. 5; M. Glibert Mem. Inst. sci. nat. Belg. No. 129 1954.

Mitrithara macphersonae p. 3 f. 1; M. bassiana p. 4 f. 2 spp. nov. south east of King Island, Bass Strait, Victoria, Australia, C. J. Gabriel. See also under Mitromorpha.

Nepotilla powelli sp. nov. 130 fathoms, east of the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 141 f. 198.

Pleurotoma (Pleurotomella) tcherniai sp. nov. Terre Adélie, J. M. Gaillard (2) Bull. mus. Hist. nat. Paris (2) 26 6 1955 p. 682 text-fig.

Rhizoconus (?) hirasei sp. nov. Kashiwajima, Tosa, Japan, T. Kuroda (1) Venus 19 1 1956 p. 3 text-fig. 10.

†Spirotropis (Typhlomangelia) kazusensis sp. nov. p. 306 f. 13; S. (T.) kazusensis inflatus subsp. nov. p. 307 f. 14 Pliocene, Tomiya, Japan, Y. Otuka.

Splendrillia benthicola p. 133 f. 192, 193; S. runcinata p. 133 f. 189; S. hermata p. 134 f. 187. Off Chatham Islands; S. jacula p. 134 f. 191, 300 fathoms, off east Otago; S. roseacincta p. 134 f. 199, 130 fathoms, east of the Forty Fours, Chatham Islands spp. nov. New Zealand, R. K. Dell.

Turris see Pleurotoma.

†Turris (Gemmulas) desiticula borealis forma nov. p. 5 pl. ii f. 2a, b, Miocene, Belgium, M. Glibert Mem. Inst. sci. nat. Belg. No. 129 1954.

Virroconus imperator sp. nov. Trinity Bay, Queensland, Australia, L. Woolacott (1) p. 72 f. 3.

Waipaoa marwicki sp. nov. 300 fathoms, Chatham Island, New Zealand, R. K. Dell p. 112, f. 113.

Xanthodaphne maoria sp. nov. 155 fathoms, S.E. of Pitt Island, Chatham Islands, New Zealand, R. K. Dell p. 136 f. 203.

Zeadmete otagoensis p. 113 f. 104; 260-350 fathoms, off east Otago; Z. ovalis p. 114 f. 103; 130 fathoms, east of the Forty Fours, Chatham Islands spp. nov. New Zealand, R. K. Dell.

# **Opisthobranchiata**

Roumanian opisthobranchs, A. V. Grossu (3); Opisthobranchiata of Lancashire and Cheshire, N. F. McMillan (1).

#### BULLACEA

Actaeon tornatilis anatomy of the pallial gonoduct and its phylogenetic significance, J. Johansson (1).

Acteon anatomy and systematic position, A. Pruvot-Fol (3).

Aglaja Renier, "[1807]" used in the Tavole serv. Class. Conn. Anim., future status to be considered, Opinion 427.

Akera bullata porphyrin pigments, Kennedy & Vevers.

Akera bullata first record from Plymouth area. Digestive system. Swimming, Morton & Holme.

Cylichna bulloidea sp. nov. off Pitt Island, Chatham Islands, New Zealand, R. K. Dell p. 144 f. 220, 221, 223.

Cylichna alba pp. 196-224; C. occulta pp. 225-239; C. magna pp. 240-250; C. cylindracea pp. 14-195; anatomy and histology, H. Lemche. Hydatina physis spawning habits, Baba, Hamatani & Hisai.

Neactaeonina inexpectata sp. nov. 300 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 147 f. 210.

Philine aperta population variations and effects of parasitism, J. M. Gaillard (3). Philine aperta micro-anatomy and histology of the digestive system, F. Starmühlner.

Rictaxis punctocolatus (Carpenter) Systematic position, fig. of radula, T. Habe (9).

Scaphander otagoensis sp. nov. off east Otago, New Zealand, in 260-350 fathoms, R. K. Dell p. 143 f. 208.

Volvulella truncata sp. nov. 260 fathoms, Chatham Islands, New Zealand, R. K. Dell p. 146 f. 211, 212.

#### APLYSIACEA

Aplysia fasciata bioelectric interactions, Arvanitaki & Chalazonitis; Aplysia effect of narcotics, Arvanitaki & Chalazonitis (1); A. neuronic biopotentials, Arvanitaki & Chalazonitis, A. & Chalazonitis, (2); A., on the giant nerve, Avanitaki-Chalazonitis, A. & Chalazonitis, N.; A. spawning habits of Japanese species, Baba, Hamatani & Hisai; A. depilans research on ovocite growth, O. Bolognan; Aplysia, nature of pigment, A. Christomanos; A. brasiliana; Adactylomela & A. juliana, description, discussion and occurrence, E. & E. Marcus; A. depilans micro-anatomy and histology of digestive system, F. Starmühlner; A. electric action in the nerve cells, L. Tauc.

Dolabella auricula cardiac nerves, A. Ebara.

### PTEROPODA

Pteropods from the Caribbean, A. D. Howard.

Cavolinia tridentata sexual differentiation in the gonads, Vitagliano-Tadini.

†Cavolina bituminata sp. nov. Mio Pliocene Waisiae, Dicton, Malay, Arch. C. Beets Leid. geol. Meded. 17 1953 p. 251, pl. f. 5-7.

†Hyolithes totoralensis sp. nov. Cambrian. El Totoral, Mendoza, Argentina, C. Rusconi (1) p. 116 f. 5.

†Middalya gen. nov. of Atlantidae genotype johnstonei sp. nov. Lower Permian Western Australia, J. M. Dickins Bull. 29 Com. Aust. dep. Nat. development Min. res. Geol. Geoph. 1956 p. 36 pl. vi f. 1-7 text-fig. 3.

### Nudibranchiata

Nudibranchiata of Saint Malo Bay, A. Franc; The preparation of specimens in extended state, G. D. Hanna; Viet-Nam Nudibranchia, J. Risbec; Nudibranchia & Ascoglossi from the Leghorn coast, Italy, Sordi & Majidi.

# SACOGLOSSA

#### ELYSIACEA

Alderia uda sp. nov. São Paulo and S. Vicente, Brazil, E. & E. Marcus (2) p. 8 f. 8-23.

Elysia marginata; E. atroviridis; E. halimedae spawning habits, Baba, Hamatani & Hisai.

Limapontia digestive system of unnamed species, F. Starmühlner.

Stiliger (Stiliger) talis sp. nov. São Paulo and S. Vicente, Brazil, E. & E. Marcus (2) p. 1 f. 1-7.

Thuridilla splendida digestive system, F. Starmühlner.

### **PLEUROBRANCHIACEA**

Berthella agassizii description of reproductive organs, E. & E. Marcus p. 20.

Euselenops luniceps from Sagami Bay, Japan. Fig. of radula, K. Baba.

Pleurobranchaea novaezealandiae spawning habits, Baba, Hamatani & Hisai; P. hamva sp. nov. S. Sebastião Island & Cananéia, Brazil, E. &. E. Marcus p. 21 f. 40-52.

### DORIDACEA

Aldisa sanguinea spawning habits, Baba, Hamatani & Hisai; A. nhatrangensis sp. nov. Hon lon, Nhatrang, Viet-Nam, J. Risbec p. 14 pl. xxii.

Archidoris montereyensis unusual carotenoids, Goodwin & Fox.

Argus Bohadsch (J.B.), 1761, direction that the suppression of the generic name by the Ruling in *Opinion* 185, be limited to suppression for the purposes of the Law of Priority, the status under the Law of Homonymy remaining unimpaired, **Opinion** 429.

Argus speciosus spawning habits, Baba, Hamatani & Hisai.

Dendrodoris (Dendrodoris) gemmacea; D. (Doriopsilla) miniata spawning habits, Baba, Hamatani & Hisai.

Discodoris notiperda sp. nov. Port Dayot, Viet-Nam, J. Risbec p. 15 f. 43-51.

Glossodoris pallescens; G. festiva; G. sibogae; G. aureopurpurea spawning habits, Baba, Hamatani & Hisai; G. coi sp. nov. beach at Hon lon, Nhatrang, Viet-Nam, J. Risbec p. 9, pl. xi f. 58, 59; pl. xxii.

Goniodoris castanea spawning habits, Baba, Hamatani & Hisai.

Homoidoris japonica spawning habits, Baba, Hamatani & Hisai.

Jorunna marchadi sp. nov. Ile Pattle, Viet-Nam, J. Risbec p. 16 f. 54-57.

Kalinga ornata general note, Pruvot-Fol (4).

Polycera (Greilada) elegans general note, Pruvot-Fol (4); P. quadrilineata digestive system, F. Starmühlner.

Phyllidia honloni p. 22 f. 71-75, 79-81, Hon lon beach, Nhatrang; P. serenei p. 24 f. 82-84, 86-89, Ile Poulo spp. nov. Viet-Nam, J. Risbec. P. catena Ile de France sp. nov., A. Pruvot-Fol J. de Conchyl. biol. Paris 96 2 July 1956 p. 70 f. vi, vii.

Phyllidiadae. A review of the family, figs., A. Prunot-Fol.

#### **AEOLIDIACEA**

Aeolidiella japonica spawning habits, Baba, Hamatani & Hisai.

Aeolidiopsis ransoni gen et. sp. nov. Aeolidinae? Kaukura Island, South Pacific, A. Pruvot-Fol (2) p. 228 f. 1-11.

Berghia coerulescens digestive system, F. Starmühlner.

Armina diqueti sp. nov. Gulf of California, U.S.A., A. Pruvot-Fol (1) p. 464 f. 8-10.

Cephalopyge trematoides general anatomy and taxonomy; C. mediterranea, C. picteti, C. orientalis, C. arabica and C. michaelsarsi are all considered synonymous with C. trematoides, J. E. Steinberg.

Cratena ornata spawning habits, Baba, Hamatani & Hisai. Dermatobranchus striatus; D. striatellus spawning habits, Baba, Hamatani & Hisai.

Favorinus japonicus spawning habits, Baba, Hamatani & Hisai.

Fimbriidae of Japan, Taki & Habe. Flabellina iodinca unusual carotenoids, Goodwin & Fox.

Hervia ceylonica; H. lineata spawning habits, Baba, Hamatani & Hisai.

Idulia (Doto) coronata digestive system, F. Starmühlner.

Melibe vexillifera spawning habits, Baba, Hamatani & Hisai.

### Pulmonata

Pulmonata Families, addenda, H. B. Baker (1); Family names in Pulmonata, H. B. Baker (3); New Caledonian pulmonates, A. Franc (2); Illustrated handbook of Swedish forms, B. Hubendick (2); Function of the radula in some pulmonates, K. Märkel; On the origin of Palmonata, J. E. Morton; Golgi bodies in the spermatogenesis, V. Nath.

#### ELLOBIACEA

Auriculodes gaziensis anatomy and biology, Knipper & Meyer.

Carychium tridentatum and Evolution of the Ellobiidae, J. E. Morton; C. tridentatum functional morphology Digestive and Reproductive Systems, J. E. Morton (2); C. tridentatum, distribution in Hungary, J. Vágvölgyi.

†Carychium minimum minor f. nov. Holocene, Velký Vřeštov, Bohemia, J. Petrbok (1) p. 8.

Cassidula angulifera. Evolution of the Ellobiidae, J. E. Morton; C. labrella anatomy, Knipper & Meyer.

Ellobiidae—biology and anatomy of six Tanganyikan species, Knipper & Meyer.

Ellobium auris-judae, auris-midae and Evolution of the Ellobiidae, J. E. Morton.

Leucophytia bidentata and evolution of the Ellobiidae, J. E. Morton.

Marinula filholi and Evolution of the Ellobiidae, J. E. Morton.

Melampus semisulcatus first record from Africa; M. fasciatus and M. semiplicatus anatomy and biology, Knipper & Meyer; M. bidentatus granifer, adansonianus, gundlachii and Evolution of Ellobiidae, J. E. Morton.

Ophicardelus costellaris in Evolution of the Ellobiidae, J. E. Morton.

Otina otis morphology and development, J. Morton (1).

Ovatella myosotis (Drap). Biology and Anatomy, figs., K. O. Meyer.

Ovatella myosotis in 'Evolution of the Ellobiidae', J. E. Morton; O. myosotis, The functional morphology, Digestive and Reproductive systems, figs., J. E. Morton (2).

Pedipes ovalis and Evolution of the Ellobiidae, J. E. Morton.

Plecotrema a conchological survey of the genus (maps and figures), B. Hubendick (3).

Pythia scarabacus, pollex and 2 spp. Evolution of Ellobiidae, J. E. Morton.

†Valenciennius annulatus in the Pliocene of Cocorova, Roumania, P. Nita.

#### **AMPHIBOLACEA**

Salinator Javanese species, E. v. B. Jutting.

#### LYMNAEACEA

Acroloxus lacustris (L.) in Merionethshire, L. Lloyd-Evans (1).

Acutorbis nom nov.; Carinorbis Mandahl-Barth 1954 non. Yen 1946 nec Conrad 1862, Mandahl-Barth, Arch. Molluskenk. Frankfurt a.M. 85 1/3 1956 p. 83.

Ancylidae, biology of middle European species, E. Frömming (4) p. 199–205.

Ancylus fluviatilis biology of populations at Windermere, R. Geldiay; A. fluviatilis breeding and growth in a mountain lochan, Isle of Skye, W. R. & M. R. Hunter (1); A. lacustris in Loch Lomond and in the Monkland Canal, W. R. Hunter (2).

Anisus (Armiger) crista (L.) and Anisus (Gyraulus) riparius (West.) distinctive characteristics, N. Odhner.

Australorbis glabratus use of radioisotopes in metabolic studies, with reference to molluscocide action, Azevedo & Gomes; A. glabratus olivaceus use of chromosome number in classification, Azevedo & Goncalves: A. glabratus control by Marsia cornuarietis and Helobdella fusca, Chernin, Michelson & Augustine; A. glabratus varying habitats in Puerto Rico, Harry & Cumbie; A. glabratus effect of stream gradient on the choice of habitat, Harry & Cumbie (1); A. centimetralis biospecific characterization and generic status, Paraense & Deslandes; A. inflexus sp. nov. Pouso Alegre, State of Minas Gerais, Brazil, Paraense & Deslandes (2) p. 149 f. 1-7; A. diagnostic characters of the Brazilian species, Paraense & Deslandes (3); A. glabratus effect of pentachlorophenol on metabolism, Weinbach & Nolan.

Biomphalaria pfeifferi use of radioisotopes in metabolic studies with reference to molluscocide action, Azevedo & Gomes; Biomphalaria infested with Chaetogaster, F. A. H. Bayer; Biomphalaria experiments on Schistosoma infection, C. C. Cridland; B. boissgi and B. truncatus. Control, S. El Din & A. El Nazar; B. glabrata (Say), anatomical figs., G. Ranson (3); B. smithi genital apparatus, G. Ranson (4); B. alexandrina. The shape of the prostate gland in geographical forms, E. Schwarz.

Bulinus tropicus and Chaetogaster F. A. H. Bayer; Bulinus, experimental Schistosoma infection, C. C. Caidland; B. truncatus seasonal rhythms in development, Gaud & Dupuy; B. truncatus Rhythms of Development, Gaud & Dupuy (1); B. contortus and Bilharziosis transmission, L. Lamy; B. cernicus (Morelet) and bilharziasis in Africa, P. L. Le Roux; B. (Physopsis) africanus B. (P.) globosus from Kenya, differentiation of forms likely to transmit Schistosoma, W. F. J. McClelland; B. (Physopsis) globosus as a vector of Bilharzia in Zambezia, T. d. Morais; Bulinus ecology of species carrying schistosomiasis in Gambia, S. Smithers; B. (Pyrgophysa) forskalii as a vector of Schistosoma haematobium, C. A. Wright; Bulinus species from Senegambia and their anatomy, figs., C. A. Wright (3); B. truncatus ecology, H. Zakaria.

Carnifex Keep, 1893 (erroneous spelling of Carinifex Binney, 1865) rejected; Carinifex Binney, 1865, validated; Carinifex Binney, 1863, rejected, Opinion 432.

Drepanotrema anatinum 491-499; D. melleum p. 527-534, shell and genital structure of Brazilian species, Paraense & Deslandes (1).

Ferrissia javana from Java, E. v. B. Jutting.

Galba palustris variations in size of successive generations, C. R. Boettger.

Gyraulus costulatus (Krauss) anatomy figs., G. Ranson (2).

Helisoma duryi normale, use of radioisotopes in metabolic studies, Azevedo & Gomes; H. duryi normale efficiency and action of various molluscocides, Azevedo & Gomes (1).

Indoplanorbis exustus note on development and environmental conditions, Kathuria, Rao & Hiregaudar-

Limnaea stagnalis structure of hermaphrodite gland, R. Aubry; L. stagnalis hermaphrodite gland, R. Aubry; L. stagnalis structure of hermaphrodite canal, Bounoure & Aubry; L. tenella Hutton, 1885, based upon juveniles of L. stagnalis (L.) of which it is a synonym p. 73; L. tomentosa Pfr. and L. leptosoma Hutton, removed to the genus Simlimnea Iredale p. 80, R. K. Dell (1); L. truncatula and L. auricularia rufescens, G. B. Kendall; L. stagnalis function of the radula, K. Markel; L. stagnalis effect of monovalent cations on the eggs, C. Raven.

Lymnaea on hermaphrodism and gland, R. Aubry (2); L. stagnalis possibility of autofecondation, R. Aubry (3); L. stagnalis variations in successive generations, C. R. Boeltger; L. peregra and truncatula occurring in gun pits on Bookham Common, C. F. J. Bensley; L. stagnalis appressa crossand self-fertilization, G. Cain; L. truncatula Müll. Anatomy of Japanese specimens compared with European material, figs., H. Itagaki; L. rubiginosa from Java, E. v. B. Jutting; L. stagnalis (L.) and associated species in Llyn Llywenan Anglesey, N. F. McMillan (5); L. auricularia (L.) Two forms in one pond, N. F. McMillan (4); L. humilis experimental infection with Fasciola hepatica, L. Mazzotti.

Lymnaeidae from Portuguese overseas provinces, Azevedo & Medeiros; L. biology of middle European species, E. Frömming (4) p. 65-134; L. development of the "mud-snail", Trousil-Linhart.

Megastropha Walker, 1918 (erroneous spelling of Megasystropha Lea, 1864) rejected, Opinion 432).

Megasystropha Lea, 1864, suppressed, Opinion 432.

Myxas all living species previously classed under this genus are shown to belong to one variable species, Simlimnea tomentosa (Pfr.), R. K. Dell (1) p. 80.

Physa Draparnaud, "[1801]", gender feminine, Direction 42.

Physa acuta efficiency and action of various molluscocides, Azevedo & Gomes (1); P. acuta note on the radula, K. Märkel p. 457 f. 6; P. rivularis experiments on the egg, V. Mancuso; P. acuta exchange of phosphorus in shells, M. Polster.

Physastra variabilis (Gray), proposed for all the living New Zealand forms of this genus, R. K. Dell (1) p. 87.

Physidae, biology of middle European species, E. Frömming (4) p. 135–157.

Physopsis africana and Chaetogaster, F. A. H. Bayer; P. globosa from the Giuba river, Italian Somaliland, E. Zavattari.

Planorbarius corneus (L.) alteration of spermatozoides, F. Alaphilippe; P. corneus use of radioisotopes in metabolic studies, with reference to molluscocide action, Azevedo & Gomes; P. corneus and P. metidjensi, use of chromosome numbers in classification, Azevedo & Gonçalves.

Planorbidae, eleven species from the area of Warsaw, S. Feliksiak; P. biology of middle European species, E. Frömming (4) p. 158-198; Planorbidae from Java, E. v. B. Jutting; Planorbidae: note on identification and ecology of schistosomiasis vectors, W. McClelland (1); Genetic approach to the systematics of Planorbidae, W. L. Paraense.

Planorbis newberryi Lea, 1858, specific name validated (sp. name of type sp. of Carinifex Binney, 1865), Opinion 432.

Planorbis susceptibility of different species to Schistosoma mansoni, Azevedo & Faro; P. crista, planorbis, and vortex occurring in crater and gun pits on Bookham Common, C. F. J. Bensley; P. corneus nerve and scent organs, figs., J. Demal; P. planorbis (L.), Westerhouses quarry pool, Glasgow district, W. R. Hunter (2); P. planorbis genital anatomy, Ranson & Medeiros.

Radix auricularia and R. peregra variations in successive generations, C. Boettger.

†Radix ovata minor, Budňany on the Berounka river; R. ovata minutissima, Únětice, ff. nov. Holocene, Bohemia, J. Petrbok (1) p. 8.

Simlimnea Iredale, now includes Limnaea tomentosa Pfr., L. leptosoma Hutton, Mysas arguta Hutton, M. ampulla Hutton, M. ampulla globosa Suter and M. ampulla waikariensis Cumber, R. K. Dell (1) p. 80.

†Stagnicola saratogensis sp. nov. North Park formation, Miocene of Wyoming, L. S. Russell J. Pal. 30 5 1956 p. 1260 text-fig. 1.

#### ONCHIDIACEA

Onchidella indolens from Brazil 76-81 pl. 23; O. remanei sp. nov. p. 81 pl. 24 f. 13-18, Tenerife, E. & E. Marcus (1); O. souriei sp. nov. Cap Blanc (Baie de l'Étoile) Tunisia, M. Gabe & M. Prenant Bull. Inst. franc. Afr. noire 17A l 1955 p. 91 f. 1-7.

### SUCCINEACEA

Boninosuccinea gen, nov. monotype Succinea punctulispira Pilsbry Endemic in the Bonin Ids., T. Habe (10) Venus 192 1956 p. 107.

Catinella (Quickella) arenaria found in Czechoslovakia, V. Ložek (3).

Neosuccinea gen. nov. Succineidae type: Succinea chinensis Pf. 1857, P. Matekin p. 902 f. 1, 2.

Oxyloma pfeifferi yamamotoi subsp. nov. Motaoshih Manchuria, T. Habe (10) Venus 19 2 1956 107.

Succinea Draparnaud, "[1801]", gender feminine, Direction 42.

Succinea elegans distribution P. Matekin (1).

Succineidae of Japan: Anatomical studies, T. Habe (10).

### ACHATINELLACEA

Tornatellina Pfeiffer, 1842, gender feminine, Direction 42.

### VERTIGINACEA

Abida frumentum from Tropical East Africa, C. Boettger (4).

Albinaria vermiculata janisadana p. 875 f. 3 Janisada Island; A. v. dragonarana p. 875 f. 4 Dragonara Island; A. v. paximadiana p. 877 f. 5, Paximadia Island subspp. nov. Dionysiades Islands, N.E. of Crete, F. E. Loosjes; A. (Albinaria) brevicollis theodori, Id. San Theodora Dodecanese p. 120 pl. 8 f. 9: A. (A.) brevicollis castellensis, I. of Rhodes p. 125 pl. 9 f. 14, 14a-15; A. (A.) brevicollis telensis I. of Piscopi p. 139 pls. 11 f. 31-32; A. (A.) brevicollis simiensis I. of Simi p. 140 pl. 11 f. 33-34; A. (A.) maculata kosensis I. of Kos p. 150 pl. 11 f. 37-38; subspp. nov. from the Dodecanese Islands, K. L. Pfeiffer Archiv. f. Mollusk. 84 4/6 1955; Albinaria of Dodecanese A. vesti (Bttgr) f. suturalis Bttgr; v. phalanga Bttgr. orientalis Bttgr. types figd., K. L. Pfeiffer (1) Arch. Molluskenk. Frankfurt a.M. 85 pl. 5; teres lara Bttgr.; v. paucicosta Bttgr.; strigata orientalis Bttgr.; pl. 7: teres famosa Bttgr.; teres armathia Bttgr.; oertzeni camura Bttgr.; privigna Bttgr.; pl. 8. Types of Albinaria, figured, K. L. Pfeiffer (1) Arch. Molluskenk. Frankfurt a.M. 85 4/6 pls. 5-8; megalostoma Bttgr.; parallelifera Bttgr., K. L. Pfeiffer (1) Arch. Molluskenk, Frankfurt a.M. 85 pls. 6; Albinaria of the Dodecanese, K. L. Pfeiffer.

Barcania gen. nov. Clausiliidae Alopiinae; genotype Clausilia bengasiana Gambetta, R. A. Brandt, Arch. Molluskenk. Frankfurt a.M. 85 4/6 121 pl. 9 f. 1 text-fig. 1; Barcania (Barcania) nizeli p. 124; pl. 9 f. 2 fig. 2-4; B. (B.) sasaensis p. 126 pl. 9 f. 3 text-fig. 5; B. (B.) kaltenbachi p. 128 pl. 9 f. 4 text-figs. 6-7; spp. nov. B. (B) k. apolloniana subsp. nov. p. 132 pl. 9 f. 6 text-fig. 8; B. (B) klugi p. 133 pl. 9 f. 7 text-fig. 9; B. (B.) nagaensis 137 pl. 10 f. 11 text-fig. 11 spp. nov. Cyrenaica, pls. 9 10; B. kaltenbachi albaensis p. 131 pl. 9 f. 5; B. klaptoczi gahhami p. 136 pl. 10 f. 9; B. klaptoczi gambettae p. 137 pl. 10 f. 10; subspp. nov. Cyrenaica, R. A. Brandt Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956.

Boninena gen. nov. Enidae Type species: Buliminus callistoderma Pilsbry Japan, T. Habe (6) p. 265 f. 3.

†Chondrina avenacea minor; C. avenacea elongata ff. nov. Hostím, Bohemia, Holocene, J. Petrbok (1) p. 8.

Clausilia Draparnaud, "[1805]", gender feminine, Direction 42; C. dubia Deap. 2200 feet Highcup Sill; C. rugosa in the Dumfriesshire Cheviots to 1100 feet, E. Blezard.

Cochlicopa nitens (Kokeil) Gallenstein and C. minima Siemashko, usually considered as varieties of C. lubrica (Müller) are two good species, A. Nilsson.

Cochlodina laminata albanica subsp. nov. Albania, S. Jaeckel (See Jaeckel and Meise) p. 28 f. 2.

Delima ornata note, V. Hudec (1).

Ena montana (Draparnaud) at Little Munden, Hertfordshire, L. Lloyd-Evans.

Euphaedusa cumingiana recondita note on anatomy, F. Loosjes (2).

Gracilinenia aequistriata sp. nov. Valle de Chanchamayo on the highway from Hacienda Naranjal near the village San Ramón to the mine "Pichita-Caluga", about 1300 m. probably, Peru, W. Weyrauch (1) Nautilus 69 4 p. 110 pl. 6 f. 2.

Herilla excedens albanica subsp. nov. Korca, Albania, S. Jaeckel (see Jaeckel and Meise) p. 29 pl. 1 f. 3.

Laciniaria (Denticularia) denticulata sprattiana nom. nov. for Clausilia denticulata spratti Boettger 1883 non Pfeiffer 1846, K. L. Pfeiffer Archiv. f. Moll. 84 4/6 1955 p. 153.

Leucochila von Martens, 1860, gender feminine, Direction 42.

Leucostigma leucostigma from Tropical East Africa, C. Boettger (4).

Luchuena gen. nov. Enidae. Type species: Buliminus eucharistus Pilsbry. Genus also includes L. reticulata (Reeve) from Amami and Okinawa; L. taivanica (Moellendorff) from Taiwan, Japan, T. Habe (6) p. 265 f. 1, 12.

Nenia (Columbinia) zischkai sp. nov. Yungas de Palmar, 700 m. typical rain forest, Bolivia, W. Weyrauch (1) Nautilus 69 4 Ap. 1956 p. 113 pl. 6 f. 3 3a; Nenia in Argentina, Anon. (2); N. argentina H. Scott in the Argentine, Anon. (3). Papillifera bidens from Tropical East Africa, C. Boettger (4).

Paraphaedusa minahassae, first record from New Guinea, F. Loosjes (1).

Pupilla alpicola found in Czechoslovakia, V. Ložek (3).

†Pupilla slovenica sp. nov. Pleistocene Sárovce, Hron river, South Slovakia, J. Petrbok (2). p. 92 text-f. 1.

Temesa pilsbryi sp. nov. El Infiernillo Rio Rimac 3360-3370 m. road from Lima nr. Oroya, Peru, W. K. Weyrauch (2) Arch. Molluskenk. Frankfurt a.M 85 4/6 1956 146 pl. 11 f. 1-4.

Torbana subgen, nov. of Barcania q.v. Clausiliidae subgenotype Barcania (Torbana) pabsti Cyrenaica R. A. Brandt Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956 p. 140 pl. 10 f. 13 text-figs. 14-15.

Truncatellina strobeli first record from Hungary, J. Vágvölgyi.

Vertigo substriata from Western Bohemia, V. Ložek (4); Vertigo ronnebyensis not specifically distinct from Vertigo aretica, H. W. Waldén.

†Vertigo praecoquis sp. nov. North Park formation, Miocene of Wyoming, L. S. Russell J. Pal. 30 5 1956 p. 1262 text-fig. 4.

Yakuena gen. nov. Enidae. Type species: Ena nesiotica Pilsbry et Hirase Japan, p. 264 f. 11 Genus also includes Y. hachijoensis (Kuroda) from Hachijojima, T. Habe (6).

Zebrina (Subzebrinus) drangiana sp. nov. Bashgul-Tal, Nuristan, eastern Afghanistan, S. Jaeckel (1) p. 344 f. 2, 3.

# ACHATINACEA

Achatina fulica occurence of disease, A. Mead; A. fulica, a criticism of Dr. Mead's paper, Nautilus 69 2 57, C. E. Pemberton.

Achatininae in the Muséum de Paris, M. Denis.

Archachatina (Tholachatina) saskai p. 345 f. 1-3 text-f. 1-3; Ihagana Forest, Iringa; A. (T.) gebhardti p. 351 f. 4-6, text-f. 5, 6. Mufindi, Iringa spp. nov. Tanganyika, E. Africa, H. Knipper.

Hohenwarthia orghidani sp. nov. Roumania, A. V. Grossu (2) p. 1579 f. 1. Lamellaxis gracilis living in greenhouses, E. Karlin.

Limicolariopsis wagneri sp. nov. p. 353 f. 7-10, Loliondo, Tanganyika, E. Africa, H. Knipper.

Opeas pumilum living in greenhouses, E. Karlin.

Rumina decollata influence of environment, E. Frömming (2).

Synapterpes (Promoussonius) propinquus sp. nov. Vulcano Puraco, Colombia, T. E. Crowley J. Conch. London. 24 4 July 1956 144 pl. 6.

#### **OLEACINACEA**

Araucania twomeyi gen. et sp. nov. Endodontidae, Rio Pascua, southern Chile, J. Parodiz p. 17 fig.

Oleacina Röding, 1798, gender feminine, Direction 42.

#### ZONITACEA

Acutidens subgen. nov. of Polygyra Say Type: P. acutedentata (W. G. Binney) Polygyridae, H. A. Pilsbry (3) p. 26.

Agriolimax reticulatus electrical activity in pedal ganglion, Hughes & Kerkut; A. reticulus sensitivity of pedal ganglion to osmotic pressures, Kerkut & Taylor; A. reticulatus (Müller) feeding on carrion, G. W. Pitchford; A. sibiriensis [sp. nov.] form of so called "Limax hyperborrus" from Jakutsk, H. W. Walden (1) Arkiv. f. Zool. (2) 10 5 1956 p. 354; A. altaicus transetorius forma nov. from Jenissei, H. W. Waldén (1) Arkiv f. Zool. (2) 10 5 1956 p. 356 [nom et figs.] and p. 361; A. reticulatus colour variation, B. Wood.

Anademus altivagus (Theobald). Origin of Golgi dictysomes, figs., Nath & Chopra; A. altivagus morphology of golgi bodies, R. Rishi.

†Anguispira holroydensis sp. nov. North Park formation, Miocene of Wyoming, L. G. Russell J. Pal. 30 5 1956 p. 1262 text-fig. 3.

Arion rufus tentacular regeneration, M. Chétail; A. ater sensitivity of pedal ganglion to osmotic pressures, Kerkut & Taylor; Arion taste and scent senses, R. Kittel; A. circumscriptus Johnson, A. rufus L. A. subfuscus Drap., A. hortensis Fér., A. intermedius Normand: genital organs, figs., M. Lamesch; A.

rufus germinal regeneration after castration, **P. Laviolette** (1); A. rufus genital system anatomy, figs., **P. Laviolette** (2); Arion rufus regeneration experiments, **P. Laviolette** (3).

†Arion kinkelini and A. hochheimensis notes on fossil species described by Wenz 1911, H. Bräm.

Baiaplecta nom. nov. = Pangania Laidlaw Proc. malac. Soc. 20 (2) 83 non Pangania Poppius 1914 Acta Soc. Sci. Fenn. 44 (3) 47 Hamiptera; F. F. Laidlaw, Arch. Molluskenk. Frankfurt a.M. 85 1/3 1956 p. 83.

Bulines from Portuguese Overseas provinces, Azevedo & Medeiros.

Charopa coma variation in the shell, H. E. Whitten.

Corilla lesleyae sp. nov. Ceylon, Ullswater Estate near Kitulgala, 2000 ft., G. A. S. Barnacle J. Conch. 24 3 Feb. 1956 95 pl. 4.

Daudebardia brevipes in Bergen, W. R. Schlickum.

Deroceras reticulatum (Müll.) D. agreste (Müll.), genital organs, figs., M. Lamesch.

Discus brunsoni Berry, in McDonald Cirque, Mission mountains, western Montana, R. A. Brunson.

Erymodon subgen. nov. of Polygyra Say Type: P. ventrosula (Pfr.) Polygyridae, H. A. Pilsbry (3) p. 20.

Helicodiscus (Hebetodiscus) singleyanus inermis H. B. Baker, figures of shell and radula and map of distribution in Europe, J. G. J. Kniper Arch. Molluskenk. Frankfurt a.M. 85 4/6 163-169.

Lehmannia marginata Müll. genital organs, fig., M. Lamesch.

Limax Linn. 1758, masculine gender, Direction 42.

Limax scent and taste senses, R. Kittel; L. flavus L. L. lencophaeus; L. maximus L.; L. albipes Dam and Moelt, M. Lamesch; L. cinercaniger Wolf. Genital organs, fig., M. Lamesch; L. marimus genital system, figs., P. Laviolette (2); L. duplex from the Tamborin grotto, Como, Italy, C. Maviglia; L. hyperboreus Westerlund, Notes from Siberian specimens, H. W. Waldén (1).

†Limax fossil found in a cave in the Urio valley, Como, Italy, Cadeo & Senna.

Malacolimax tenellus Nilsson, genital organs, fig., M. Lamesch.

Megapallifera subgen. nov. type Pallifera mutabilis Hubricht (1951 Naut. 65, 57), L. Hubricht (2) Nautilus 69 4 April 1956 p. 126.

Mesodon normalis (Pils.) a distinct species, L. Hubricht (3).

Milax rusticus Mull. genital organs, figs., M. Lamesch.

Milax sowerbii anatomy, fig., A. R. Schouten.

Monophysis subgen. nov. of Polygyra Say Type: P. behri (Gabb) Polygyridae, H. A. Pilsbry (3) p. 25.

Nipponarion gen. nov. genotype N. carinatus sp. nov. His-amatsu-yama, Tottori City, Japan, Yamaguchi & Habe Venus 18 4 1955 pp. 237, 238.

Oxychilus Fitzinger, 1833, validated, gender masculine, Opinion 431.

Oxychilus glaber note on the radula K. Märkel p. 457 f. 6; O. cellarius (Müll.) a sinistral example from Sweden, H. Schlesch (1); O. draparnaldi, H. A. Schmidt.

Polygyra bicruris insculpta p. 21; P. hindsi heteroea p. 24 pl. 2 f. 2 subspp. nov. Nayarit; P. idiogenes p. 30 pl. 3 f. 1, 1a, b sp. nov. Northern Mexico; P. (Solidens) p. 27; P. (Acutidens) p. 26; P. (Monophysis) p. 25; P. (Erymodon) p. 20 subgen. nov., H. A. Pilsbry (3); P. scintilla sp. nov. Wallacy Co. along the railroad 15 m. N. of Raymondville, Texas, Pilsbry & Hubright Nautilus 69 Jan. 1956 p. 94 pl. 5 f. 3, 3a.

†Slug granules from post glacial deposits in the Lea Valley, J. F. Hayward.

Solidens subgen. nov. of Polygyra Say Type: P. anilis (Gabb) Polygyridae, H. A. Filsbry (3) p. 27.

Systrophia stenogyra zischkai a new locality in Bolivia, W. Blume; S. (Systrophia) siolii sp. nov. Serra Formosa nr. Mulata, Monte Alegre, Staat. Pará. Brasil, F. Haas Archiv. f. Mollusk. 84 1/3 1955 103 text-figs 4-6.

Ventridens intertextus eutropis Pilsbry living in Cheatham Co., Tennessee, M. C. Teskey.

Vitrea contracta from Central Bohemia, V. Ložek (5).

Vitrina Draparnaud "[1801]", gender feminine, Direction 42.

Vitrina the Palaearctic species and V. pellucida islandica subsp. nov. N.W. Island Isafjördur, L. Forcart Archiv. f. Mollusk 84 4/6 1955 p. 156 pl. 12 f. 3, 6.

Vitrinidae in the Dolomites, L. Forcart (1).

Zonitoides arboreus living in greenhouses, E. Karlin.

### ARIOPHANTACEA

Degneria subgen. nov. of Helicarion, for Helicarion boranensis Piersanti Corydon Memorial Museum, Nairobi, Verdcourt Proc. Malac. Soc. London 32 1/2 Aug. 1956 p. 72 text-fig. 1.

Machrochlamys (Parvatella) ghorbandensis sp. nov. Ghorband-Tal, Afghanistan, S. Jaeckel (1) p. 347 f. 5, 6.

Videnoida subgen. nov. of Videna for Trochomorpha gouldiana Pilsbry, T. Habe (2) Venus 18 4 1955 pp. 231–234 f, 2.

### **ACAVACEA**

Strophocheilidae of the Museum National d'Histoire natrelle de Paris, E. Salmon.

Strophocheilus (Strophocheilus) pudicus roseolabris p. 33 pl. 3 f. 3; pl. 24 f. 3; S. pudicus debilis p. 34 pl. 16 f. 1; pl. 17 f. 2; S. planidens fusoides p. 43 p. 9 f. 2; Pl. 12 f. 2; pl. 25 f. 2 subspp. nov. Brazil; S. atavus p. 47 pl. 8 f. 3 sp. nov. Uruguay; S. (Megalobulimus) oblongus albescens nom. nov. p. 73 for Bulimus oblongus var. alba E. A. Smith, 1894, non. B. oblongus var. albus Bland and Binney 1872; S. oblongus perelongatus p. 77 pl. 22 f. 3 Colombia; S. oblongus elongatus p. 78 pl. 1 f. 4; pl. 2 f. 4 Uruguay; S. oblongus conicus p. 79 pl. 6 f. 1; pl. 19 f. 2 pl. 23 f. 1, 3 Brazil; S. oblongus musculus p. 82 pl. 12 f. 6; pl. 20 f.1; pl. 31 f. 4 Paraguay subspp. nov.; S. terrestris oliveirai p. 117 pl. 5 f. 2; S. sanctipauli pygmaeus p. 138 pl. 15 f. 5; S. sanctipauli eyerdami p. 139 pl. 24 f. 2; S. granulosus abbreviatus p. 164 pl. 27 f. 4; S. granulosus foreli p. 165 pl. 28 f. 3 Brazil subspp. nov., J. C. Bequaert.

#### BULIMULACEA

Anostoma baileyi sp. nov. Salobra, Matto Grosso, Brazil, A. Solim (1) Notulae Naturae No. 287 1956 4 pl. 1 f. 1-3.

Bostryx (Platybostryx) fisheri sp. nov. below Tarma on the Tarma-Chanchamayo road, Peru, H. A. Pilsbry Nautilus 69 Jan. 1956 p. 92 pl. 5 f. 4, 5.

Bulimulidae from the rivers of Alabama, Georgia and Florida, U.S.A., Clench & Turner p. 122; Bulimidae from Java, E. v. B. Jutting.

Bulimulus (Scansicochlea) hyltonscottae p. 59 f. 1-4; B. (S.) strobeli p. 62 f. 5 spp. nov. San Luis Province, Argentina, J. Parodiz (1); B. (Scansicochlea) catamarcanus p. 77 f. 6; B. (S.) rudisculptus p. 78 f. 7 spp. nov. Catamarca Province, Argentina, J. Parodiz (2); B. (Bulimulus) araozi sp. nov. Tingo Maria, Peru, W. K. Weyrauch (2) Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956 p. 149 pl. 11 f. 8.

Cerion Röding 1798, neuter gender, Direction 42.

Cerion eximium lerneri subsp. nov. Bimini Islands, Bahamas, W. J. Clench (5) p. 1 f. 1, 2; Cerion geographic variation and hybridization in Bahama populations, Mayr & Rosen.

Cyclodontina (Cyclodontina) inflatus salobrensis subsp. nov. Salobra, Matto Grosso, Brazil, A. Solem (1) Notulae Naturae No. 287 1956 7 pls. 1 f. 22.

Macroceramus loeryi sp. nov. Puerto Rico M. K. Jacobson Arch. Molluskenk. 84 1/3 1955 97 text-figs. 1-3.

Naesiotus elegantulus p. 4 pl. 1 f.1; N. andivagus p. 5 pl. 1 f. 2; N. gracillimus p. 6 pl. 1 f. 3; N. pilsbryi p. 6 pl. 1 f. 4; N. haasi p. 7 pl. 1 f. 5, 5a; N. zilchi p. 9 pl. 1 f. 6 spp. nov. Peru, W. Weyrauch.

Partula taeniata genetic diversity, D. W. Bailey.

Stenostylus zilchi sp. nov. Peru, W. K. Beyrauch (2) Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956 p. 156 pl. 11 f. 18.

Thaumastiella subgen. nov. of Thaumastus to include T. sarcochrous and T. glyptocephalus (Pilsbry), W. Weyrauch p. 11.

Thaumastus (Thaumastiella) subgen. nov. Bulimulidae, W. Weyrauch p. 11

#### HELICACEA

Arianta arbustorum function of radula K. Markel p. 454 f. 2.

Bradybaena (Chosenelix) problematica (Pilsbry) 1927, formerly known as Chloritis (C.) problematica, T. Habe (7) p. 192.

Cathaica rufispira hispida subsp. nov. Sarckanda Mts., Badakschan, Afghanistan, S. Jaeckel (1) p. 349 f. 8.

Cepaea nemoralis (L.) up to 1600 feet on Cross fell, E. Blezard; C. nemoralis effect of temperature on histological evolution of the ovotestis, J. Bouillon; C. nemoralis genetic stability of populations, C. Goodhart; C. nemoralis, shell banding patterns in two colonies in Ontario, W. Judd; C. nemoralis, on a colony and variation in same as regards colour and banding, M. A. Landman.

Epiphragmorphora birabeni p. 93 f. 1; E. villavilensis p. 94 f. 1 spp. nov. Catamarca Province, Argentina, J. Parodiz (3); E. ormeai sp. nov. Peru, W. K. Weyrauch (2) Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956 p. 158 pl. 11 f. 19.

Euhadra amaliae studies on the eye Hara & Hara.

†Euomphalia obvoluta unidentata var. nov. Holocene, Bohemia, J. Petrbok (1) p. 8.

Euparypha Hartmann, 1844, rejected, Opinion, 431.

Euparypha pisana Müll. ssp. arietina Rossm. from S. Spain, C. Sacchi (3).

Helicella Lamarck, 1812, rejected Opinion 431.

Helicella Férussac, 1821, validated: type species designated as Helix itala Linn. 1758, Opinion 431.

Helicella caperata (Mont) up to 1200 ft on Newbiggin Fell, E. Blezard.

Helicellinae Hesse, 1926, family group name validated; Helicellinae Chenu, 1859, family-group name rejected, Opinion 431.

†Helicodonta obvoluta at Bobbitshole, Ipswich, A. G. Davies (1).

Helix Linn. 1758, gender feminine, Direction 42.

Helix incurva Renier "[1804]", the specific name, as used in the above combination, rejected, Opinion 427

Helix cartusiana Müller (O.F.), 1774 (specific name of type species of Monacha); H. cellaria Müller (O.F.) 1774 (specific name of type species of Oxychilus); H. itala Linn., 1758 (specific name of type species of Helicella); H. pisana Müller (O.F.) 1774 (specific name of type species of Theba), above species validated; H. rhodostoma Draparnaud, "[1801]" rejected, Opinion 431.

Helix pomatia experiments on the heart, L. Acolat; H. pomatia and effect of drugs on the heart, L. Acolat (2); H. pomatia experiments on the heart, Acolat & Cornier; H. pomatia spermatocyte structure, J. André; H. pomatia holochemical localization of & McGeeβ-glucuronidase, Billett Russell: H. pomatia, the use of glucosamine in experiments on the heart, Binet, Jullien, Ripplinger & Carnot; H. adspersa experiments on heart. M. Corda and M. Corda (1): H. adspersa experiments on cardiac cycle before and during stimulation of estrinsic nerves, M. Corda (2); H. pomatia L. Chemical studies and alimentation, Dastugue, Bastide and Moinet; H. pomatia, aspersa composition of blood-pigments, E. Eliassen; H. pomatia structure, A. Fabricius; H. pomatia L. structure, Grassé, Carasso & Favard; H. pomatia, chromosome ultrastructure. The spermatozooid, Grassé, Carasso & Favard (1); H. poma tia structure of mitochondria in spermatozoide, Grassé, Carasso & Favard (2); H. pomatia spermatogenesis, Grassé, Carasso & Favard (3); H. pomatia golgi elements of the neurones, Gresson, Threadgold & Stinson; H. aspersa, the first in Hawaii, Y. Kondo; H. aspersa a second specimen in Hawaii, Y. Kondo (2); H. nemoralis function of the isolated heart. Jullien. Cardot & Vieille-Cessay; H. pomatia experiments on the heart movements, Jullien, Ripplinger & Guyon; H. pomatia survival of the isolated heart at low temperature in different media, Jullien, Cardot & Rerat; H. pomatia experiment on heart, Jullien, Ripplinger & Joly; H. pomatia influence of various substances on heart ventricle action, Jullien, Ripplinger & Joly; H. pomatia experiments on inhibition of the heart and the action of the visceral nerve, Jullien, Ripplinger & Joly; H. pomatia experiments on heart activity, Jullien, Ripplinger & Cardot; H. pomatia

experiments on automatism of the heart, Jullien, Ripplinger & Cardot; H. pomatia production by the heart of a substance equivalent to acetylcholine. Jullien, Ripplinger & Cardot; H. pomatia reports of acetylcholine production by the heart, Jullien, Ripplinger & Cardot; H. pomatia Nervous systems, Jullien & Ripplinger; H. pomatia, nerve structure, Jullien & Ripplinger; H. pomatia, experiments on heart, & Ripplinger; H. pomatia Jullien atropine-calcium antagonism. Jullien & Ripplinger; H. pomatia, course of exerction, K. Nejedlý; H. pomatia blood-sugar level relative to temperature, Pavlović & Bać; H. adspersa experiments on the heart, relative to Starling's Law, Peruzzi & Corda; H. pomatia influence on isolated heart, of extract of Ecbalium elaterium, Pora & Rosca; H. aspersa distribution and. dispersal by human agency, W. J. Rees; H. pomatia action of K+ and C++ irons on activity of the heart, Ripplinger & Joly; H. pomatia experiments on the heart, Ripplinger & Joly (1); H. morphology and special technique, A. L. Rogue; H. pomatia, electrical effects observed in cellular ganglions, L. Tauc (1); H. pomatia reaction of the isolated heart to acetylcholine, Velluda, Ticsa & Csutak.

†Helix (Archhelix) betieri sp. nov. p. 129 pl. 1 f. 9, 10, 13a-c; H. (A.) betieri conica var. nov. p. 129 pl. 1 f. 8; H. (A.) tectiformis sp. nov. p. 127 pl. 1 f. 2, 4, 5, 6, 7a, b, 11a-c; Miocene from environs of Gambatta, David & Jodot Bull. Soc. Hist. Nat. Afrique Nord 46 3/4 1955.

Hygromia (Ponentina) subvirescens Bellamy between Beer Head and Branscomb, S. Devon, A. G. Davis (2).

†Hygromia (Ponentina) subvirescens (Bellamy) in Holocene deposit at Haslyn Bay near Padstow, Cornwall, A. G. Davis (3).

Iberus note on systematics, Garcia San Nicolás & Alvarado.

Jacosta Gray (J.E.), 1821, suppressed, see Opinion 431.

†Leptaxis neivai sp. nov. nom. nud. Miocene, Porto Santo Island, Atlantic ocean, N.E. of Madeira, Silva, G. H. da p. 35.

Leucochroa candidissima polymorphism, J. Altes; L. candidissima shell polymorphism, J. Altes (1).

Monacha Fitzinger, 1833, validated Opinion 431.

Murella (Ambigua) fuscolabiata from Tropical East Africa, C. Boettger (4).

Nipponochloritis subgen. nov. of Trichochloritis subgenotype Chloritis fragilis Gude, T. Habe (2) Venus 18 1955 pp. 226, 232.

Opica fuscolabiata brunellae subsp. nov. Pizzo Salvatesta, Messina Province, Sicily, C. F. Sacchi (2) p. 3 f. 1 2.

Planatella Clessin, 1876, rejected, Opinion 431.

Psadariella subgen. nov. of Solaropsis subgenotype Solaropsis (Psadariella) palizae sp. nov. Peru, W. K. Weyrauch (2) Arch. Molluskenk. Frankfurt a.M. 85 4/6 1956 pp. 159-160 pl. 11 f. 15-17.

Pyrenaearia on the anatomy and systematics of the Spanish species; P. cantabrica cavadongae p. 44 f. 14; P. carascalensis parva p. 56, Sierra del Coli, province of Lérida vars. nov.; P. poncebensis sp. nov. p. 46 f. 5, 16, 22, 23, 25–27 Spain, A. O. Lopez.

†Tachea (= Cepaea) hortensis f. minor depressa subform. nov.; T. (= C) hortensis minutissima form. nov. Holocene, Smíchov, Bohemia, J. Petrbok (1) p. 8.

Theba Risso, 1826, validated Opinion 431.

Theba carthusiana in Mecklenburg, Germany, H. A. Schmidt (1).

Trichia erjaveci hajlensis subsp. nov. Hajla-Planina, Macedonia, S. Jaeckel (see Jaeckel & Meise) p. 29 pl. 1 f. 4.

Xerophila Held, 1837, rejected, Opinion 431.

Xeropicta gyroides from Tropical East Africa, C. Boettger (4).

Yakuchloritis subgen. nov. of Trichochloritis subgenotype Chloritis albolabris Pilsbry and Hirase, T. Habe (2) Venus, 18 1955 p. 224, 232 f. 11.

#### STREPTAXACEA

Artemon pilsbryi sp. nov. Salobra, Matto Grosso, Brazil, A. Solem (1) Notulae Naturae No. 287 1956 11 pl. i f. 4-6. Gonaxis kibweziensis and its economics relative to Achatina, A. R. Mead (1).

Gulella (Pupiquiella) pupa mueriensis p. 108 f. 16; G. tudes minor p. 122 f. 23; vars. nov.; G. (Paucidentina) oloitae p. 114 f. 19; G. bolocoensis p. 124 f. 24; G. (Costigulella) pooensis p. 129 f. 26; G. (Molarella?) fernandensis p. 132 f. 27; G. (M?) mongolae p. 134 f. 28; G. escalerai p. 136 f. 29 a, b; G. cacuminata p. 138 f. 30; spp. nov. Fernando Póo Island, W. Africa, López & Rocandio; G. (Primigalella) ndamanyiluensis sp. nov. Tanganvika. West Usambaras. Mt. Ndamanyilu, 5 m. west of Malindi, 5,800-7000 ft., Venmans Basteria 20 4/5 Dec. 1956 p. 74 text-figs. 7, 8, 9; G. intrusa sp. nov. East Usambaras, Mt. Bomola, Tanganyika, Verdcourt (1) Proc. Malac. Soc. London 32 1/2 Aug. 1956 p. 69 text-figs. 4, 5; G. usagarica msambaa subsp. nov. W. Usambaras in Coryndon Museum, Verdcourt Basteria 20 4/5 Dec. 1956 p. 73.

Haplotrema concavum (Say) and H. kendeighi Webb described as a subspecies of H. concavum, are distinct species, L. Hubricht (1).

Ptychotrema (Ennea) bonyomae p. 99 f. 12; P. (E.) malaboensis p. 101 f. 13; spp. nov. Fernando Póo Island, W. Africa, López & Rocandio.

Streptaxis (Streptartemon) extraneus sp. nov. Serra Formosa bie Mulata, Monte Alegre Staat Pará, Brazil, F. Haas Archiv. f. Mollusk. 84 1/3 1955 102 text-figs. 1-3.

Streptostele (Streptostele) mínima sp. nov. Fernando Póo Island, W. Africa, A. O. de Z. López & A. O. de Z. Rocandio p. 87 f. 4.

### SCAPHOPODA

Spermiogenesis in some species, A. Franzén (1); Scaphapoda secretions, M. Gabe (1).

†Cadulus (Dischides) yatalensis sp. nov. Pliocene, Australia, N. H. Ludbrook p. 4 pl. 1 f.3, 4 [nom. nud. Tr. roy. Soc. S. Aust. 77 1954: p. 58.]

Dentalium Linn. 1758, neuter gender, Direction 42.

Dentalium vulgare population variations and effect of parasitism, J. M. Gaillard (3); D. (Tesseracme) hancocki sp. nov. Off west coast of Mexico, W. K. Emerson (1) p. 3 f. 1. †Dentalium (Fissidentalium) mawsoni p. 2 pl. 1 f. 5, 6; D. (Antalis) denotatum p. 3 pl. 1 f. 7-9 sp. nov. Pliocene, Australia, N. H. Ludbrook [nom nud. Tr. roy. Soc. S. Aust. 77 1954: p. 58.]

†Siphonodentalium (Pulsellum) adelaidense sp. nov. Pliocene, Australia, N. H. Ludbrook p. 4 pl. 1 f.1, [nom. nud. Tr. roy. Soc. S. Aust. 77 1954: p. 58.]

# Pelecypoda or Lamellibranchiata

Fine structure of the shells of some lamellibranchs, G. Athanassopoulos; Spanish Cretaceous Lamellibranchia, J. Bataller (1): Physiology of nutrition and Water conservation, W. v. Buddenbrock pp. 241-255; Spermiogenesis in some species, A. Franzén (1); Naming Carboniferous non-marine lamellibranchs, R. Eager; Lamellebranchiata histological notes on neuro-secretion in numerous species, M. Gabe (2); Lamellibranchs from Kerguelen and Heard Islands, J. M. Gaillard; Lamellibranchs from the Thyrrhenian sea, off the Tuscan coast, some listed for the first time from this locality, B. Gianfranco; Structure and diversity of form in the lamellibranchs, H. Janus; Fluviatile and lacustrine Pelecypoda from Victoria, Australia, R. C. Kershaw; growths and decline of population and distribution, D. Nicol; Taxonomic significance of gaps in morphology, D. Nicol (2); Observations on stomach and digestive divitacula, G. Owen; Stomach types of Protobranchia and Septibranchia, R. D. Purchon (3); Lamellibranch larvae in the North Sea. C. B. Rees; Lamellibranchs of the genera Inoceramus, Didymotis, Liopistha, Lopha, spp. from the La Suma formation of Western Venezuela, Rutsch & Salvador; Pelecypoda from Strangford Lough and neighbouring coasts, Ireland, G. Williams.

### **PRIONODESMACEA**

†Edmondia sibirica spec. nov. Verkhoyansk ridge, Russia, Permian, Kulikov, M. V. C.R. Acad. Sci. Moscow 101 6 1955 p. 1102.

†Grammysia quadrata p. 273 pl. 1 f. 6, 7; G. obliqua p. 274 pl. 1 f. 8, 9; spp. nov. Silurian Pentland Hills nr. Edinburgh, A. Lamont Proc. roy. soc. Edinb. 65B 3 1954.

#### NUCULACEA

Austrotindaria benthicola p. 13 f. 14, 15; A. flemingi p. 14 f., 12, 13, 16 spp. nov. 330 fathoms south-east of Pitt Island, New Zealand, R. K. Dell.

Huxleyia Adams, April, 1860 has precedence over Cyrilla Adams, June, 1860. Cyrillista Iredale, 1929 and Diabolica Jousseaume, 1897 are synonyms, H. E. Vokes.

Jupiteria wolffi sp. nov. 155 fathoms, South-east of Pitt Island, New Zealand R. K. Dell, p. 13, f. 17, 18.

Neilo rugata sp. nov. c. 300 fathoms, E.-N.-E. of Taiaroa Head, New Zealand, R. K. Dell p. 14 f. 62, 63.

Nucinella S. Wood, 1861, the type of the genus is N. miliaris "Deshayes" Wood [non Deshayes] = Pleurodon ovalis Wood, from the Pliocene of England. Pleurodon Wood, 1840 non Harlan 1831, Nuculina d'Orbigny, 1844 non Porro 1837, Cyrillona Iredale, 1929 and Neoplorodon Hertlein and Strong, 1940 are synonyms of Nucinella. H. E. Vokes J. Paleont. 30 3 1956 pp. 652-671.

†Nucinella oregonia sp. nov. p. 657-658, text-f. 2A-2C, Upper Eocene, Nestucea formation of Oregon, H. E. Vokes J. Paleont. 30 3 1956 pp. 652-671.

Nucula strangeiformis sp. nov. 220-300 fathoms, Chatham Rise, New Zealand, R. K. Dell p. 9 f. 2, 3.

†"Nucula" iwayai sp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, K. Ichikawa p. 269 pl. 10 f. 7; Nucula aegyptiaca p. 115 pl. 9 f. 4-9; N. chavani p. 116 pl. 9 f. 14-16 spp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud; N. prepercrassa sp. nov. Cretaceous. Ochillee Creek, Georgia, U.S.A., L. Stephenson (1) p. 234 pl. 39 f. 1-5.

†Nuculana lyonsensis p. 8 pl. 1 f. 7-9; N. thomasi p. 10 pl. 1 f. 1-2, 10; spp. nov. Lower Permian, Western Australia, J. M. Dickins Bull. 29 Com. Austr. dep. Nat. Develop. Min. Res. Geol. Geoph. 1956; N. insolita sp. nov. Devonian, Buchan Caves limestone, Victoria, Australia, J. A. Talent p. 48 pl. 4 f. 9.

Nuculidae—anatomy of the stomach and digestive diverticula, G. Owen (1).

†Palaeoneilo tenelliformis sp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa in Ichikawa p. 271 pl. 10 f. 8.

†Prilukiella gen. nov. p. 138 type: P. janischewskyi sp. nov. p. 140 pl. 18 f. 1-16; P. janischewskyi infurcatidentata var. nov. p. 141 pl. 18 f. 11, 12. Tatarian Stage, Triassic. Lower Sukhona River, Russia, M. Plotnikov.

Saccella hedleyi morioria subsp. nov. 155-330 fathoms, Chatham Rise, New Zealand, R. K. Dell p. 12 f. 10, 11.

†Trigonucula gen. nov. p. 267, Nuculidae, Genotype: T. sakawana sp. nov. p. 267, 268, pl. 10 f. 1-3; T. sakawana tokombensis p. 269 pl. 10 f. 4-5; T. sakawana lata p. 269 pl. 10 f. 6 vars. nov. Triassic, Sakawa Basin, Shikoku, Japan, K. Ichikawa.

Yoldia hyperborea limatuloides subsp. nov. Fossfjord (Arnarfjord) N.W. Iceland, Ockelmann Medd. Grønland 107 7 1954 p. 11 pl. 1 f. 2 pl. 2 f. 1, 2.; G. myalis, sapotilla, limatula, hyperborea, and h. limatuloides key to species, W. K. Ockelmann.

†Yoldia hikoshimensis sp. nov. Tertiary, south of Watase, Hikoshima, Yamaguchi Prefecture, Japan, K. Hirayama p. 84, p. 102 pl. 6 f. 1.

#### ARCACEA

Anadara inflata guanine-deaminating activity and nitrogen metabolism, S. Ishida; A. inflata distribution of uricase activity, Ishida, Tsuzuki & Nitahara; A. antiquata (L.) A. a. scapha Meuschen new to Seychelles, Rost & T. Soot-Ryen; A. inflata researches on haemoglobin, Yagi, Mishima, et al.

Arca subdisparilis nom. nov. for Arca disparilis Orb., D. Nicol (1) J. Paleont. 28 I 1954 p. 96.

†Arca sampelayoi sp. nov. Jurassic. Maranchón, Castell, Concha & Revilla; A. turonica konkensis p. 30 pl. 2 f. 1-4; A. noae usboica p. 30 pl. 1 f. 10-14 subspp. nov. Miocene. Turkmenstan & W. Kazakhstan, Merklin & Nevesskaya.

Austrosarepta benthicola sp. nov. 260-350 fathoms, off E. Otago coast, New Zealand, R. K. Dell p. 15 f. 26, 29.

Barbatica (Cucullaearca) ovata new to Seychelles, Rost & T. Soot-Ryen.

†Breviarca unisulcata sp. nov. Cretaceous, Shishi-jima, southwest Japan, M. Amano p. 66 pl. 1 f. 6-8; B. subinflata p. 236 pl. 39 f. 6-8. Ochillee Creek, Georgia; B. symmetros p. 236 pl. 40 f. 1, 2. Hurstboro-Marvyn road, Russell County, Alabama spp. nov. Cretaceous, U.S.A., L. Stephenson (1).

†Cucullaea (Megacucullaea) nov. sp. aff. C. (M.) eminens Cox Kimmeridgian, Katrol series, Kutch, India, S. Agrawal (2) p. 14 pl. 1 f. la-b; C. (Cucullaea) coxi p. 59 pl. 8 f. 3a-b; C. (Megacucullaea) sp. nov. af. C. (M.) eminens, Cox; p. 61; S. K. Agrawal (1) Ann. Cent. et Doc. Paléont. No. 19 1956.

Cucullaeidae review of genera and subgenera with types, D. Nicol (1).

†Cyrtodontarca note on structure, N. Yakovlev.

†Gignouxiarca gen. nov. Arcidae; G. orientalis sp. nov. Cretaceous. Moghara Massif, Sinai Egypt, I. G. Mahmoud p. 117 pl. 9 f. 17, text-f. 55.

†Glycymeris idensis p. 267 pl. 38 f. 1-5; G. flabellata p. 268 pl. 38 f. 9-13. Miocene. Miyagi Prefecture; G. compressa p. 269 pl. 38 f. 6-8 Akashiba formation. Fukushima Prefecture; G. osozawensis p. 270 pl. 38 f. 14 a-c, Pliocene, Yamanashi Prefecture. spp. nov. Tertiary, Japan, S. Kanno.

Glycymeris sect. Pectunculus Lamark 1799; Glycymeridae p. 794 and Pectunculus Da Costa synonym of Dosinia fam. Veneridae p. 888.

Hochstetteria sculpturalis sp. nov. 130 fathoms off the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 17 f. 21, 22.

†Mnataia gen. nov. p. 4, Genotype: M. přibyli sp. nov. p. 5 f. 1-9, Devonian. Holyně, Bohemia, Prantl & Růžička.

†Pectunculus (Axinaea) obtusatoformis sp. nov. Tortonian Miocene. S.W. Russia, V. Ghoretzkii p. 200 pl. 2 f. 4.

Stenzelia nom. nov. for Trinacriella Macociel 1937 non Del Guerico 1913 nec Parona 1933 "type (of Twinacriella): Pectunculus perplanus Courad," F. S. Macneil J. Paleont. 28 2 1954 p. 217. †Trigonarca inflata sp. nov. Cretaceous. Hurstboro-Marvyn road, Russell County, Alabama, U.S.A., L. Stephenson (1) p. 237 pl. 40 f. 4-8.

[1956]

Tucetoma subtilis sp. nov. off the south shore of Bermuda at about 100 fathoms, D. Nicol (3) Nautilus 70 2 Oct. 1956 51 pl. 3 f. 1, 2.

### MYTILACEA

Amygdalum papyria Conrad in Maryland waters of Cheshapeake Bay, J. F. Allen (1).

†Arcomytilus arnali sp. nov. Cretaceous. Valencia, Spain, J. Bataller p. 285 1 fig.

Austromytilus gen. nov. Mytilidae: genotype: Mytilus rostratus Dunker, New South Wales, Australia, C. F. Laseron (1) p. 265 f. 4, 5.

Botula foliata behaviour within boring, form and method of boring figs., C. M. Yonge (1).

Brachidomtes Aragomite percentage, H. A. Lowenstan.

†Chondrodonta munsoni from the Cretaceous. Yugoslavia, M. Pleničar.

Fluviolanatus amarus sp. nov. coastal lagoons, N.S.W. Australia, C. Laseron (1) p. 274 f. 46-49.

†Hilgardia gen. nov. Mytilidae: genotype. Hilgardia compressa sp. nov. Cretaceous, Ripley Formation, Union County, Mississippi, L. W. Stephenson J. Paleont. 30 3 1956 p. 752–753, text-f. 1A–D.

Lithophaga calcifer from coral New South Wales, Australia, L. Woolacott; Lithophaga plumula, Habitat mantle and siphon, behaviour in boring, method of boring and nature of boring, C. M. Yonge (1).

†Modiola manzouriana sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 104 pl. 6 f. 1 text-f. 49.

Modiolaria commensal with Ascidians, A. Bourdillon; M. marmorata. Quantitative aspects of filter feeding, C. B. Jørgensen.

†Modiolopsis finlayi sp. nov. Silurian, Portland Hills, nr. Edinb., A. Lamont Pr. roy. Soc. Edinb. 65B 3 1954 p. 282 pl. 1 f. 1. Modiolus peronianus p. 269 f. 22-24, Port Hacking; M. cottoni p. 270 f. 25-28 30-50 fathoms off Twofold Bay spp. nov. N.S.W., Australia, C. Laseron (1).

Musculus alganus sp. nov. N.S.W., Australia, C. F. Laseron (1) p. 268 f. 13, 14.

†Musculus conditus sartasi p. 98 pl. 29 f. 6-10; M. naviculoides maximus p. 99 pl. 29 f. 21-23 subspp. nov. Miocene. Turkmenstan and W. Kazakhstan, Merklin & Nevesskaya.

†Myoconcha piveteani sp. nov. Callovian Kutch India Agrawal (1) Ann. Cent. ét. Doc. Paléont. No. 19 1956 p. 93 pl. 10 f. 13a-c.

Mytilaster lineatus in the Caspian Sea, L. Zenkevitch.

Mytilus Linn. 1758, gender masculine Direction 42.

Mytilus edulis oxidation of L-amino acids, Blaschko & Hope; M. edulis v. galloprovincialis, conditions necessary for spawning H. Bouxin; M. edulis growth of forms at Boulogne-sur-Mer. H. Brienne: M. edulis diegensis population variations in California, W. R. Coe; M. edulis, M. galloprovincialis general biology and culture, H. A. Cole in M. Graham p. 162-168; M. californianus growth rates in different populations, P. Dehnel; M. edulis seasonal variation of chemical contents. F. Fraga; Mytilus chemical contents, Glahn, Manchon & Roche. M. edulis culture in Holland, B. Havinga; M. edulis mechanism of anterior byssus retractor muscle, Hoyle & Lowy; M. edulis quanine-deaminating activity and nitrogen metabolism, S. Ishida; M. edulis distribution of uricase activity Ishida, Tsuzuki & Nitahara; M. edulis egg maturation, K. Iwata; Mytilus: On the muscle, figs., J. Lowy; M. edulis L., on neuro-secretion, P. Lubet; M. edulis effect of removal of nervous centres on emission of gametes, P. Lubet (2); M. edulis, P. Lubet (1); M. edulis influence of Mytilicola intestinalis on the gonad development, H. Mann; M. edulis attacked by Mytilicola intestinalis, Meyer-Waarden & Mann; M. edulis culture in the German Waddensea, Meyer-Waarden; M. edulis relation between zinc and sperm motility, T. Mizuno; M. edulis adaptation to salinity fluctuations, M. Motwani; M. edulis osmotic and ionic regulation, W. T. W. Potts (1); M. galloprovincialis Lmk. sexuality, E. Ricci; M. edulis difference in size of forms from the White & Okhotsk Seas, Russia, A. Savilov (1); M. edulis studies on the acrosome, Wada, Collier & Dan.

†Mytilus razmyslovae sp. nov. Miocene. Turkmenstan and W. Kazakhstan, Merklin & Nevesskaya p. 96 pl. 28 f. 1, 2.

†Opokiella gen. nov. p. 92, type: O. tschernyschewi p. 93 pl. 10 f. 8, 9; O. tetraedroides p. 94 pl. 10 f. 6, 7 spp. nov. Tatarian stage, Triassic? Sukhona River, Russia, M. Plotnikov (1).

Septifer australis sp. nov. 14 fathoms off Long Reef, N.S.W., Australia, C. F. Laseron (1) p. 266 f. 6, 7.

†Stutchburia muderongensis sp. nov. Lower Permian Western Australia, J. M. Dickins Bull. 29 Com. Aust. dep. Nat. Develop. Min. res. Geol. Geoph. 1956 p. 14 pl. 1 f. 12–17.

Volsella metcalfi early life history, H. Yoshida.

†Volsella weaveri sp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro p. 192 pl. 2 f. 4.

### **PTERIACEA**

†Actinopecten (?Tolmaia) midlothiana sp. nov. Silurian Pentland Hills, nr. Edinburgh, A. Lamont Proc. roy. Soc. Edinb. 65B 3 1954 p. 279 pl. 1 f. 3, 4.

†Actinopteria meridionalis sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 68, pl. 7 f. 19.

Atrina vexillura Born new to Seychelles, Rost & T. Soot-Ryen; A. pectinata japonica early life history, H. Yoshida.

†Avicula cygnipes Phillips 1829, Avicula longicosta Stutchbury 1839, figured, **H. Hölder.** 

†Aviculopecten netschajewiformis sp. nov. Verkhoyansk ridge, Russia, Permian, Kulikov, M. V. C.R. Acad. Sci. Moscou 101 6 1955 p. 1102.

†Avicolupinna (?) Sibirica sp. nov. Verkhoyansk ridge, Russia. Permian, Kulikov, M. V. C. R. Acad. Sci. Moscou 101 6 1955 p. 1102. †"Cassianella" dubia sp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa (1) p. 184 pl. 6 f. 10a-b.

Electroma tragulata from coral at Queensland, Australia, L. Woolacott.

†Entomonotis multistriata p. 255 pl. 9 f. 11, 14; E. tenuicostata p. 259 pl. 9 f. 6, 7 spp. nov.; E. zabaikalica intermedia p. 258 pl. 10 f. 19-20; E. tenuicostata mabara p. 260 pl. 9 f. 8 vars. nov. Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa (2).

†Halobia aotii nom. nov. p. 185 for H. mullistriata Kobayashi and Aoti, 1943; H. kashiwaiensis sp. nov. p. 185 pl. 6 f. 11, Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa (1).

†Inoceramus from the Trans-Carpathian flisch, Tertiary, Vyalov & Pasternak.

†Leioptera jacksonensis p. 46, pl. 3 f. 15; L. buchanensis p. 47, pl. 3 f. 14; pl. 4 f. 8 spp. nov. Devonian. Buchan caves, Victoria, Australia, J. A. Talent.

Meleagrina margaritifera containing a fish trapped inside it's shell and covered with nacre, A. Zänkert.

†Modiomorpha tenuilineata p. 48, pl. 4 f. 3-6; M. concentrirugosa p. 48 pl. 4 f. 1, 2 spp. nov. Devonian. Buchan Caves, limestone, Victoria, Australia, J. A. Talent.

†Oxytoma scanica (Lundgren) in the Planorbis-Zone in Wurtemberg, figs., H. Hölder; O. longicostata (Stutchbury) Middle Lias, England, figd., H. Hölder; O. koniensis sp. nov. Rhaetian, Triassic. Okhotsk Sea, River Vilighi, Koni Peninsula, Russia, I. Tuchkov p. 189 pl. 3 f.1-4.

Parviperna nucleus Lmk; P. dextata Krauss new to Seychelles, Rost & T. Soot-Ryen.

Pinna sunricata L. new to Seychelles, Rost & T. Soot-Ryen; P. fragilis Quantitative aspects of filter feeding, C. B. Jørgensen.

Pinctada martensi physiology of sensory responses, I. Honjo; P. martensii use of radioautography to study pearl formation, Okada, Tsujii, Watabe & Yoshii; P. maxima damage to shell

by boring animals, Takemura & Okutani; P. martensii, dahllite as a constituent of prodissoconch, N. Watabe; P. martensii shell structure, Watabe & Wada.

†Plesiopinna gen. nov. p. 70 Pinnidae. genotype P. atriniformis sp. nov. p. 71 pl. 1 f. 1-5. Cretaceous, Shishi-jima, southwest Japan, M. Amano.

†Posidonia permica sp. nov. Peruvian East Greenland, N. D. Newell Medd. Grønland 110 4 1955 pp. 16, 21 pl. 2 fs. 5, 6; Posidonial camerata sp. n. carboniferous Frankenberg, E. Paproth Paläont. Z. 27 3-4 1953 p. 191 pl. 12 f. 11.

†Promyalina groenlandica sp. nov. Permian East Greenland, N. D. Newell Medd. Grønland 110 4 1955 pp. 17 25 pl. 5 f. 11-13.

†Pseudoptera securiformis sp. nov. Cretaceous. Alabama and Georgia, U.S.A., L. Stephenson (1) p. 238 pl. 40 f. 9, 10; pl. 41 f. 12-14.

Pteriidae, Mesozoic forms in the Pacific area, R. S. Allan p. 367.

†Pterinea concentrilamellata sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 69 pl. 7 f. 30, 31.

†Pterochaenia fragilis (Hall) a pseudoscalar pelecypod from the Devonian of the Ural, B. Nalivkin.

Vulsella living in a sponge, K. v. Haffner.

#### PECTINACEA

†Camptonectes khatschiljorense spec. nov. Tadzhikidsan depression, Khochilor. Paleogene, Korobkov, I. A. Trud. Leningrad Soc. Nat. 68 2 1951 p. 67 pl. I f. 3-6.

Chlamys (Radulopecten) Callovian Kutch, India, S. K. Agrawal (1) Ann. Cent. ét Doc. Paléont. No. 19 1956 p. 71 pl. 8 f. 12; C. varia growth, Audouin & Letaconnoux; C. varia general biology, Letaconnoux & Audouin; C. varia L. On neuro-secretion, P. Labet; C. varia effect of removal of nervous centres on emission of gametes, P. Lubet (2); C. cuneola Reeve; C. cfr. rubromaculata Swby. new to Seychelles, Rost & T. Soot-Ryen; C. farreri growth and reproduction, Tchang-Si, Tsi shi (1); C. opercularis distribution and growth in Danish and Faroese waters, E. Ursin.

†Chlamys mojsisovicsi sp. nov. p. 165 pl. 5 f. 1-5; C. mojsisovicsi toyamai var. nov. p. 166 pl. 5 f. 6-8 Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa; Chlamys zimistanensis sp. nov. p. 91 pl. 3 f. 1, 2 South Western Fergana. Paleogene; C. necopinata sp. nov. p. 92 Paleogene; C. (?) foediaurita sp. nov. p. 93 pl. 2 f. 18 Fergana, Dzhalal-abad, Changir-tash. Paleogene; C. sumsarica Vialov var. fastosa var. nov. Fergana, Sokh, Pishtan, Changir-tash, Tadzhi-kistan depression, Aruk-tan, Tereklitan p. 58 pl. 4 f. 2–4, 6, 8, 10, 11; C. incelebrata sp. nov. p. 99 pl. 3 f. 21 Fergana, Rishtan, Paleogene; C. abominosa sp. nov. p. 69 pl. 1 f. 17, 18 Tadzhikistan depression, Aruk-Tan, Russia. Paleogene; C. vialovi sp. nov. p. 80 pl. 2 f. 3-9 Isfara, Tadzhikistan depression, Tutkaul, Khochilor, Vakhshe river valley. Paleogene; C. alajensis sp. nov. p. 79 pl. 2 f. 2 Tadzhikistan depression, Estrom-tan Russia; C. estimiensis sp. nov. p. 77 pl. 2 f. 1 Tadzhikistan depression, Estim-tan, Russia. Paleogene; C. edecimata sp. nov. p. 76 pl. 1 f. 7, 7a Tadzhikistan depression, Shilabad, Russia. Paleogene; Chlamys suzakiensis sp. nov. p. 72 pl. 1 f. 8-11 Tadzhikistan depression, Khochilor, Aruk-tan, Tutkaul, Russia. Paleogene; C. schira-badica, sp. nov. p. 74 pl. 1 f. 12–16 Tadzhikistan depression, Shirabad, Russia. Paleogene; C. sokhiensis sp. nov. p. 83 pl. 2 f. 10-17 Fergana; Isfarc, Sokh, Selrokho. Paleogene; C. dichoticosta sp. nov. p. 90 pl. 3 f. 14, 16 Fergane, Kara-Darya. Paleogene, Korobkov, I. A. Trud. Leningrad Soc. Nat. 68 2 1951. C. subvacaensis p. 190 pl. 3 f. 1; C. wayensis p. 189 pl. 2 f. 15; pl. 3 f. 2, 3 spp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro; C. biarritzensis barbieri var. nov. Oligocene, Belloua, Basse-Kabylie, Algeria, P. Muraour p. 318 pl. 3 f. 1, 2; C. (Radulopecten) drewtonensis sp. nov. Callovian, Jurassic. Kellaways Rock, E. Yorkshire, England, J. Neale, p. 371, pl. 28; C. kannogawaensis sp. nov. Tertiary. Tanzawa massif, Kanagawa Prefecture, Japan, M. Shibata p. 231 pl. 33 f. 4 a, b.; C. (Gigantopecten) latissima from the Neogen of Moravia, V. Zázvorka.

Coralichlamys acroporicola from coral at Queensland, Australia, L. Woolacott.

Cyclopecten powelli p. 23 f. 34, 35, 610 metres, Galathea Station 626, Tasman Sea; C. mestayerae p. 24 f. 36, 39, 75 fathoms, off North Cape, New Zealand spp. nov., R. K. Dell.

†Harpax note on ecology, V. Bodui-levskii.

†Lima (Acesta) omorii sp. nov. Kadonosawa formation, Miocene. Shiratori river, Iwate Prefecture, Japan, S. Aoki p. 189 pl. 29 f. 1; L. (Plagiostoma?) kuromagariensis p. 179 pl. 6 f. 18a-b; L. naumanni p. 177 pl. 6 f. 13-15 spp. nov.; L. naumanni obliqua p. 178 pl. 6 f. 16, 17 var. nov. Upper Triassic, Sakawa Basin, Shikoku Japan, Kobayashi & Ichikawa (1); L. (Mantellum) difficilis sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 106 pl. 6 f. 6 text-f. 50; L. acutecostata sp. nov. Rhaetian Triassic. Koni Peninsula, Okhotsk Sea, Russia, I. Tuchkov p. 192 pl. 3 f. 10.

†Limatula rathodi p. 86 pl. 9 f. 4; L. onithali p. 86 pl. 10 f. 14; spp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956.

†Limea? almelae p. 37 pl. 5 f. 4; L.? riosi p. 38 pl. 5 f. 5; spp. nov. Jurassic Maranchón, Castell, Concha & Revilla.

†Neithea coxi sp. nov. Cretaceous. Moghara Massif, Sinai Egypt, I. G. Mahmoud p. 105 pl. 6 f. 2-5.

Parvanussium maorium sp. nov. off E. Otago coast, New Zealand, 260-350 fathoms, R. K. Dell p. 20 f. 30, 31.

†Patinopecten matumoriensis (Nakamura) originally described as a subsp. of P. kimurai (Yokohama) by M. Nakamura, 1940, now raised to specific rank, K. Masuda (2) p. 221, 223 pl. 31.

Pecten maximus general biology and culture, H. A. Cole in M. Graham p. 175-177; P. maximus populations at Brest, France, L. Faure; P. jacobeus chemical content, M. López-Benito; P. yessoensis habitats and spot mortality in Mutsu Bay, Japan, G. Yamamoto; P. (Chlamys) terumachii sp. nov. off Naha City, Ryakyen Is., 2-5 fathoms, J. H. Webb Venus 19 1 1956 22 text-figs.

†Pecten rochi sp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956 p. 75 pl. 9 f 1, 2; P. (Chlamys) alternans sp. nov. Eccene. Northern Aral, Russia, A. K. Alexeiev p. 26 pl. 1 f. 11-13; P. pseudo-pandorae; P. olisiponensis spp. nov. [nom. nud. in list p. 37.] Miocene, Lisbon, J. C. Berkeley-Cotter; P. mancus sp. nov. p. 94 pl. 3 f. 22-27 South Fergane, Shurab, Russia. Paleogene, Korobkov, I. A. Trud. Leningrad Soc. Nat. 68 2 1951; P. karadarjense sp. nov., var. dividicosta var. nov. Tadzhikistan depression Aruk-tan. Russia Paleogene, Korobkov, I. A. Trud. Leningrad Soc. Nat. 68 2 1951 p. 89 pl. 3 f. 10, 12, 13, 15; P. subitus sp. nov. Paleogene. Aruk-Tan, Russia. Tadzhikistan Depression p. 85 pl. 3 f. 5-7, 9, 17; P. kardarjense sp. nov. p. 86 pl. 3 f. 3, 4, 8, 11, 18 Fergana: Kara-Darya. Paleogene, Korobkov, I. A. Trud. Leningrad Soc. Nat. 68 2 1951; P. koniensis sp. nov. Rhaetian, Triassic. Koni Peninsula, Okhotsk Sea, Russia, I. Tuchkov p. 195 pl. 4 f. 4-7; P. (Chlamys) kaneharai from the Miocene of Sendai, Japan, K. Masuda (1); P. (Chlamys) miyatokoensis from the Miocene of Sendai, Japan, K. Masuda (3).

128

Pectinacea from the Tertiary of East Africa, Persia, and the Mediterranean region, Eames & Cox.

†Plagiostoma? polita sp. nov. Permian East Greenland, N. D. Newell Medd. Grønland 110 4 1955 pp. 16 23 pl. 4 f. 1–3.

†*Plicatula kantilaljii* sp. nov. Callovo-Oxfordian Kutch, India, **Agrawal** (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956 p. 83 pl. 9 f. 10 a-c.

†Pseudamussium ignotum sp. nov. Aruk-Tan, Russia. Tadzhikistan Depression, Paleogene, I. A. Korobkov p. 89 pl. 3 f. 19.

†Spondylopecten rogeri sp. nov. Jurassic, Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 p. 73 pl. 8 f. 4a-b 5-7.

†Streblochondria? maynci sp. nov. Permian East Greenland, N. D. Newell Medd. Grønland 110 4 1955 pp. 17 20 pl. 1 f. 15, 16.

†Tosapecten gen. nov. p. 166, Pectinidae, genotype Pecten (Velopecten) suzukii Kobayashi, 1931; T. suzukii paucicostatus p. 169 pl. 5 f. 16; T. suzukii

inflatus p. 169 p. 5 f. 17-18; T. pseudohiemalis mabarus p. 170 pl. 5 f. 20 vars. nov.; T. pseudohiemalis p. 169 pl. 5 f. 19 sp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa.

†Velata hartzi sp. nov. Lias, Jurassic. E. Greenland, A. Rosenkrantz p. 79–83 f. 1–7; "Velata" infrequens p. 163 pl. 5 f. 21; "V." sumeriensis p. 164 pl. 5 f. 22, 23 spp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobavashi & Ichikawa.

### ANOMIACEA

Anomia ephippium note on mimiery L. Acolat (1); Anomia sol Reeve, new to Seychelles, Rost & T. Soot-Ryen; A. cytaeum anatomy, J. Tanaka.

†Anomia fossil byssus from the Miocene. Germany, A. Papp (2); A. preolmstedi sp. nov. Cretaceous. Alabama and Georgia, U.S.A., L. Stephenson (1) p. 242 pl. 41 f. 9-11.

### **OSTREACEA**

†Alectryossia haidingeriana (Munich 1853), notes, H. Hölden (1).

Crassostrea virginica population variations in Long Island Sound, U.S.A., W. R. Coe; C. virginica spawning and egg production, Davis & Chanley; C. virginica osmotic behaviour, Fingerman & Fairbanks; C. margaritacea culture in South Africa, P. Korringa (1); C. virginica relative intensity of setting in Long Island Sound, U.S.A., Loosanoff & Nomeiko; C. virginica influence of spermatic hormones on spawning and water flow, Nelson & Allison; C. virginica nature of the pigment cells and relation of their numbers to fungus disease, Stein & Mackin.

†Exogyra alexandri sp. nov. Jurassic. Maranchón, Castell, Concha & Revilla; E. trivialis sp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro p. 185 pl. 1 f. 2, 7; E. complicata sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 111 pl. 7 f. 1-10; pl. 8 f. 1-12; text-f. 52-54.

Gryphaea angulata, G. virginica general biology and culture, H. A. Cole in M. Graham p. 148-162; G. angulata chemical composition and nutritional value, Graça. Almeida & Costa; G. cytochemical studies on the eggs, H. Kobayashi; G. angulata breeding habits,

H. Vilela; G. angulata stage of sexual maturity in forms from the river Tagus, H. Vilela (2); G. angulata culture in Portugal, H. Vilela (1).

†Liogryphaea oblonga p. 191 pl.-text ii f. 1, 2; L. ampla p. 194 pl.-text ii f. 3, 4; spp. nov. Jurassic E. of Paris Basin, Charles & Maubeuge (1) Bull. Ser. géol. Fr. 6 2 4-6 1953; Liogryphaea revision of the species in the museum at Luxembourg, Charles & Maubeuge.

Ostrea edulis growth in rivers of north-west Spain Andreu & Arté; Ostrea cultivation, biology and note on the systematics, of Philippine species, Blanco, Villaluz & Montalban; O. virginica persistant activity cycles, F. A. Brown, Ralph & others; O. edulis general biology and culture, H. A. Cole in M. Graham p. 148-162; O. edulis control of pests and diseases, Cole & Hancock; Ostrea bactereo-logical control, A. Grijns; O. gigas guanine-deaminatian and nitrogen metabolism, S. Ishida; O. gigas artificial breeding in tanks, Imai, Hatanaka, et. al.; O. gigas distribution of uricase activity, Ishida, Tsuzuki & Nitahara; O. edulis experimental breeding in Japanese waters, Imai, Sakai & Okada; O. denselamellosa cytochemical studies on the eggs, H. Kobayashi; Ostrea cultivation in South Africa, P. Korringa (1); O. edulis mechanism of food movement in the gut of the Larva, R. H. Millar; O. (Crassostrea) madrasensis seasonal gonad changes, K. V. Rao; Ostrea effect of salt, vinegar and heat on the coliforms, E. Salafranca; O. edulis method for seed-oyster collecting, M. Thiéblemont; O. edulis reproduction in the region of Marennes-Oléron, France, P. Trochon; O. gigas content and distribution of cobalt, Tsuchiya & Makuta; O. gigas effects of 1A and fluoride on ciliary activity of gills, I. Usuki; O. gigas effects of cyanide and azide on ciliary activity of the gill, I. Usuki (1); O. gigas effect of malonate and 2, 4-dinitrophenol on cilia of Ostrea gigas gills, I. Usuki (2); O. gigas glycolytic intermediates in the gills, Usuki & Okamura; Ostrea: the effect on spatfall, of controlling barnacle settlement, Waugh & Ansell; O. edulis breeding experiments, P. Walne

†Ostrea konbo Mabuchi (MS) from the Cretaceous of Hokkaido, Japan, I. & S. Hayasaka; O. pharaonis sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 108 pl. 6, 7-12, text-f. 51; O. edulis quantitative aspects of filter feeding, C. B. Jørgensen; O. (Lopha) ucheensis sp. nov. Cretaceous. Russell & Dallas Counties, Alabama, U.S.A., L. Stephenson (1) p. 238 pl. 43 f. 1-5.

†Pycnodonta magnei sp. nov. Cretaceous, Algeria, p. 244 pl. 1, S. Freneix in Freneix & Magné.

### TRIGONIACEA

†Myophorella (Promyophorella) obsoleta p. 2 pl. 1 f. 2, Honmura, Kochi Prefecture; M. (P.) imamurai p. 3 pl. 1 f. 3 Kiritani, Toyama Prefecture; M. (P.?) hashimotoi p. 3 pl. 1 f. 4-7, Kurisaka, Tokushima Prefecture; M. (Haidaia) gracilenta p. 4 pl. 1 f. 8, Arinoki, Kochi Prefecture spp. nov. Jurassic, Japan, T. Kobayashi (2).

†Myophoria nakajimensis p. 180 pl. 6 f. 4; M. okunominetaniensis p. 181 pl. 6 f. 7-9; M. umenokiensis p. 183 pl. 6 f. 6; M. dieneri p. 181 pl. 6 f. 1 spp. nov.; M. dieneri longa p. 181 pl. 6 f. 3, var. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa (1).

†Neoschizodus (Okunominetania) kawarensis sp. nov. Triassic. Kyoto Prefecture, Japan, K. Nakazawa p. 245 pl. 4 f. 11-15.

†Schizodus kennedyensis sp. nov. Lower Permian Western Australia, J. M. Dickins Bull. 29 Com. Aust. dep. Nat. development Misc. res. geol. Geoph. 1956 p. 30 pl. v f. 1–10.

†Setotrigonia subgen. nov. of Steinmanella Cretaceous. Japan, Kobayashi & Amano p. 206.

†Steinmannella (Setotrigonia) subgen.
nov. p. 206. subgenotype: S. (S.)
shinoharai sp. nov. p. 207 pl. 15 f. 1-3;
S. (Yeharella) subgen. nov. p. 200.
subgenotype: Trigonia japonica Yehara:
S. (Y.) jimboi p. 204 pl. 13 f. 4; S. (Y.)
lymani p. 203 pl. 13 f. 3 spp. nov.; S. (Y.)
japonica obsoleta var. nov. p. 202; p. 14 f.
4, 5. Cretaceous. Japan, Kobayashi &
Amano.

Trigonia note on fossil and recent forms, D. McMichael (2).

†Trigonia (Trigonia) elongata var. mebhaensis p. 97 pl. 10 f. 2a, b; T. (T.) e. habhoensis p. 97 pl. 10 f. 4 9a-b, 10; vars. nov.; Trigonia (Indotrigonia) katrolensis sp. nov. p. 98 (see Bull. Soc. Géol. Fr. (6) 6 1-3), Jurassic, Kutch, India, Agrawal (1) Ann. Cent. Et Doc. Paléont. No. 19 1956; T. (Indotrigonia) katrolensis sp. nov. Kimmeridgion, Jurassic, Katrol Series, Cutch, India, S. Agrawal (2) p. 15 pl. 1 f. 2, 3a-b; Trigonia spp. in a new Albian deposit in the Cerbier's orientales, figs., Casteras & Villatti; T. pocilliformis from Mesozoic sandstone, Hokkaido, Japan, S. Imanishi: T. (Minetrigonia) katayamai sp. nov. Upper Triassic, Sakawa Basin, Shikoku, Japan, Kobayashi & Ichikawa (1) p. 184; T. (Vaugonia) from the Cayetano Formation, Jurassic, of Cuba, K. Krömmelbein; T. douvillei nom. nov. for T. orientalis H. Douvillé 1916, I. G. Mahmoud p. 120; T. nalivkini sp. nov. Rhaetian, Triassic. Koni Peninsula, Okhotsk Sea, Russia, I. Tuchkov p. 185 pl. 2 f. 1.

†Trigonioides dentition, hinge-structure and relation to similar pelecypods, T. Kobayashi (3).

†Vaugonia yambarensis sp. nov. Jurassic, Tetori series. Fukui Prefecture, Japan, T. Kobayashi (2) p. 1 pl. 1 f. 1a, b.

† Yeharella subgen. nov. of Steinmanella Cretaceous. Japan, Kobayashi & Amano p. 200.

### UNIONACEA

Anodonta Lamarck 1799, gender feminine, Direction 42; A. minima Millet in the Corfe River, Dorset, A. G. Davis (4); A. minima Millet in Dorset, A. G. Davis (5); A. cygnea effect of 3, 4-benzopyrene, L. Haranghy; A piscinalis and A. cygnea anatomy of the gills, R. Kilias; A. muscle structure of heart, Kirpichnikova; A. cygnea The rate of urine production, W. T. W. Potts; A. cygnea Osmotic and Ionic regulation, W. T. W. Potts (1).

†Anodonta getica sp. nov. Pliocene, Oltenia, Roumania, Rosculescu-Soverth p. 205, 206 pl. 1 f. 1-4; A. (Haasiella) saidovi sp. nov. Triassic, Dzhunger depression, Russia, M. N. Saidov p. 96.

†Anthracosia (Carbonicola) carbonaria A. cf. thuringensis from the inferior Permian of Roumania, G. Răileanu. Caelatura hauttecoueri ruellani. Bgt. fig. digestion and production of H<sub>2</sub>SO<sub>4</sub>, G. R. Fisk.

†Carbonicola artifex p. 45 pl. 1 f. vii, Text-fig. 2g; C. declinata p. 48 pl. 1 f. x text-fig. 2a; C. torus p. 54 pl. 1 f. xi-xiii text-fig. 2f; spp. nov. Carboniferous Pennine District, R. M. C. Eager (1).

†Cuneopsis sinensis, sp. nov. Triassic, Dzhungen depression, Russia, p. 94; C. lacustris sp. nov. Triassic Dzhunger depression Russia, p. 95, Saidov, M. N.

Elliptio memichaeli sp. nov. (E. fraternus auctt., in part. Non Lea, 1852). Choctawhatchee River, west of Miller Cross Roads, Florida, U.S.A., Clench & Turner p. 170 pl. 7 f. 1, 2.

Fusconaia escambia sp. nov. Escambia river, near Century, Florida, U.S.A., Clench & Turner p. 152 pl. 7 f. 3, 4.

Haasodonta gen. nov. Unionidae subfam. Rectidentinae, genotype Hyridella fannyae Johnson 1948, Nautilus 62 47-48, D. F. McMichael Nautilus 70 2 Oct. 1956 p. 47 pl. 2 f. 3-5.

Lamellidens corrianus intracellular digestion of proteins, D. Kamat; L. corrianus phosphateses in digestive diverticula and amoebocytes, D. Kamat (1).

Leiovirgus guppyi aipianus subsp. nov. Aipiana on S. Joseph's River (= Angabunga River) New Guinea, D. F. McMichael Nautilus Philad. 70 2 Oct. 1956 p. 44 pl. 2 f. 1, 2.

Margaritana auricularia and M. margaritifera prehistoric finds in Saxony, R. Hertel.

Margaritifera problem of nomenclature, D. McMichael (1); M. vulgaris chemical investigations on pearl formation, Venkataraman & Chari.

†Megalodon anmandykanensis sp. nov. Vilighi, Okhotsk Sea, Russia. Rhoetian, Triassic, I. Tuchkov p. 187 pl. 2 f. 5.

Naiades, list of types in the Mus. Comp. Zool. Harvard, R. Johnson.

†Palaeanodonta? carinata sp. nov. Tatarian stage, Triassic? Sukhona River, Russia, M. Plotnikov (1) p. 95 pl. 10 f. 10. †Palaeomutela? inconcinna sp. nov. Tatarian stage, Triassic? Sukhona river, Russia, M. Plotnikov (1) p. 91 pl. 10 f. 1, 2, 4, 5.

†Plicatounio radieri sp. nov. Cretaceous Gao, French Sudan, J. Goumard p. 253 text-f. 1-3.

Potamida corrugata alkaline phosphatase activity in the gills, C. Nambudri.

Unio Philipsson 1788, proposed validation, A. E. Ellis; U. pictorum and U. tumidus effect of 3, 4-benzopyrene, L. Haranghy; U. Philipsson, 1788, support for proposed validation, H. Watson.

†Unio uralensis p. 162 pl. 1 f. 1; U. riphaeicus p. 162 pl. 1 f. 2 spp. nov. Triassic. Eastern slope of the Urals, B. Chernuishev; U. biarmicus; U. tertius; U. carino-plicatus; U. grigorovici; U. serrato-radiatus spp. nov. Quaternary lower Kama river basin, Russia, G. Goretzki p. 34; U. wetzleriformis sp. nov. Miocene (Sarmatian) to Pliocene. Tebea, Brad-Baia Basin, Rumania, N. Macarovici; U. wetzleriflabellatiformis from the Pliocene of the Salaj basin, Roumania, M. Paucă; U. horgossica; U. simulator; U. bunbensis spp. nov. Triassic, Dzhungar depression, Russia, M. N. Saidov p. 94.

Unionidae of B. H. and S. H. Wright, W. J. Clench (3); Unionidae of New Guinea, notes and figs., D. F. McMichael.

Velesunio sentaniensis (Haas), V. ovatus (Haas), V. wilsonii (Lea), (new combinations) Unionidae of New Guinea, D. F. McMichael.

Westralunio flyensis Tap-Can, New Guinea (new combination), D. F. McMichael.

#### ASTARTACEA

†Anthonya apicalis shishijimensis subsp. nov. Cretaceous, Shishi-jima and Miyanohara, southwest Japan, M. Amano p. 75 pl. 2 f. 6-12.

†Astarte (Neocrassina) rajnathi sp. nov. p. 433 Pl. 21 f. 1-4; A. (Pruvostiella) subgen. nov. p. 435 genotype: A. (P.) freneixae sp. nov. p. 436 pl. 21 f. 5, 6; Jurassic, Cutch, N.W. India, S. Agrawal; A. notica sp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro p. 197 pl. 3 f. 7; A. choiensis sp. nov. Cretaceous. Sakhalin Island, Siberia, E. Liverovskaya p. 97 pl. 13

f. 9, 10; A. (Nicaniella) extensa badiensis var. nov. p. 106 pl. 10 f. 11 Callovian S. of Badi; A. (Asatarte) pandei p. 102 pl. 10 f. 8, Callovo-Oxfordian; A. (Pruvostiella) francizae p. 108; A. (Neocrassina) rajnathi p. 104 Callovo-Oxfordian spp. nov. of the Jurassic, Kutch, India, S. K. Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956.

†Crassatella chilena sp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro p. 199 pl. 2 f. 2; C. fruscogorensis sp. nov. [nom. nud?] in list p. 172, M. Pasić (1) Zborn. Radova Geol. Inst. Beograd. 4 1952.

†Pacificor subgen. nov. of Venericardia subgenotype Venericardia mulleri Verategui sp. nov. Lodo formation Paleocene, California, P. Verastegui Palaeontogr. Amer. 3 25 1953 pp. 17, 20 pl. 1 f. 6-9.

†Pleurophorus siglanensis sp. nov. Rhaetian, Triassic. Koni Peninsula, Okhotsk Sea, Russia, I. Tuchkov p. 201 pl. 5 f. 6.

†Pruvostiella subgen. nov. of Astarte Jurassic. Cutch, India, S. Agrawal p. 435; Pruvostiella subgen. nov. of Astarte for Astarte (Pruvostiella) freneixae sp. nov. [q.v.], S. K. Agrawal (1) Ann. Cent. ét Doc. Paléont., No. 19 1956 p. 108.

†Venericardia (Pacificor) [subgen; nov. q.v.] mulleri p. 20 pl. 1 f. 6-9. · V. (P.) nelsoni p. 21 pl. 2 f. 1, 5; pl. 3 f. 5; V. (P.) susamaensis p. 22 pl. 5 f. 1–4; V. (P.) durhami p. 23 pl. 7 f. 1, 2; V. (P.) hertleini p. 24 pl. 15 f. 1, 2, 6; V. (P.) orcgonensis p. 25 pl. 9 f. 7-9; V. (P.) aegentea p. 25 pl. 1 f. 10-14; V. (P.) diabloensis p. 27 pl. 5 f. 5-7, 7a; V. (P.) gabbi p. 30 pl. 19 f. 1, 5, 6; V. (P.) weaveri p. 31 pl. 21 f. 3, 4; V. (P.) transversaria p. 37 pl. 2 f. 2-4; V. (P.) taliaferroi p. 38 pl. 1 f. 15, 16; V. (P.) lisa p. 39 pl. 21 f. 1, 2; V. (Glyptoactis) keenae p. 41 pl. 1 f. 1-5; V. (G.) memastersi p. 42 pl. 13 f. 2, 3; V. (G.) domenginica p. 43 pl. 13 f. 1; V. (G.) marksi p. 44 pl. 19 f. 2-4; V. (Venericor) simiana p. 47 pl. 4 f. 1-4; V. (Leuroactis) schencki p. 50 pl. 4 f. 6-8; V. (L.) alisoensis p. 52 pl. 10 f. 1-3; V. (L.) vokesi p. 61 pl. 14 f. 1-3; V. (L.) popenoci p. 62 pl. 22 f. 1-3; spp. nov. Paleocene and Eocene of Peru, P. Verástegui Paleontogr. Amer. 3 25 1953.

### CARDITACEA

Pleuromeris ultima sp. nov. 130 fathoms, off the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 28 f. 23-25.

#### SPHAERIACEA

Corbicula variation due to environment, figs., T. Hayashi.

†Cyrena microvenulina; C. karamaica spp. nov. Cretaceous, Dzhungar depression, Russia, M. N. Saidov p. 92.

Pisidium obtusale (Lmk.) occurring in Craters and Gun pits on Bookham Common, C. F. J. Benslay; P. pseudosphaerium Favre and Iford, Sussex, G. P. Dance (1); P. tenuilineatum; P. hibernicum; P. pseudosphaerium; P. lilljeborgi species from Czechoslovakia, V. Ložek (3).

†Pisidium obtusale lapponicum Clessin from Swanscombe, Kent, S. P. Dance; P. vinecutianum and notes of Pleistocene occurrence with associated species, H. Zeissler Arch. Molluskenk. Frankfurt a.M. 85 4/6 p. 197.

Sphaerium Scopoli, 1777, neuter gender, Direction 42.

Sphaerium corneum meiosis, H.-G. Kevl.

†Sphaerium pseudolacustre sp. nov, Triassic, Dzhunger depression, Russia. Saidov, M. N. p. 94.

#### CYPRINACEA

†Anisocardia (Collignonicardia) subgen. nov.; A. (C.) simplex sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud.

†Collignonicardia sub-gen. nov. of Anisocardia Cretaceous: subgenotype simplex sp. nov. [see supra]. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 121.

Cyprina islandica neurohormones, J. Welsh (1).

†Cyprina moreti sp. nov. p. 120 pl. 9 f. 26-30; C. moreti orientalis var. nov. p. 121 pl. 10 f. 1-4 Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud; C. islandica from the Quarternary of Sicily, A. Malatesta.

†Vesicomya (Waisiuconcha) margotae sp. nov. Mio Pliocene, Waisioe, Baton, Malay Arch., C. Beets Laid. geol. Meded. 17 1953 p. 255 pl. 1 f. 11-15.

### DREISSENACEA

Dreissena polymorpha effect of 3, 4-benzopyrene on metabolism, L. Haranghy; D. polymorpha protozoan parasites of specimens from the Firth of Vistula, Poland, Z. Raabe.

†Dreissena Kolesnikovi sp. nov. Lower Kama basin, Quaternary Russia, Goretzki, G. I. p. 33.

†Congeria ramphophora vösendorfensis p. 159 pl. 13 f. 26-28; C. partschi leobersdorfensis p. 172 pl. 16 f. 8, 9; C. partschi globosatesta p. 172 pl. 16 f. 3; C. partschi firmocarinata p. 174 pl. 16 f. 4; C. subglobosa longitesta p. 174 pl. 16 f. 4; C. subglobosa longitesta p. 177 pl. 19 f. 2; subspp. nov., A. Papp Mitt. Geol. Ges. Wien. 44 1953; C. praeornithopsis p. 167 pl. 15 f. 1, 2; sp. nov., A. Papp Mitt. Geol. Ges. Wien. 44, 1953 C. subdigitifera sp. nov. nom. nud. Miocene. Paša Bunar Valley, Yugoslavia, P. Stevanović p. 27.

#### LUCINACEA

Anodontia edentula (L.) new to Seychelles, Rost & T. Soot-Ryen.

Codahia (Lentellarcia) interrupta Lmk. new to Seychelles, Rost & T. Soot-Rven.

†Dentilucina doderleini from the Pliocene of Sicily, A. Malatesta.

Jugonia (Epicodakia) divergens Phil. new to Seychelles, Rost & T. Soot-Ryen.

†Lucina subporrecta sp. nov. Cretaceous Antofagasta, Chile, Leanza & Castellaro p. 200 pl. 2 f. 6-9.

†Miltha kolesnikovi sp. nov. Miocene. Turkmenstan and W. Kazakhstan, Merklin & Nevasskaya p. 73 pl. 21 f. 4-7.

Mytiloides anabarensis sp. nov. Anabar region, Russia Turassic, Voronetz, N. S. & Laptinskaya, E. S. (1) C.R. Acad. Sci. Moscou 100 5 1955 956.

†Sphaera rogeri sp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Palcont. No. 19 1956 p. 111 pl. 11 f. 3-5.

†Sphaeriola pruvosti sp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956 p. 114 pl. 10 f. 6. †Tancredia gigantea sp. nov. Anabar region. Jurassic, Russia, Voronetz, N. S. & Laptinskaya, E. S. (1) C. R. Acad. Sci. Moscou 100 5 1955 p. 956.

#### ERYCINACEA

†Cocculinella salisburyensis sp. nov. Pliocene, Australia, N. H. Ludbrook p. 26 pl. 2 f. 1 [nom. nud. Tr. roy. Soc. S. Aust. 77 1954 p. 58.]

Coriareus neozelanicus sp. nov. 330 fathoms, S.E. of Pitt Island, New Zealand, R. K. Dell p. 30 f. 42, 43, 47.

Lasaea rubra filtering, feeding and digestion, Ballantine & Morton.

Notolepton triangulare sp. nov. 120–125 fathoms, N.E. of Kaingaroa, Chatham Islands, New Zealand, R. K. Dell p. 30 f. 41.

Rochefortula quadrata sp. nov. 130 fathoms, off the Forty Fours, Chatham Islands, New Zealand, R. K. Dell p. 29 f. 44, 46.

Scintillona benthicola sp. nov. 330 fathoms, S.E. of Pitt Island, Chatham Islands, New Zealand, R. K. Dell p. 31 f. 40.

#### CHAMACEA

†Agria falconi sp. nov. Cretaceous, Jamaica, L. Chubb p. 13 pl. 2 f. 3; text-f. 3.

†Biradiolites forbesi sp. nov. Cretaceous, Jamaica, L. Chubb p. 15 pl. 3 f. 1-2.

Chama Linn. 1758, proposed addition to the "Official List of Generic Names in Zoology," L. R. Cox; C. limbata Lmk. new to Seychelles, Rost & T. Soot-Ryen.

†Gyropleura shaviensis sp. nov. Cretaceous, Jamaica. Apparent first record of genus from Western Hemisphere, L. Chubb p. 7 pl. 1 f. 4-6.

†Hippurites (Orbignya) ceibarum sp. nov. Cretaceous, Jamaica, L. Chubb p. 19 pl. 4 f. 9, 10 text-f. 4.

†Matheronia semirugata from the Cretaceous of Roumania, Atanasiu & Semaka; M. rougonensis, sp. nov. lower Valaujiniau Grand Canon Verdon, Mongin & Trouve Bull. Soc. Géol. France 6 3 1-3 1953 p. 231 pl. iii f. 5 and text-figs. 3-4.

†Mitrocaprina multicanaliculata sp. nov. Cretaceous, Jamaica, L. Chubb p. 11 pl. 1 f. 14.

†Monopleura jamaicensis sp. nov., Cretaceous, Jamaica, L. Chubb p. 6 pl. 1 f. 1-3.

†Parastroma maldonensis sp. nov. Cretaceous, Jamaica, L. Chubb p. 21 pl. 5 f. 1-6.

†Plagioptychus trechmanni p. 8 textf. 1, 2 (placed by Trechmann in Caprina jamaicensis Whitfield); P. zansi p. 9 pl. 1 f. 7-9; P. minor p. 10 pl. 1 f. 10-13 spp. nov. Cretaceous, Jamaica, L. Chubb.

†Praeradiolites verseyi sp. nov. Cretaceous, Jamaica, L. Chubb p. 14 pl. 2 f. 4-6.

Radiolitidae. List of Cretaceous forms from the North Pyrenees, G. Astre (1).

 $\dagger Rudistes$  early work by Lapeirouse, G. Astre.

†Sauvagesia mcgrathi p. 17 pl. 3 f. 3-4; S. fluminisagni p.18 pl. 4 f. 3, 7, spp. nov. Cretaceous, Jamaica, L. Chubb.

†Thyrastylon gen. nov. p. 36 Radiolitidae, genotype: Radiolites adhaerens Whitfield, 1897. Also includes Biradiolites coryi Trechmann, 1924; B. semiannulosus Trechmann, 1924, from Jamaica and B. persicus Douvillé 1904 from Persia. Cretaceous, L. J. Chubb (1).

### CARDIACEA

†Arcicardium bosphoranum sp. nov. Pliocene. Kerchenski, Crimea Peninsula, Russia, A. Eberzin p. 19 pl. 1 f. 2-4.

†Caladacna steindachneri euxinica subsp. nov. Pliocene, Kerchenski, Crimea Peninsula, Russia, A. Eberzin p. 30 pl. 5 f. 11-14.

Cardium edule factors affecting growth-rate, H. Cole; C. edule general biology and culture, H. A. Cole in M. Graham p. 168-175; C. edule observations on variation, P. Mars; C. edule in Iceland, fig., I. Oskarsson (1).

†Cardium transcaucasicum p. 127 pl. 2 f. 1, 2; C. agdschakendense p. 129 pl. 1 f. 4, 5; Azerbaijan; C. gulistanense p. 131 pl. 2 f. 3, 4; C. dimidiatum p. 132 pl. 2 f. 5-8; Armenia spp. nov. Cretaceous. Caucasus, Russia, N. Bobkova; C. sartasense sp. nov. Miocene. Turkmenstan and W. Kazakhstan, Merklin & Nevesskaya p. 46 pl. 8 f. 14-17; C. (Trachycardium) ochilelanum sp. nov. Cretaceous. Ochillee Creek, Georgia, U.S.A., L. Stephenson (1) p. 242 pl. 41 f. 1, 2.

†Caspia plaekovalevskii sp. nov. Lower kamar basin Quarternairy Russia, Goretzki, G. I. p. 33.

†Caspicardium gen. nov. Cardiidae, Turkmenia, Asserbaydjan, Astagyeva, K. A. Bull. Soc. Nat. Moscou. Geol. 30 3 1955 94.

†Chartoconcha tenuissima sp. nov. Pliocene. Kerchenski, Crimea Peninsula, Russia, A. Eberzin p. 23 pl. 5 f. 5a, b.

†Conocardium howitti sp. nov. Devonian, Buchan Caves limestone, Victoria, Australia, J. A. Talent p. 51, f. 15; C. angelicum sp. nov. p. 67 pl. 7 f. 23-25. Marble Creek, Thomson River, Victoria, Australia; C. cresswelli p. 68 nom. nov. for Pleurorhynchus bellulus Cresswell (non Barrande) 1893, Talent & Philip.

†Corculum (Ctenocardia) weberi sp. nov. Mio-Pliocene Waisioe, Buton Malay Arch., C. Beet Leid. geol. Meded. 17 1953 p. 253 pl. 1 f. 8-10.

†Cypricardinia multilamellosa sp. nov. Devonian, Buchan Caves limestone, Victoria, Australia, J. A. Talent p. 50 pl. 4 f. 7.

† Didacna devexa Salo-Manitcluterrivers Quaternary Russia? p. 35; D. emendate Primanu part of Pricaspian Quaternary Russia p. 35; D. tanaitica Don and Warsaw Manich Quaternary. Russia p. 40; D. parvuloides Premanich part of Pricaspian Quaternary. Russia p. 35, spp. nov. Popov, G. I. (1) Bull. Soc. Nat. Moscou geol. 30 7 1955; D. tanaitica Dono-Manich bay, Russia. Quaternary, p. 144; D. emendata Quaternary, Drevneevski horizon, Russia Quaternary p. 145; D. lissitzini Quaternary, Urundzhiksk horizon, Russia, p. 145; D. sokolovi Quaternary, Drevneevski horizon. Russia p. 145, spp. nov. Popov, G. I. C.R. Acad. Sci. Moscou. 101 1 1955; D. adacnoides p. 78 pl. 1 f. 11–14; pl. 2 f. 9; D. plana p. 78 pl. f. 4-7; p. 2 f. 6; D. porsugelica p. 78 pl. 1 f. 8-10; pl. 2 f. 8 spp. nov. Khazar deposits, Quaternary. Cheleken, Russia, Turkmenia, L. A. Nevesskaya.

Fragum fragum (L.) new to Seychelles, Rost & T. Soot-Ryen.

†Granocardium (Ethmocardium) pomeyroli from New Caledonia, now thought to be Cretaceous, R. Pomeyrol.

Hemicardia nivale Reeve new to Seychelles, Rost & T. Soot-Ryen.

Hippopus hippopus spawning habits, S. Wada.

[1956]

Laevicardium australe Swby. new to Seychelles, Rost & T. Soot-Ryen; L. subdiprosopum sp. nov. nom. nud. Miocene. Yugoslavia, P. Stevanovié p.26; L. praeinflatum p. 193 pl. 22 f. 4-6; L. edlaueri p. 194 pl. 23 f. 7-9; spp. nov. L. ornatum biseptum subsp. nov. p. 196 pl. 22 f. 23-25, A. Papp Mitt. geol. Ges. Wien. 44 1953.

†Liopistha elegantulata nom. nov. for Cardium elegantulum Roemer, 1849 (non Beck, 1842) Austin Chalk, Cretaceous, Texas, H. E. Vokes (1) p. 764.

†Monodacna viennensis p. 200 pl. 22 f. 19-21; M. vösendorfensis p. 201 pl. 13 f. 1-3; spp. nov., A. Papp Mitt. Geol. Ges. Wien. 44 1953; M. euxinica sp. nov. Primarily part of Pricaspian Quaternary Russia 1955, Popov, G. I. (1) p. 35.

†Oraphocardium alatoplanum infracinmericum p. 69 pl. 12 f. 13, 14, Kerchenski, Crimea; O. alatoplanum transcaucasicum p. 69 pl. 12 f. 5-9 Duab, south of Samarkand. subspp. nov. Pliocene Russia, A. Eberzin.

†Oxydacna buguntyrica p. 36 pl. 6 f. 6a, b; O tertiana p. 37 pl. 6, f. 9-12; Kuban Province; O.(?) ambrae p. 39 pl. 7 f. 1, 2 Ambra river spp. nov. Pliocene Russia, A. Eberzin.

†Panticapaea praeduboisi sp. nov. Pliocene. Kerchenski, Crimea Peninsula, and Kuban Russia, A. Eberzin p. 42 pl. 7 f. 3-8.

†Paradacna andrussowi p. 53 pl. 9 f. 18a, b; P. deformis p. 55 pl. 9 f. 12-14 sspp. nov. Kerchenski Crimea and Kuban; P. deformis kubanica subsp. nov. Kuban Pliocene Russia, A. Eberzin.

†Pteradacna pracedentula sp. nov. Kerchenski, Crimea, Russia. Pliocene, A. Eberzin p. 88 pl. 15 f. 1-2.

†Stenodacna praeangusticostata sp. nov. Kuban. Kerchenski, Crimea, Russia. Pliocene, A. Eberzin p. 91 pl. 16 f. 2-7.

Trachyeardium alternatum Swby.; T. leucostomum Boru; T. orbita Swby.; T. pectiniform Born.; T. rubicundum Reeve new to Seychelles, Rost & T. Soot-Ryen.

Tridacna squamosa and T. serrifera, spawning habits, S. Wada; T. deresa Symbiosis, Habitat, Feeding, figs., C. M. Yonge.

### VENERACEA

Callista brevisiphonata as a cause of food poisoning, Asano, Takayamagi & Kitamura.

†Callista chinensis takagii subsp. nov. Miocene Noto Peninsula, Japan, K. Masuda Fr. Proc. Palaeont. Soc. Japan N.S. No. 20 1955 p. 121 pl. 19, f. 8.

†Clementia japonica sp. nov. Miocene Noto Peninsula, Japan, K. Masuda Tr. Proc. Palaeont. Soc. Japan N.S. No. 20 1955 p. 121 pl. 19 f. 8.

Dosinia exolete general note, I. Kristensen.

†Dosinia chikuzenensis Nagao, systematic consideration, p. 109; D. nomurai Otuka, and D. nagaii Otuka, proposal to place these in the synonomy of D. chikuzenensis p. 112, K. Hirayama.

†Dosiniopsis jerofeievi sp. nov. Cretaceous. Sakhalin Island, Siberia, E. Liverovskaya p. 98 pl. 13 f. 7, 8.

Gafrarium aduncum Römer new to Seychelles, Rost & T. Soot-Ryen.

Katelysia (Nipponomaria) [q.v.], J. Itoigawa.

Meretrix meretrix lusoria guanine-deaminating activity and nitrogen metabolism, S. Ishida; M. meretrix lusoria distribution of the uricase activity and aspects of purine metabolism. Ishida & Tsuzuki; M. meretrix lusoria cytochemical studies on the eggs, H. Kobayashi.

Nipponomarcia, should be separated as a genus from Katelysia, for N. nakamurai Ikebe as type, J. Itoigawa Venus 18 3 1955 178, 184.

Petricola pholadiformis functional morphology, R. Purchon.

Pitas (Pitarina) obliquata Hanley new to Seychelles, Rost & T. Soot-Ryen.

†Similivenus insolida nom. nov. for Venus solida Deshayes 1825 non Schroeter 1802 A. M. Keen J. Paleont. 28 2 1954 p. 218.

Tapes litteratus L. one broken ex. only; T. deshagesi H. new to Seychelles, Rost & T. Soot-Ryen; T. semidecussata polymorphism, figs., R. F. Shaw.

†Tapes (Amygdala) miyamurensis sp. nov. Miocene, Kyoto Prefecture, Japan, J. Itoigawa p. 189 pl. 1 f. 2, 3.

Tivela stultorum population fluctuations in California, W. R. Coe.

Venerupis philippinarum distribution of the uricase activity, Ishida & Tsuzuki; V. philippinarum guaninedeamination and nitrogen metabolism, S. Ishida; V. semidecussata cytochemical studies on the eggs, H. Kobayashi; V. semidecussata population density effects, S. Ohba.

Ventricolaria nom. nov. for Ventricola Römer 1867 with type Venus rigida Dillwyn (Caribbean sp.), A. M. Keen J. Paleont. 28 2 1954 p. 218.

Venus mercenaria persistant activity cycles, F. A. Brown, Ralph & others; V. mercenaria spawning and egg production, Davis & Chanley; V. mercenaria acclimatized in Brittany, L. Marteil; V. mercenaria environmental effect on growth, Pratt & Campbell; V. mercenaria effects of acetylcholine on the isolated heart, J. Welsh; V. mercenaria cytochemistry, S. I. Zacks.

†Venus sobieskii; V. sobieskii var. media, from the Miocene of the south west area of the Russian Platform, L. Kudrin.

### MACTRACEA

Atactodea trigona Desh. new to Seychelles, Rost & T. Soot-Ryen.

Caecella chinensis artificial activation of the eggs, N. Sawada.

†Cymbophora ochilleana sp. nov. Cretaceous. Ochillee Creek, Georgia, U.S.A., L. Stephenson (1) p. 244 pl. 44 f. 1-3.

Donacilla seychellarum sp. nov. Seychelles (Mahé) H. Rost and T. Soot-Ryen a single specimen, Acta borealia (A) No. 8 1955 p. 21 text fig. 3 [v. poor).

†Eastonia rugova in ancient rejectamenta of the Rhone Delta, M. Paulus.

Mactra Linn. 1767, gender feminine, Direction 42.

Mactra sorallina cinerea thrown up in quantity after storm, L. W. Grensted; M. sulcataria distribution of uricase activity, Ishida, Tsuzuki & Nitahara: Mactra; experimental activatia of the egg, K. Iwata; M. stultorum mass mortality at Lowestoft, Nov. 21, 1950, H. E. Jenner.

†Mactra vitaliana turkmena subsp. nov. Miocene. Turkmenstan, Merklin & Nevesskaya p. 92 pl. 27 f. 1-6.

Pteropsella nom. nov. for Pteropsis Conrad, 1860 (non Rafinesque, 1814), H. E. Vokes (1) p. 763.

Spisula use of sodium taurocholate as a lysin, J. George.

#### TELLINACEA

†Abra alba euxinica subsp. nov. Miocene. Turkmenstan and Western Kazakhstan, Merklin & Nevesskaya p. 82 pl. 23 f. 1-5.

Donax population fluctuation in California, W. R. Coe; Donax polymorphic colouration, D. Nolte; D. (Latona) cuneata L. D. (L.) faba Gmeh. new to Seychelles, Rost & T. Soot-Ryen.

Jactellina (Loxoglypta) clathrata Desh. = rhomboidea Q. and G. non Gmel. Tellina perrieri Bertin is now Pharaonella perrieri Bertin (syn. is T. consanguinea Sow). T. parvula Bertin is Semelangulus pervula fam. Sunedidae, T. Habe (1).

Leptomya rostrata H. Ad. new to Seychelles, Rost & T. Soot-Ryen.

†Linearia orientalis sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 128 pl. 12 f. 6-8 text-f. 56.

Macoma baltica ecology, N. Miloslavskaya (1).

†Macoma nagaoi sp. nov. Tertiary, south of Watase, Hikoshima, Yamaguchi Prefecture, Japan, K. Hirayama p. 85, p. 115 pl. 8 f. 27.

Psammotaea violacea Lmk. new to Seychelles, Rost & T. Soot-Ryen.

Quadrans gargadia L. new to Seychelles, Rost & T. Soot-Ryen.

Sanguinolaria diphos amino acids of the crystalline style, V. Venugopalan.

Scrobicularia plana mechanism of siphon extension, Chapman & Newell.

Semelangulus kurodai (Makiyama) Japan, is closely allied to S. parvula (Bertin), T. Habe (1).

Tellina serrata Renier "[1804]", specific name, as used in above combination, rejected, Opinion 427.

Tellina perrieri Bertin, T. gibba Bertin, T. parvula Bertin, nomenclature and systematic notes, T. Habe (1).

†Tellina hamadai p. 122 pl. 19 f. 10-11; T. notaensis p. 122 pl. 19 f. 12; spp. nov. Miocene Note Peninsula, Japan, K. Masuda Fr. Proc. Palaeont. Soc. Japan N.S. No. 20 1955.

Thyellisca nom. nov. for Thyella Adams, 1866 (non. Wallengren, 1858), H. E. Vokes (1) p. 762.

#### SOLENACEA

†Cultellus orientalis sp. nov. Eccene. Northern Aral, Russia, A. K. Alexeiev p. 23 pl. 3 f. 1, 4.

Glauconomidae of Japan, figs., Taki & Habe.

Scissuliua dispar Conrad, new to Seychelles, Rost & T. Soot-Ryen.

Solen candidus Renier, "[1804]", the specific name as used in the above combination, rejected, Opinion 427.

Solen siliqua experiments on foot muscle, Bargoni & Atzori.

### SAXICAVACEA

Panope bitruncata Conrad, A living specimen taken at St. Augustine, Florida, M. C. Johnson fig.

#### MYACEA

†Caryocorbula? veatchi p. 245 pl. 44 f. 4-9; C.? georgiana p. 246 pl. 44 f. 14-16 spp. nov.; C.? veatchi longa p. 246 pl. 44 f. 10-13 var. nov. Cretaceous. Georgia and Alabama, U.S.A., L. Stephenson (1).

†Cuneamya holmesi sp. nov. Silurian Pentland Hills nr. Edinburgh, A. Lamont Proc. roy. Soc. Edinb. 65B 3 1954 p. 275 pl. 1 f. 5.

Mya Linn. 1758, gender feminine, Direction 42.

Mya arenaria population dynamics, J. C. Ayers; M. arenaria mechanism of siphon extension, Chapman & Newell; M. arenaria and the natural control of its predator Lunatia heros by Nassarius trivittatus, A. H. Clarke jnr. (1).

Nya pictorum typographical error for Mya pictorum H. Watson Bull. zool. Nom. 12 1956 p. 308.

†Platyodon klinghardti from the Miocene of West Africa, Darteville & Roger; P. nipponica sp. nov. Miocene. Japan, Uozumi & Fujie p. 361-368 pl. 4 f.1-6; pl. 5 f. 4-6.

†Pseudomyalina gen. nov. Myalinidae genotype obliqua sp. nov. Lower Permian, Western Australia, J. M. Dickins Bull. 29 Com. Aust. dep. Nat. development Min. res. Geol. Geoph. 1956 p. 25, 26 pl. iii f. 1–7 text fig. 2.

#### GASTROCHAENACEA

Gastrochaena rostrata from driftwood at Manat, Philippine Islands, G. Ablan (1).

# ADESMACEA

Barnea manilensis collected from driftwood at Manat, Philippine Islands, G. Ablan; (1) B. candida Oology (and technique) and development, J. Pasteels.

Martesia striata note on the biology, Ganapati & Nagabhushanam; M. striata digestive enzymes, Ganapati & Nagabhushanam (1); M. striata water filtration, R. Nagabhushanam (1); M. striata—anatomy and biology, R. Purchon (2).

Pholadidae of Japan, figs., Tahi & Habe.

Pholas dactylus luminescence, J. & J. M. Gaillard.

Teredo Linn. 1758, gender feminine Direction 42.

Teredo navalis shell structure and ecology, P. Deschamps; T. navalis note on its biology, Ganapati & Nagabhushamam; Teredo biological studies, C. E. Lane; T. (Psiloteredo) tondiensis sp. nov. port of Tondi, South India, Nair & Gurumani; T. navalis, methods of control, L. Popescu; T. navalis in coastal waters of Maryland, U.S.A., Scheltema & Truitt; Teredo note on life history and control, F. G. W. Smith; T. (Lyrodus) samoaensis occurence round N. China coast. Morphology, Tchang, Tsi & Li.

Xylophaga washingtona Bartoch and notes on the genus, R. D. Turner.

#### PANDORACEA

†Cercomya peruviana sp. nov. Jurassic Nasca area Peru, L. R. Cox (4) J. Pal. 30 5 1956 p. 1185 pl. 128 f. 7.

Ectorizma neozelanica sp. nov. 300 fathoms, east Otago coast, New Zealand, R. K. Dell p. 43 f. 49.

†Homomya antofagastensis p. 195 pl. 4 f. 1, 6, 7; H. subandina p. 195 pl. 4 f. 3-5; spp. nov. Cretaceous. Antofagasta, Chile, Leanza & Castellaro.

Mactromya mehrotrai sp. nov. Callovo-Oxfordian Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956 p. 116 pl. 11 f. 7a-b.

Pandora inaequivalvis and P. pinna structure and adaptation figs., J. A. Allen; P. albida parasitic castration by Cercaria melanocystea, Arvy & Gaillard.

†Pholadomya japonica sp. nov. Cretaceous, southwest Japan, M. Amano p. 80 pl. 2 f. 1-5.

†Pleuromya harringtoni sp. nov. Cretaceous. Antofagasta, Chile. Leanza & Castellara p. 194 pl. 4 f. 2; P. wakasana sp. nov. Nabae Group, Triassic. Kyoto Prefecture, Japan, K. Nakazawa p. 239 pl. 3 f. 1–3.

†Sanguinolites? inflexum sp. n. Carboniferous Frankenberg, E. Paproth Palëont. Z. 27 3-4 1953 p. 187 pl. 12 f. 8.

#### POROMYACEA

Cuspidaria morelandi p. 39 f. 48, 50; C. morioria p. 40 f. 52, 53 spp. nov. 130 fathoms, east of Forty Fours, Chatham Islands, New Zealand, R. K. Dell.

†"Cuspidaria" ayabensis sp. nov. Nabae Group, Triassic, Kyoto Prefecture, Japan, K. Nakazawa p. 239 pl. 3 f. 1-3

Euciroa delectabile sp. nov. 330 fathoms, S.E. of Pitt Island, Chatham Islands, New Zealand, R. K. Dell p. 42 f. 58, 59.

†Praecardiomya gen. nov. Genotype: P. siniatica [as sinaitica in title] sp. nov. Cenomanian, Cretaceous. Eastern Sinai, Egypt, G. H. Awad (1) p. 7 f. 1-8.

Ptychomya Agassez interpretation of the Genus, A. Chavan.

#### CLAVOGELLACEA

Brechites penis general anatomy, R. Purchon (1).

# **CEPHALOPODA**

RECENT

Cephalopoda. Locomotion of pelagic forms, Anon (1); C. food of Sperm Whales, Akimushkin (1); Cephalopoda of Eastern Seas of Russia, Akimushkin (2): Physiology of nutrition, W. v. Buddenbrock pp. 279–283; Spermiogenesis in some species, A. Franzén (1); Ink ejection in the escape reaction of a squid, D. Hall; On the method of, and changes in coloration, W. Holmes; Deep-sea photographs as a means of learning more about squids, N. Mackintosh; Origin, development and structure of cephalopod eyes, G. da F. Sacarrão (1). Notes on various cephalopods, H. Wendt.

Architeuthis from Japan. Structure of unidentified species, E. Iwai; A. dux specimens from the Trondheim fjord two 1954, and one 1928, fig., E. Sivertsen.

Argonauta Linn. 1758, gender masculine, Direction 42.

Argonauta argo organs associated with optic tract, Boycott & Young (1);

Argonauta notes on the various species, A. H. Jones.

Bathypolypus sponsalis P. and H. Fischer common in the Mediterranean. A study of the species, 12 figs., K. Wirz.

Desmoteuthis hyperborea (Steenstrup) is identical with Teuthowenia megalops (Prosch) which represents a younger stage of D. hyperborea. The species should now be known as D. megalops (Prosch), B. J. Muus p. 7.

Doryteuthis siborgae p. 146 Bay of Sanama, E. coast of Sula Besi f. 16, 18 pl. 1 f. 4; D. pickfordi p. 149 f. 19-21 pl. 1 f. 3 S. coast of Eastern Flores, spp. nov., W. Adam (1) Silboga Exped. Monogr. 35c 1954.

Loligo pealii organs associated with optic tract, Boycott & Young (1); L. pealei blood pigments, E. Eliassen; Loligo cation transport in giant axons, Hodgkin & Keynes.

Lolliguncula abulati sp. nov. Red Sea, "Calypso", W. Adam Ann. Inst. Oceanogr. Monaco N.S. 30 1955 p. 185 pl. 1 f. 1.

Taonius unidentified specimen from Japan, E. Iwai.

Tremoctopus violaceus organs associated with optic tract, Boycott & Young (1); Tremoctopus chromatophore genesis, G. Sacarrão (2).

Eledone moschata blood pigments, E. Eliassen; E. cirrhosa spermiogenesis, A. Franzén (2).

Galiteuthis armata probable first record from Hokkaido coasts, Japan, T. Hikita; Galiteuthis taken from stomach of sperm whale in the North Pacific, E. Iwai.

Macrotritopus problems of classification, W. J. Rees (1).

Nautilus an abnormal shell, fig., J. Sornay; Nautilus undosus J. de C. Swby., fig., J. S. Turner.

Octopus vulgaris reactions to shape and duration of memories, Boycott & Young: O. vulgaris organs associated with optic tract, Boycott & Young (1): O. vulgaris Lmk. A memory system, Boycott & Young (2); O. vulgaris blood pigments, E. Eliassen; O. vulgaris experiment on neuro-sensory cells, figs. Martoja & May; O. vulgaris brachial innervation compared with Sepiola rondeleti, Martoja & May (1); O. vulgaris innervation of arms, Rossi & Graziadei; O. vulgaris note of legends associated with it, W. Tavolga; O. vulgaris tactile sense of blinded animals. M. J. Wells; O. vulgaris regulation of circulation, J. de Wilde; Octopus visual responses before and after training, J. Z. Young.

Ommastrephes pteropus, use of deepsea photography to learn more of squids, N. Mackintosh.

Sepia Linn. 1758, gender feminine, Direction 42.

Sepia officinalis visual perception of movement, P. Boulet; S. officinalis L. P. Boulet (1); S. officinalis organs associated with the optic tract, Boycott & Young (1); S. officinalis blood pigments, E. Eliassen; Sepia cation transport in giant axons, Hodgkin & Keynes; S. officinalis potassium permeability of giant axons, Hodgkin & Keynes (1); S. officinalis washed up in numbers at Lowestoft, H. E. Jenner; S. officinalis experiments on the liver, Jullien, Cardot, Ripplinger & Claudey; S. officinalis chromatophores and colour change, G. Sacarrão; S. officinalis comparison of redox-pigments with ommochrome group of insect pigments, I. Schwinck; S. officinalis early development, H. Wendt.

Sepiola rondeleti brachial innervation compared with Octopus vulgaris, Martoja & May (1).

Teuthowenia megalops (Prosch) is identical with and represents a younger stage of Desmoteuthis hyperborea (Steenstrup). The species should now be known as Desmoteuthis megalops (Prosch), B. J. Muus p. 7.

#### FOSSIL

Spanish Cretaceous Cephalopoda, J. Bataller (1); Cretaceous cephalopods from Muzo emerald mines, Boyaca, Colombia, S.A., H. Bürgl p. 384 pl. 2; Ammonite succession in the Blue Lias of Somerset, England, D. Donovan (1); Cephalopoda of the Pennsylvanian of America, A. G. Unklesbay.

†Acrioceras debelmasi p. 106 pl. 9 f. 7, 8; text-f. 16a; A. ganeshai p. 107 pl. 4 f. 20; text-f. 16c; A. lakhmiae p. 108 pl. 8 f. 10 text-f. 16d; A. maheswariae p. 108 pl. 8 f. 17 text-f. 16e: A. muckleae p. 109 pl. 9 f. 3 text-f. 17a; A. ramkrishnai p. 109 pl. 5 f. 17 pl. 8 f. 12; A. shankariae p. 110 pl. 6 f. 9 text-f. 17c; A. sugrivai p. 111 pl. 8 f. 18 text-f. 17e; A. taraae p. 112 pl. 9 f. 9 text-f. 17f; A. piveteaui p. 116 pl. 8 f. 9 text-f. 18; A. thorali p. 117 pl. 10 f. 2; A. breistrofferi p. 120 pl. 7 f. 3; A. pruvosti p. 123 pl. 9 f. 11; A. spathi p. 124 pl. 7 f. 20 spp. nov.; A. tabarelli sarasini p. 103; A. tabarelli uhligi p. 103; A. seringei multicostata p. 114 pl. 5 f. 16; A. puzosianum drumensis p. 115 pl. 9 f. 4; A. terveri multicostata p. 118 pl. 8 f. 7; A. terveri manteostata p. 118 pl. 7 f. 9; pl. 8 f. 13; A. meriani dilatata p. 122 pl. 8 f. 1; A. meriani tabarelliformis p. 122 pl. 7 f. 21 vars. nov; A. (Paraspinoceras) pulcherrimum latispira var. nov. p. 127 pl. 10 f. 5; A. (Aspinoceras) monmothoi sp. nov. p. 129 pl. 10 f. 3; A. (Protacrioceras) subgen. nov. p. 129; A. (P.) fromageti p. 129 pl. 7 f. 8; A. (P.) oosteri p. 131 pl. 8 f. 15 spp. nov.; A. (P.) ornatum alpina p. 133 pl. 7 f. 13; A. (P.) ornatum flexuosa p. 133 pl. 4 f. 18; A. (P.) ornatum tuberculata p. 133 pl. 7 f. 12 vars. nov. Lower Cretaceous, south east France, S. Sarkar.

†Actinocamax groenlandicus sp. nov. Cretaceous. West Greenland, T. Birkelund p. 5 pl. 1 f. 1-3; Actinocamax microscopical investigation, G. Göke.

†Agathiceras asselicum prima morpha nov. Yurezan river, S. Urals, Russia. Schwagerina horizon, Permian, S. Maksimora p. 26 pl. 3 f. 5–8.

†Amaltheus striatus nom. nov. for Ammonites depressus Simpson, 1843, non Ammonites depressa Bruguière, 1789, M. Howarth p. 161.

Ammonoidea. Literature on Palaeozoic and Mesozic forms from the Pacific area, R. S. Allan p. 378–381.

†Anaptychus bubnoffi sp. nov. Devonian. Thuringia, Germany p. 44 pl. 1 f. 2; A. trauthi nom. nov. p. 44=A. latus Clarke non Woodward 1882; A. bickensis Trauth and A. bickensis curvivexa Trauth 1935=Crickites holzapfeli Wedekind 1913; see p. 36 and 80, K. J. Müller.

†Anarcestina suborder nov. of Ammonoidea, Miller & Furnish J. Paleont. 28 5 1954 p. 687.

†Anaspyroceras cultellus sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 69 pl. 6 f. 7-9.

†Anthracoceras mooreae sp. nov. Carboniferous. Carluke, Scotland, E. D. Currie p. 574 pl. 4 f. 21, 22 text-f. 5x.

†Aphroditiceras subgen, nov. of Salfeldiella. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 80.

†Arcestes kiparisovae sp. nov. Rhaetian, Triassic. Vilighi river, S.E. coast of Okhotsk Sea, Russia, I. Tuchkov p. 203 pl. 5 f. 8.

†Arieticeratinae subfam. nov. of Hildoceratidae, Hyatt. New name for Sequenziceratidae Spath 1924, M. Howarth p. 166.

†Arnioceras arniokridion nom. nov. for Ammonites kridion Quenstedt 1858, non Hehl in Zieten, 1830, O. Walliser p. 207.

†Aspidoceratacea Superfam. nov. Ammonoidea. Aspiroceratidae Zittel, Peltoceratidae Spath and Berriasellidae Spath, Sazonov, N. T. Bull. Soc. Nat. Moscou 30 4 1955 p. 117.

†Asteroceras marstonense at Marston Magna, a note, and its occurrence at Vale of Marshwood, W. D. Lang. †Aturia cf. alabamensis (Morton) in The Eocene of Cyrenaica, figs., A. Haas (1); "Aturia" matsushitai sp. nov. Palaeogene, Kyushu Island, Japan, T. Kobayashi (1) p. 243 text-f. a-d.

†Ammonites listeri Sowerby (J.), 1812: A. sphaericus Sowerby (J.), 1814. These specific names validated, Opinion 420; A. mammillatus Schlotheim, 1813 to be interpreted by reference to the specimen figured by Spath (L.F.) in 1923 as figures 3a and 3b on Pl. 4 of Monograph of the Gault Ammonoidea and to be the type species of the genus, Douvilleiceras de Grossouvre, 1893, Opinion 422; A. cymodoce d'Orbigny, 1850, the nominal species to be interpreted by the lectotype selected in 1896, that is the specimen illustrated by d'Orbigny as f. 3 and 4 on his plate 202, Opinion 426 p. 260; A. royerianus d'Orbigny, 1841, suppression of the specific name; A. cornuelianus d'Orbigny, 1841 (specific name of type species of Cheloniceras Hyatt), the specific name validated, Opinion 428.

†Bactritina suborder nov. of Nautisloidea Miller & Furnish J. Paleont. 28 5 1954 p. 687.

Balatonites semilaevis from the Debarzone of the Pelagoniden (?Triassic) of Macedonia, W. Medwenitsch.

†Balearites labrousseae p. 144 pl. 11 f. 11; B. nolani p. 144 pl. 6 f. 11; B. paramhansi p. 145 pl. 11 f. 6; B. shankariae p. 146 pl. 11 f. 12; B. tuberculatus p. 146 pl. 11 f. 3; B, koechliniformis p. 147 pl. 10 f. 7 spp. nov.; B. balearis unituberculata var. nov. p. 142 pl. 11 f. 5 Lower Cretaceous. south east France, S. Sarkar.

†Belemnitella mucronata [Schloth] mut. senior Nowak in S. Bayern and notes on zone, figs., H. Hagn; Belemnitella abnormalities of structure in two species, S. Koltuipin.

†Belemnites listeri Mantell, 1822, specific name suppressed; B. minimus Miller (J.S.), 1826, specific name validated, Opinion 430; Belemnites microscopical examination, G. Göke; B. (Hibolites) escragnollensis sp. nov. Neocomian maritime Alps, M. Delattre Bull. Sci. géol. Fr. 6 2 4-6 1953 p. 283 pl. xiv f. 1-3; B. cornuclianus note on systematics, V. Stchepinsky.

†Belemnoteuthis rosenkrautzi sp. nov. Cretaceous, West Greenland, T. Birkelund p. 17 pl. 1 f. 9.

†Berbericeratinae subfam. nov. of Macrocephalitidae, to include (with certain reservations) Berbericeras, Schwandorfia and Krumbeckia Jurassic G. Westermann p. 263.

†Beudanticeras revoili elegans var. nov. Cretaceous, Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 98 pl. 5 f. 19, 20.

†Biasaloceras sauclum p. 77 pl. 4 f. 13; B. striatum p. 78 pl. 4 f. 14 spp. nov. Lower Cretaceous. Crimea and North Caucasus, V. Drushchitz.

†Bifoveoceras gen. nov. Nautiloidea genotype: Orthoceras bifoveatum Noetling, 1884. Cambrian, Estonia, Z. Balashov p. 237 pl. 3 f. 1-3.

†Bigotitinae subfam. nov. of Parkinsoniidae; Type; Bigotites haugi (Nicol.) Jurassie, G. Westermann p. 271.

†Bisiphytes (Cenoceras) imlayi Alaska p. 322 pl. 35 f. 1, 2; B. (C.) lupheri Wyoming p. 323 pl. 34 f. 3, 4: spp. nov., Rierdon formation Jurassic, B. Kummel J. Paleont. 28 3 1954 p. 320.

†Burenoceras barnesi sp. nov. Ordovician Texas, Unklesbay (1) J. Paleont. 28 5 1954 pp. 637 641 pl. 68 f. 11-13.

†Cadoceras tschubukulachensis sp. nov., Vorontzev, N. B. & Laptinskaya, E. B. p. 146.

†Cadomitinae subfam. nov. of Stephanoceratidae, Jurassic, G. Westermann p. 251.

†Caicedia subgen. nov. of Pulchellia subgenotype: Ammonites caicedi Karsten, H. Bürgl (2) p. 66.

†Canavaria Gemmellaro, 1886. Designation of type species, p. 167, as Harpoceras (Dumortieria) haugi Gemmellaro, 1885; C. af. cultraroi first record from Britain, Domerian of Yorkshire, p. 168, M. Howarth.

†Ceratites various species from the Trochitenkalk (Miocene) of the Saar, Germany, E. Schneider; C. unusual forms from the Muschelkalk, R. Wenger.

†Cheloniceras Hyatt, 1903 (genotype by designation, Ammonites cornuelianus d'Orbigny, 1841) validated, Opinion 428.

Cheloniceratidae Spath, (L.F.) 1923, to be placed on the "Official List of Family-Group Names in Zoology", Opinion 428.

†Chondroceras (Chondroceras) wrighti minor subsp. nov. p. 61 pl. 3 f. 2-3 Germany; C. (C.) polytomum p. 62 pl. 3 f. 4a-c Spain; C. (C.?) densicostatum p. 65 pl. 3 f. 5, 6 Suiss Jura. spp. nov; C. (Schmidtoceras) subgen. nov. p. 66; subgenotype: C. (S.) schmidti p. 69 pl. 3 f. 7, 8; pl. 4 f. 2–5; pl. 5 f. 1–5 and subsp. p. 71; C. (S.) schindewolfi p. 80 pl. 7 f. 3-7; pl. 8 f. 1-3; C. (S.) arkelli p. 84 pl. 8 f. 4-9; pl. 9 f. 1-3; C. (S). crassum p. 87 pl. 9 f. 4-5; C. (S.) evolutum p. 88 pl. 9 f. 6-9; C. (S.) tenue p. 91 pl. 10 f. 1-4; C. (S.?) antiquum p. 94 pl. 10 f. 7a-d; Germany; C. (S.??) gracile "[n. subgen.]" p. 96 pl. 10 f. 8; pl. 11 f. 1, 2 Swiss Jura; C. (S.) ibericum p. 93 pl. 10 f. 5-6 Spain; C. (Defonticeras?) parvumbilicum p. 104 pl. 12 f. 1, 2 England spp. nov. Bajocian, Jurassic; C. (Schmidtoceras) schmidti multicostatum p. 73 pl. 5 f. 1-5; C. (S.) orbignyanum crassicostatum p. 79 pl. 5 f. 8; pl. 6 f. 3-5; C. (S.) arkelli gerzense p. 86 pl. 8 f. 9; Pl. 9 f. 1-3; Germany; C. (S.) orbignyanum polypleurum p. 77 pl. 6 f. 6; pl. 7 f. 1, 2 Swiss Jura; C. (S.) Schindewolft hispanicum p. 82 pl. 8 f. 1-3 Spain; subspp. nov. Bajocian, Jurassic, G. Westermann (1) Chondroceras (?) (Praetulites) subgen. nov. Type: C. kruizingai sp. nov. holotype "Sphaeroceras gadohense" (Kruizinga, 1926, non Boehm), G. Westermann p. 257.

†Cicatrites hokodzensis sp. nov. Aptian, Cretaceous. North Caucasus, Russia, N. Luppov (1) p. 72 pl. 8 f. 4.

†Cluthoceras gen. nov. p. 560 Goniatitidae: Genotype C. truemani p. 561 pl. 4 f. 1, 2; text-f. 4a, b; C. neilsoni p. 563 pl. 4 f. 3, 4 text-f. 4d-f: spp. nov. Carboniferous. Bishopbriggs, Scotland, E. D. Currie.

†Coeloceras (Préperonoceras) subgen. nov. p. 136 subgenotype C. (P.) brancoi sp. nov. p. 136 a single specimen as Stephanocerus n. sp." Branco descr. & fig. see Branco 1879 Abh. Geol. Spec. Elsass.-Lothringen 2 1 66 pl. 1 f. 5, Jurassic, Lorraine, France, P. Maubeuge.

†Conocerina cloudi sp. nov. Ordovincian Texas, Unklesbay (1) J. Paleont. 28 5 1954 pp. 637, 642 pl. 68 f. 20. Ubirch and Foerste 1935; Denison Univ. Bull. (J. Sci. Labs.) 30 268.

†Coroniceras (Paracoroniceras) charlesi sp. nov. Lias, probably from Stuttgart. Type figured by Reynès, pl. 16 f. 1, 2, D. Donovan p. 12.

†Cotteswoldia spathi sp. nov. (Cotteswoldia sp. Buckman; non Harpoceras opalinum Rein. 1942) Aalenian, Jurassie. Luxembourg, P. Maubeuge (2) p. 78 pl. 2 (see also Arch. Inst. Gr. Duc. Luxemb. N.S. 181 949 pl. xvii); C. angulata p. 367 pl. 1; C. pseudogranjeani p. 369 pl. 2; spp. nov. Jurassic. Luxembourg, P. Maubeuge (3).

†Cravenoceras scoticum p. 575 pl. 2 f. 22-24; text-f. 6a; C. gairense p. 577 pl. 4 f. 8-10 text-f. 6b, c spp. nov. Carboniferous, Scotland, E. D. Currie.

†Creniceras crenatum from the Jurassic of Clucy & Tarcenay, France A. Caire; A. Caire (1); C. crenatum Brug. its occurrence in the Oxfordian at Clucy (Doubs), A. Gaire (3).

†Crimites glomulus sp. nov. Sakmara deposit, Aktyubinsk Province. Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 81 pl. 6 f. 8-10.

†Crioceras baylei p. 36 text-f. 4b; C. krenkeli p. 37 pl. 5 f. 5; text-f. 4c; C. ramkrishnai p. 38 pl. 1 f. 1; text-f. 4d; C. krishnaae p. 39 pl. 1 f. 2; text-f. 5a; C. loryi p. 40 pl. 5 f. 2 text-f. 5b; C. piveteaui p. 41 pl. 4 f. 12 text-f. 5c; C. rogeri p. 42 pl. 4 f. 14 text-f. 6a; C. schindewolfi p. 43 pl. 1 f. 9 text-f. 6b; C. stahleckeri p. 43 pl. 1 f. 5; text-f. 6c; C. viallii p. 44 pl. 4 f. 19 text-f. 6d; C. andersoni p. 46 pl. 4 f. 23 text-f. 7a; C. basseae p. 46 pl. 5 f. 4; text-f. 7b; C. karakaschi p. 48 pl. 4 f. 13 text-f. 8a; C. nowaki p. 49 pl. 4 f. 11 text-f. 8b; C. shibaniae p. 49 pl. 4 f. 9 text-f. 8e; C. sornayi p. 50 pl. 1 f. 7; text-f. 8c; C. sahnii p. 56 pl. 8 f. 16; C. kaliae p. 60 pl. 5 f. 24; C. mahadevai p. 62 pl. 5 f. 7; C. pseudoangulicostatum p. 63 pl. 2 f. 7; C. barrabei p. 64 pl. 5 f. 19; C. inermis p. 64 pl. 3 f. 6 pl. 5 f. 8; C. vishnui p. 68 pl. 3 f. 5; C. arkelli p. 69 pl. 3 f. 9; C. shibaae p. 69 pl. 5 f. 12; C. seitzi p. 70 pl. 3 f. 3; C. anglesensis p. 71 pl. 5 f. 3; C. durgaae p. 7 pl. 5 f. 15; C. gignouxi p. 74 pl. 3 f. 103 C. matsumotoi p. 74 pl. 3 f. 2; "C.'; astieri p. 161 pl. 7 f. 29 spp. nov.; C. duvali sarasini p. 36; C. sornayi densicostata p. 51 pl. 4 f. 8; C. sornayi tuberculata p. 51 pl. 4 f. 15 text.-f. 8d; C. krishnaae tuberculata p. 55 pl. 5 f. 9; C. binelli tuberculata p. 58 pl. 4 f. 2, 4; C. fallauxi tuberculata p. 59 pl. 5 f. 20; C. koechlini recticostata p. 61 pl. 4 f. 7; C. koechlini tuberculata p. 61 pl. 5 f. 10; C. mahadevai tuberculata p. 63 text-f. 9; C. pseudoangulicostatum gracilis p. 63 pl. 2 f. 5; C. villersianum bituberculata p. 67 pl. 3 f. 8; C. villersianum trituberculata p. 67 pl. 5 f. 14 vars. nov. Lower Cretaceous, south east France, S. Sarkar.

Cyclobactrites gen. nov. genotype: Bactrites carinatus Münster Devonian, Germany, Shimanski, V. N. (1) Bull. Soc. Nat. Moscou Geol. 1 1955 p. 97.

†Cymatoceras yabei sp. nov. Nagase formation, Upper Cretaceous, Kôti Prefecture, Japan, Ozaki & Katto p. 63 pls. 11–12 f. 1–4.

Cyrthoceratites depressa Goldfuss, fig., J. S. Turner.

†Daixites attenuatus sp. nov. Sakmara deposit in the Aktyubinsk Province of the Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 49 pl. 1 f. 1, 2.

†Damesites laticarinatus sp. nov. Cenomanian, Cretaceous. Ishikari Province, Hokkaido, Japan, Saito & Matsumoto p. 192 text-f. 1-4.

†Delépinoceras gen. nov. (Demorphoceratidae) genotype Dimorphoceras thabassoide Delépine, Miller & Furnish J. Paleont. 28 5 1954 p. 690.

† Deltocymatoceras gen. nov. Nautilidae: genotype: Nautilus leiotropis Schlüter, 1876. Cretaceous, Germany, B. Kummel (1) p. 438 pl. 22 f. 1, 2.

†Demarezites gen. nov. Cyclolobidae genotype: Waagenoceras oyensi Gerth 1950 Tae Vei deposits, Timor, V. E. Ruzhentzev (1) p. 703.

†Desmoceras difficile in the Argonian and Ancyloceras, P. Gidon.

†Diceras speciosum (Münster) of Württemberg notes. O. F. Geyer.

†Dissimilites piveteaui sp. nov. Lower Cretaceous, south east France, S. Sarkar p. 139 pl. 10 f. 9; pl. 11 f. 7.

†Dorsetensia haydeni sp. nov. Bajocian Tibet, W. J. Arkell (2) Geol. Mag. 90 5 1953 p. 334 pl. xiv f. 9a, b; D. lotharingica p. 168 pl. 10, Ludres. D. excavata p. 169 pl. 11, Ottange. D. thilense p. 172 pl. 12, Thil. spp. nov. Jurassic. France, P. Maubeuge (1).

†Doricranites from the Triassic of Turkmenia, Russia, T. Astakhova.

†Douvilleiceras de Grossouvre, 1893, to be placed on the Official List of Generic Names in Zoology, Opinion 422; D. coronatum Rouchadzé, 1932, (specific name of type species of Megatyloceras Humphrey), the specific name validated, Opinion 428; Douvilleiceras de Grossouvre notice of proposal of Ammonites manmillatus Schlotheim as neotype, F. Hemming; Douvilleiceras sp. An early Cretaceous Ammonite occurring in graywache-type. Sandstone along southern shore of the inlet to San Francisco Bay, Schlocher, Bonilla & Imlay.

Douvilleiceratidae Parona & Bonarelli, 1897 (type genus: Douvilleiceras de Grossouvre, 1893), to be placed on the Official List of Family-Group Names in Zoology, Opinion 422.

†Dufreynoya texana media p. 13 pl. 2 f. 1; pl. 3 f. 1; D. texana sanctorum p. 13 pl. 1 f. 2; pl. 3 f. 5 vars. nov. Aptian Cretaceous. Santander, Colombia, H. Bürgl (1).

†Emericiceras bouladouxae p. 79 pl. 4 f. 8; pl. 9 f. 2; text-f. 12a; E. joliboisi p. 81 pl. 5 f. 11 text-f. 12b; E. journoti p. 82 pl. 5 f. 22 text-f. 12e; E. mahakaliae p. 82 pl. 6 f. 2 text-f. 12d; E. monopujaae p. 83 pl. 6 f. 4; text-f. 12e; E. paramahansai p. 83, pl. 6 f. 1 text-f. 12f; E. villoutreysae p. 84; E. coheni p. 86 pl. 7 f. 2; E. dolloi p. 87 pl. 5 f. 18; E. gerthi p. 87 pl. 4 f. 6, 21; E. foxi p. 90 pl. 7 f. 18; E. imlayi p. 91 pl. 6 f. 17; E. otto haasi p. 95 pl. 6 f. 5; E. paronai p. 97 spp. nov.; E. emerici uhligi p. 77; E. emerici flexuosa p. 77 pl. 7 f. 10; text-f. 11a; E. barremense multicostata p. 85 pl. 6 f. 3; E. barremense recticostata p. 86 text-f. 13; E. clausum flexuosa p. 90 pl. 6 f. 7; E. thiollieri bituberculata p. 93 text-f. 14; E. t. multicostata p. 93 pl. 4 f. 22; pl. 10 f. 6; E. otto haasi tuberculata p. 95 pl. 6 f. 6; E. emerici taraaen p. 96 pl. 7 f. 6 vars. nov. Lower Cretaceous south east France, S. Sarkar.

Encoiloceralidae fam. nov. Nauti. loidea, Shimanski, V. N. & Erlanger, A. A., Bull. Soc. Nat. Moscou Geol-30 3 1955 p. 96.

†Engonoceras wilkinsoni sp. nov. Albian, Cretaceous. Oregon, U.S.A., E. Packard p. 399 text-f. 1. †Eoasianites trapezoidalis sp. nov. p. 11 pl. 1 f. 7-12 text-fig. 6-9; E. subhanieli alta 'morpha nov.' p. 15 pl. 2 f. 4-6. River Yurezan, S. Urals, Russia. Schwagerina horizon, Permian, S. Maksimora.

†Eotetragonites breistrofferi sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 86 pl. 5 f. 10.

†Epicymatoceras gen. nov. Nautilidae genotype: Nautilus vaelsensis Binckhorst 1861. Cretaceous, Germany and Belgium B. Kummel (1) p. 439 pl. 23 f. 1, 2.

†Eremoceras sansabaense sp. nov. Ordovician, Texas, Unklesbay (1) J. Paleont. 28 5 1954 pp. 637, 649 pl. 71 f. 9, 10.

†Eulytoceras rotundum sp. nov. Lower Cretaceous. Crimea and North Caucasus, V. Drushchitz p. 86 pl. 5 f. 15.

†Eutrephoceras montanensis sp. nov. Jurassic Montana, B. Kummel, J. Paleont. 28 3 1954 p. 320 pl. 34 f. 1, 2; E. egitoense sp. nov. Senonian (Cretaceous), Benguela, Angola, Miller & Carpenter p. 34 pl. 1 text-f. 2.

†Ezilloella gen. nov. p. 263 Vascoceratidae. Genotype: E. ezilloensis sp. nov. p. 264 pl. 3 f. 4 text-f. 10. Upper Cretaceous. Aboine river, Nigeria, R. A. Reyment.

†(Fuciniceras)? bajocense sp. nov. Jurassic. France? P. Maubeuge (1) p. 164 pl. 4.

†Garantiana tetragona pseudotetragona p. 174 pl. 14, Domèvre en Haye; G. tetragona renardi p. 175 pl. 15, Maxéville vars. nov. Jurassic, France, P. Maubeuge (1).

†Gastrioceras Hyatt, 1844, genotype Ammonites listeri Sowerby (J.) 1812) validated, Opinion 420.

†Germanites (Germanoides) subgen. nov. Jurassic, G. Westermann, p. 251.

†Germanoides subgen. nov. of Germanites; subgenotype: G. nodosus Westermann, 1954. Jurassic, G. Westermann p. 251.

†Glaphyrites rarus sp. nov. River Yurezan, S. Urals, Russia. Schwagerina horizon, Permian, S. Maksimora p. 10 pl. 2 f. 1-3 text-f. 5.

†Gombeoceras subtenue p. 261 pl. 4 f. 4, text-f. 3f. Bauchi Province; G.? bulbosum p. 263 pl. 4 f. 2, text-f. 3g, 9, Bornu Province, spp. nov. Upper Cretaceous, Nigeria, R. A. Reyment. †Goniatitacea from the Devonian, H. K. Erben.

†Goniatites de Haan, 1825 genotype species: Ammonites sphaericus Sowerby (J.) 1814) to be placed on the Official List of Generic Names in Zoology, Opinion 420; Goniatites and successions of zones in N. Africa, G. Delépine.

Gymnites cf. bosnensis from the Pelagoniden (? Triassic) of Macedonia, W. Medwenitsch.

†Gyroceratites dorsolamellatus sp. nov. Devonian Europe, H. K. Erben Neues Jb. Geol. Paleont. 98 2 1953 p. 185 pl. 17 f. 7-9 text-fig. 3.

†Hammatoceras ferrugineuse sp. nov. Jurassic, Luxembourg, P. Maubeuge (1) p. 154 pl. 4.

†Hecticoceras (Brightia) nodosum recurvum subsp. nov. p. 22 pl. 4 f. 2; H. (B.) tenuinodosum sp. nov. p. 23 pl. 2 f. 12; H. (Zieteniceras) subgen. nov. p. 24, subgenotype: H. zieteni de Tsytovitch 1911; H. (Lunuloceras) michailowense sp. nov. p. 43 pl. 1 f. 10; H. (Rossiensiceras) tsytovitchae sp. nov. p. 50 pl. 2 f. 1; H. (R.) cf. rossiense teisseyrei "n. nom.", subsp. nov. p. 52 pl. 4 f. 25; H. (R.) metomphalum savoiense "n. nom.", subsp. nov. (see pl. 4 f. 30) p. 57, pl. 2 f. 10; H. (Putealiceras) arkelli p. 66 pl. 3 f. 6; H. (P.) schalchi p. 67 pl. 3 f. 4 spp. nov.; H. (P.) krakoviense transiens p. 70 pl. 3 f 5; H. (P.) punctatum arcuatum p. 72 pl. 3 f. 10; H. (P.) punctatum exile p. 73 pl. 3 f. 11 subspp. nov. Callovian, Jurassic. Blumberg, South Baden, Germany, A. Zeiss.

†*Heinzia* (*Heinzia*) colleti sp. nov. Barremian, Cretaceous. Santander, Colombia, **H. Bürgl** (2) p. 80 pl. 23 f. 1-5.

†Hettneria subgen. nov. of Pulchellia subgenotype: P. selecta Gerhardt, H. Bürgl (2) p. 70.

†Holcophylloceratinae subfam. nov. of Phylloceratidae, V. Drushchitz p. 131.

†Homolsomites, Craspeditidae. The range of the genus and species included described under other generic names, R. W. Imlay (1).

†Hoplites Neumayr with Ammonites dentatus Sow 1821 as lectotype proposal for plenary powers to establish the name, M. Breistroffer. "Hoplitoides" adkimsi sp. nov. Tissotiidae Lower Turonian, Mexico, Kummel & Decker J. Paleont. 28 3 1954 pp. 310, 316 pl. 32 f. 6 pl. 33 f. 3 text-figs. 7, 8.

†Hypophylloceras (Hyporbulites) aegyptiacum sp. nov. Cretaceous, Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 84 pl. 5 f. 8, 9.

†Imitoceras bertchogurense sp. nov. Carboniferous. Ber-Choghura, Kirghiz Steppe, N. of Aral Sea, Russia, E. Balashova p. 193 pl. 11 f. 17–25; pl. 12 f. 10.

†Infragarantiana gen. nov. Perisphinctidae; genotype: I. primitiva (Wetzel, 1936) Jurassic, G. Westermann p. 268.

†Infraparkinsonia gen. nov. Perisphinctidae genotype: I. inferior (Bentz 1928) Jurassic, G. Westermann p. 268.

†Jacobella and Hoplites auct. availability of name, M. Breistroffer; Jacobella a propos the name Cenomanaster for Jacobella Mercier 1935, C. W. Wright (1).

†Jaubertites gen. nov. Ancyloceratidae. Genotype: J. dubius nov. (ex. Jaubert in coll.) p. 99 pl. 8 f. 3; J. collignoni sp. nov. p. 99 pl. 7 f. 25; J. dubius tuberculata var. nov. p. 99 pl. 7 f. 11, Lower Cretaceous, south east France, S. Sarkar.

†Jouaniceras numidicum sp. nov. Coniacian Algeria, David & Sornay C.R. Acad. Sci. Paris 242 4 1956 p. 532.

†Juresanites kazakhorum sp. nov. Sakmara deposit. Aktyubinsk Province, Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 68 pl. 3 f. 1-4; pl. 4 f. 1.

†Kamerunoceras gen. nov. p. 250 Collignoniceratidae: Type species. Accuthoceras eschii Solger. Upper Cretaceous Mungo River, Cameroons, Nigeria, R. A. Reyment.

†Kellawaysites basseae sp. nov. Callovian, Kutch, India, Agrawal (1) Ann. Cent. ét Doc. Paléont. No. 19 1956 p. 33 pl. 3 f. 6 text-fig. 9.

†Kinkeliniceras sornayi sp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. ét Doc. Paléont. No. 19 1956 p. 41 pl. 6 f. 2.

†Knemiceras spathi p. 45 pl. 1 f. 1-11; pl. 2 f. 1 text-f. 18-27; K. collignoni p. 60 pl. 3 f. 2, 3 text-f. 31-34; K. rittmanni p. 63 pl. 3 f. 4-13 pl. 4 f. 1 text-f. 35-38; K. deserti p. 68 pl. 4 f. 3-5 text-f. 39-41; spp. nov. K. spathi orientalis p. 55 pl. 2 f. 2-7 text-f. 28, 29 var. nov. Cretaceous. Moghara Massif, Sinai, Egypt; K. aegyptiacum nom. nov. p. 57 for Kn. "Uhligi" ap. H. Douvillé, 1916, I. G. E.-D. Mahmoud.

†Kosmoceras (Galielmites) media pp. 222, 226 pl. 2 f. 1-4, pl. 3 f. 3, 4; K. (G.) nodosum pp. 222, 229 spp. nov., J. H. Callomon (1) Philos Trans. 239B 664 1955.

Kossmaticeratidae in Patagonia, Argentina, G. Cecioni (1).

†Laevaptychus species from the Jurassie of southern Roumania, Răileanu, Bădălută & Pelin.

†Laevilamellaptychus crassissimus from the Italian Jurassic, G. Cuzzi.

†Lamedaptychus, Jurassic species from southern Roumania, Răileanu, Bădălută & Pelin.

†Lamellaptychus species from the Jurassic of Italy, G. Cuzzi.

†Leopoldia? paynensis from the Cretaceous of Patagonia, G. Cecioni.

†Lepthoplites cantabrigiensis rarecostate [error for rarecostata] p. 47 pl. 5 f. 3, 4; L. pseudoplanus subdiscoidea p. 50 pl. 5 f. 9 vars. nov. Kopet-Dagh, Trans-Caspia, Russia. Cretaceous, Albian, A. Ghlazunova.

†Leptoceras (?Lytocrioceras) sabaudianum non-turberculata var. nov. Lower Cretaceous, south east France, S. Sarkar p. 140 pl. 9 f. 13; pl. 10 f. 14.

†Leptosphinctes (Praeleptosphinctes) subgen. nov. subgenotype: L. jaworskii sp. nov. = "Morphoceras nov. sp. indet. cf. defrancei d'Orb." in Jaworskii, 1926, G. Westermann p. 268.

†Levisoceras magnum sp. nov. Ordovician Central Texas, Unklesbay (1) J. Paleont. 28 5 1954 pp. 637, 651 pl. 70 f. 14, 15.

†Ludwigia eschense sp. nov. p. 389 pl. 16 Jurassic. Luxembourg, P. Maubeuge (3).

Lycetticeras Gen. nov. genotype L. lycetti sp. nov. Middle Bathonian, Great Oolite, Minchhampton, W. J. Arkell (1) Geol. Mag. 90 1 1955 p. 37 pl. 1 f. 2.

Lytoceras Suess 1865, neuter gender, Direction 42.

†Macrocephalitacea superfam. nov. Ammonoidea. Macrocephalitidae, Buck., Cardioceratidae, H. Douvillée and Pachyceratidae, Buck., N. T. Sazonov p. 117.

†Medlicottia vetusta p. 151 pl. 12 f. 1-7; M. intermedia p. 153 pl. 13 f. 4-7 spp. nov. Rivers, Aktasta and Sakmara, Russia. Kurmainskii horizon, Permian V. Ruzhentzev (3).

†Megapronorites gen. nov. p. 60 Pronoritidae; M. sakmarensis sp. nov. p. 61 pl. 1 f. 1-9. Namurian, Carboniferous. S. Urals, Russia, V. Ruzhentzev (3).

Megatyloceras Humphrey, 1949, to be placed on the "Official List of Generic Names in Zoology," Opinion 428.

†Metapronorites cuncilobatus sp. nov. p. 69 pl. 11 f. 5-8. Upper Carboniferous. Sakmara, Russia, V. Ruzhentzev (3).

†Mexicoceras gen. nov. Cyclolobidae Type: Waagenoceras cumminsi var. guadalupense Girty, 1908. Delavare Mountain, Texas, V. E. Ruzhentzev (1) p. 701 f. la.

†Mikasaites gen. nov. p. 174 Kossmaticeratidae, sub.-fam. Marshallitinae. genotype: M. orbicularis sp. nov. p. 175 pl. 16 f. la-d, 2a-c. Cretaceous, Hokkaido, Japan, T. Matsumoto (1).

†Mimosphinetinae subfam. nov. Type Mimosphinates Lichenberg 1931 genotype M. tripartitons Eich., H. K. Erben Neu. Jb. Geol. Palaont. 93 2 1953 p. 191 pl. 18 f. 3-7.

†Morrisceratinae subfam. nov. of Macrocephalitidae, to include genera *Morrisiceras* and *Lycetticeras*. Jurassic G. Westermann p. 262.

Muensteroceras emendation of the generic name Munsteroceras Hyatt, 1884, Opinion 391.

†Neoaganides tabantalensis sp. nov. Sokmara deposit in the Aktyubinsk Province of the Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 58 pl. 1 f. 7.

†Neodimorphoceras (?) armstrongi sp. nov. Carboniferous. Scotland, E. D. Currie p. 593 pl. 3 f. 13 text-f. 2 m.

†Neomorphoceras gen. nov. genotype Ammonites chapnisi Oppel pl. 4 Upper Oxfordian, W. J. Arkell (1) Geol. Mag. 90 1 1953 p. 38 pl. 1 f. 4.

†Neopharciceras kurbatovi gen. nov. sp. nov. Pharciceratidae fam. nov. Rudnii Altag Devonian, Bogoslovski, B. I. p. 95.

†Neopronorites prior p. 78 pl. 3 f. 8-10; N. carboniferus p. 80 pl. 4 f. 1-8; spp. nov. Upper Carboniferous. Sakmara, Russia, V. Ruzhentzev (3).

†Neposiella gen. nov. Ammonitacea N. nepos H. Douvillé; N. spathi sp. nov. p. 101 pl. 5 f. 23, 24 text-f. 47, 48. Cretaceous, Moghara Massif, Sinai, Egypt, I. G. Mahmoud.

†Nicklesia nodosa p. 50 pl. 9 f. 4, 5; Pl. 10 f. 1; pl. 3 f. 7 sp. nov. Chipatá, Santander; N. dumasiana retrocurvata p. 50 pl. 9 f. 2, 3 Boyacá; N. didayana multifida p. 54 pl. 11 f. 4, 5 Bolivar subspp. nov. Barremian, Cretaceous. Colombia, H. Bürgl (2).

†Nicklesiella subgen. nov. p. 55 of Pulchellia subgenotype. P. (N.) leivaensis sp. nov. p. 56 pl. 11 f. 10, 11. Cretaceous Colombia, H. Bürgl (2).

†Nordiceras gen. nov. p. 1104 Pharciceratidae; genotype: Prolecanites timanicus Holzapfel 1899. Lower Franconian. Timan, Kolima, Russia, B. I. Bogoslovski (1).

†Obtusicostites rajnathi p. 37 pl. 4 f. 2a, b. O.(?) spathi p. 38 pl. 5 f. 1a, b spp. nov. Callovian Kutch, India, Agrawal (1) Ann. Cent. ét Doc. Paléont. No. 19 1956.

†Oecoptychiinae subtam. nov. of Morphoceratidae. Jurassic, G. Westermann p. 273.

†Onitshoceras gen. nov. p. 248 Desmoceratidae: genotype species: O. matsumotoi sp. nov. p. 249 pl. 3 f. 1, 2; pl. 5 f. 7. Upper Cretaceous. Onitsha Province, Nigeria, R. A. Reyment.

†Oosterellidae, from the North African Cretaceous, Basse & Delga.

†Orthoceras Brünnich 1771 emend. nov. genotype: Orthoceratites regularis Schlotheim, 1820, Z. Balashov p. 233.

†Pachyvascoceras Furon 1935, previously a sub-genus of Vascoceras Choffat, now given generic rank, p. 257. Type species (here designated): Vascoceras (Pachyvascoceras) crassum

Furon 1935; P. costatum p. 257 pl. 3 f. 6 pl. 4 f. 3 pl. 5 f. 2; P. proprium p. 258 pl. 5 f. 1; P. globosum p. 259 pl. 3 f. 3 pl. 5 f. 4; spp. nov.; P. proprium plenum subsp. nov. p. 258 pl. 5 f. 5. Upper Cretaceous. Nigeria, R. A. Reyment.

†Paracenoceras sornayi sp. nov. Callow-Oxfordian, Kutch, India, Agrawal (1) Ann. Cent. Ét. Doc. Paléont. No. 19 1956 p. 14 pl. 1 f. 4.

†Podagrosiceratidae fam. nov. p. 621 Podagrosiceras gen. nov. p. 621, genotype: P. athleticum sp. nov. p. 622 pl. 1 f. 1. Aalenian, Jurassic. Neuguén region, Argentina, Maubeuge & Lambert.

†Paragastrioceras strigosum sp. nov. Sakmara deposit. Aktyubinsk Province. Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 72 pl. 5 f. 1-3.

†Parkinsonitacea superfam. nov. Ammonoidea. Marphoceratidae Buck., Cosmoceratidae, Hyatt, Proplanulitidae Buck., and Aulacostephanidae fam. nov., Sazonov, N. T. p. 117.

†Phaedrysmocheilus gen. nov. genotype: Nautilus subaratus Keyserling, 1860. Pleuronautilidae, Shimanski, V. N. & Erlanger, A. A. Bull. Soc. Nat. Moscou Geol. 30 3 1955 p. 96.

†Pharciceratidae fam. nov. Ammonoidea. *Pharciceras*, Hyatt, *Synpharciceras*, Schindewolf, and *Neopharcicras* gen nov. Rudnoi Altaya, Russia. Devonian p. 95, B. I. Bogoslovski.

†Phyllopachyceras katschiense p. 121 pl. 11 f. 42; P. segne p. 127 pl. 12 f. 43; P. ectocostatum p. 128 pl. 13 f. 52-54 P. crassum p. 130 pl. 13 f. 51 spp. nov. Lower Cretaceous. Crimea and Northern Caucasus, V. Drushchitz.

†Pictonia Bayle, 1878: P. laylei Salfeld, 1913 designated as type. The generic and specific names validated, Opinion 426.

†Placenticeras benningi sp. nov. Cretaceous, Alabama, U.S.A., L. Stephenson (1) p. 247 pl. 44 f. 21, 22; pl. 45 f. 7-11.

†Platyclymenia (Platyclymenia) grossi p. 71 pl. 2 f. 21; P. (P.) quiringi p. 74 pl. 2 f. 25 spp. nov.; P. (P.) subnautilina schleizi p. 74 pl. 2 f. 24 subsp. nov. Devonian, Thuringia, Germany, K. J. Müller. †Pleydellia buckmani sp. nov. (non Harpoceras opalinum Rein, 1942) Aalenian, Jurassic. Luxembourg. P. Maubeuge (2) p. 76 pl. 2 (see also, Arch. Inst. Gr. Duc. Luxemb. N.S. 18 1949 pl. xvii); P. pseudoaalense p. 378 pl. 9; P. falcifer p. 379 pl. 9; P. funcki p. 379 pl. 10; P. arkelli p. 380 pl. 10; P. (Walkericeras) pseudoarcuata p. 382 pl. 12; P. (W.) pseudolotharingica p. 383 pl. 13; P. (W.) dudelangense p. 385 pl. 14; spp. nov. Jurassic. Luxembourg, P. Maubeuge (3) P. argentina sp. nov. Aalenian, Jurassic. Neuguén region, Argentina, Maubeuge & Lambert p. 622, pl. 1 f. 2-6.

† Poecilomorphus lauxi sp. nov. Jurassic. Luxembourg, P. Maubeuge (1) p. 163 pl. 3.

†Praeleptosphinctes subgen. nov. of Leptosphinctes subgenotype: L. jaworskii sp. nov. = "Morphoceras nov. sp. indet. cf. defrancei d'Orb." in Jaworskii, 1926, G. Westermann p. 268.

†Praetulites subgen. nov. of Chondroceras (?) subgenotype C. kruizingai sp. nov. = "Sphaeroceras gadohense" (Kruizinga, 1926, non Bochm), G. Westermann p. 257.

†Préperonoceras "nov. gen." [subgen. nov.] of Coeloceras [q.v.], P. Maubeuge p. 136.

†Procerozigzag gen. nov. type Stephanoceras crassizigzag Buckman 1892 vara., W. J. Arkell (1) Geol. Mag. 90 1 1953, p. 37.

†Progeronia gen. nov. type Perisphinctes progeron v.Ammon 1875 p. 181 pl. i, f. 2 Lower Oxfordian, S. France, W. J. Arkell (1) Geol. Mag. 90 1 1953 p. 38.

†Prolecanitina suborder nov. of Thurmonoidea, Miller and Furnish J. Paleont. 28 5 1954 p. 687.

† Protacrioceras subgen. nov. of Acrioceras Ancyloceratidae. Lower Cretaceous, south east France, S. Sarkar p. 129.

†Protetragonites mediocris p. 97 pl. 6 f. 21; P. karakaschi p. 99 pl. 6 f. 22, 23 spp. nov. Lower Cretaceous. Crimea and North Caucasus, V. Drushchitz.

†Prothalassoceras serratum sp. nov. p. 23 pl. 3 f. 1, 2; P. biforme latilobata var. nov. p. 25 pl. 3 f. 3, 4. Yurezan river, S. Urals, Russia. Schwagerina horizon, Permian, S. Maksimora. P. umbilicatum sp. nov. Sakmora deposit, Aktyubinsk Province, Russia. Permian?, V. Ruzhentzev (4) p. 59 pl. 1 f. 8, 9; pl. 6 f. 1.

†Protogrammoceras a. bassanii Domerian Jurassic, Yorkshire, First record from Britain, M. Howarth p. 169.

†Pseudaganidinae subfam. nov. Nautilidae. To include genera *Pseudaganides* Spath and *Pseudonautilus* Meek, B. Kummel (1) p. 389.

†Pseudaspidoceras paganum sp. nov. Upper Cretaceous. Bauchi Province, Nigeria, R. A. Rayment p. 253 pl. 4 f. 1 text-f. 3h, 4.

†Pseudolillia murvillensis 'gen. et sp.' nov. Jurassic. Murville, France, P. Maubeuge (1) p. 150 pl. 1 pl. 2 f. 1.

Pseudonautilidae fam. nov. Nautiloidea, Shimanski, V. N. & Erlanger, A. A. Bull. Soc. Nat. Moscou geol. 30 3 1955 p. 96.

†Pseudotetragonites gen. nov. Lytoceratidae: genotype: P. kudrjavzevi sp. nov. Lower Cretaceous. Crimea and North Caucasus, V. Drushchitz p. 80 pl. 7 f. 25.

†Pseudothurmannia angulicostata tuberculata p. 155 pl. 6 f. 13; P. mortilleti rotunda p. 156 pl. 6 f. 12 vars. nov.; P. simionescui sp. nov. p. 157 pl. 11 f. 1 Lower Cretaceous, South East France; P. picteti nom. nov. for specimen figured by Pictet under the name Amm. angulicostatus D'Orb.; p. 156, S. Sarkar.

†Pseudowalkericeras hussignyensis gen. et sp. nov. Jurassic. Hussigny, France, P. Maubeuge (1) p. 151 pl. 2 f. 2.

†Psiloceras from the Mesozoic of central Switzerland, R. Trümpy.

†Psilotissotia chalmasi andina subsp. nov. p. 85 pl. 26 f. 5 Santander; P. maxima sp. nov. p. 86 pl. 26 f. 1, 2 Boyacá. Cretaceous, Colombia, H. Bürgl (2).

†Ptychophylloceras varshneyi sp. nov. Callovian Kutch, India, Agrawal, L. (1) Ann. Cent. Ét. Doc. Paléont. No. 19 p. 16 pl. 1 f. 3a, b.

†Pulchellia (Nicklesiella) subgen. nov. p. 55 subgenotype: P. (N.) leivaensis p. 56 pl. 11 f. 10, 11; pl. 2 f. 5 Boyaca; P. (N.) pseudokarsteni p. 57 pl.

12 f. 1-4; pl. 3 f. 12, 13. Santander spp. nov.; P. (Semipulchellia) subgen. nov. p. 57 subgenotype: P. (S.) communis p. 59 pl. 12 f. 6-12; P. (S.) elegans p. 58 pl. 12 f. 5; P. (S.) robusta p. 60 pl. 13 f. 1, 2 spp. nov. Santander; P. (Pulchellia) riedeli p. 64 pl. 13 f. 9, 10 Cundinamarca; P. (P.) radians p. 65 pl. 13 f. 14; pl. 14 f. 1, 2. Santander spp. nov.; P. (P.) galeata ornata p. 63 pl. 13 f. 7, 8 subsp. nov.; P. (Caicedia) subgen. nov. p. 66 subgenotype: Ammonites caicedi Karsten; P. (C.) royoi p. 68 pl. 15 f. 1, 2; pl. 4 f. 1 sp. nov.; P. (Hettneria) subgen. nov. p. 70 subgenotype: P. selecta Gerhardt; P. (H.) orbignyi p. 73 pl. 16 f. 4. Santander sp. nov. Barremian, Cretaceous. Colombia, H. Bürgl (2); P. (Heinzia) ouachensis from the lower Cretaceous of N.-W. Caucasus. First record for the area, N. Luppov; P. and other Barremian genera from Trinidad, R. W. Imlay.

†Punctaptychus punctatus from the Italian Jurassic, G. Cuzzi; P. species from the Jurassic of southern Roumania Răileanu, Bădălută & Pelin.

†Rasenia Salfeld, 1913: R. involuta (Salfeld MS.) Spath, 1935 designated as type. Generic and specific names validated, Opinion 426.

†Reinecheia ravana jhuraensis var. nov. Collovian Kutch, India, Agrawal (1) Ann. Cent. Et. Doc. Paléont. No. 19 1956 p. 31 pl. 1 f. 2a, b; R. (Reineckeites) alemannica sp. nov. p. 85 pl. 3 f. 12; R. (R.) britannica n. nom. for Kellawaysites multicostatus Peticlere sp. Buckman; p. 86; R. (R.) corroyi p. 88 nom. nov. for R. stuebeli Steinmann, Corroy; R. (R.) aff. jeanneti nom. nov., p. 89, for R. hungaricus Till, Jeannet, A. Zeiss.

†Renziceras gen. nov. type Hildoceras nausikaae Renz. Toarcian, Epiraes Greece, A. J. Arkell (1) Geol. Mag. 90 1 1953 p. 36 pl. f. 3.

†Sagittoceras costulatum from Sauerland, Germany, A. Kraemer.

†Salfeldiella (Aphroditiceras) subgen. nov. p. 80; S. (A.) aphrodite venus var. nov. p. 80 pl. 5 f. 1 text-f. 43; S. (Goretophylloceras) moreti sp. nov. p. 81 pl. 5 f. 2-4 text-f. 44, 45. Cretaceous. Moghara Massif, Sinai, Egypt, Mahmoud, I. G. E.-D. Sandbergeroceras emendation of the generic name Sandbergeoceras Hyatt, 1884, Opinion 392.

†Sphaeropharciceras gen. nov. Pharciceratidae. Type S. sandbergerorum sp. nov. (= Goniatites tridens Sandberger (pars)) Lower Franconian, Triassic? West Germany, B. I. Bogoslovski (1) p. 1104.

†Schmidtoceras subgen. nov. of Chondroceras Jurassic. Europe, G. Westermann (1) p. 66.

†Semipulchellia subgen. nov. p. 57, of Pulchellia Type: P. (S.) communis sp. nov. p. 59 pl. 12 f. 6-12. Barremian, Cretaceous. Santander, Colombia, H. Bürgi (2).

†Solgerites tuberculatus sp. nov. Upper Cretaceous. Numan Division, Nigeria. R. A. Reyment p. 267 pl. 2 text-f. 11 a, b.

†Sphaeroceras brongniarti terpartitum subsp. nov. Jurassic, Calvados, France, G. Westermann (1) p. 33 pl. 14 f. 3-5.

†Spathicrioceras gen. nov. Heteroceratidae Genotype: S. sornayi sp. nov. p. 160 pl. 8 f. 8. Lower Cretaceous, South East France, S. Sarkar.

†Spathites gen. nov. Vas-coceratidae for chispaensis sp. nov. Lower Turonian, Mexico, Kummel & Dicken J. Paleont. 28 3 1954 pp. 310, 311 pl. 30 f. 1, 2 pl. 31 f. 1-15 text-fig. 1.

†Spectoniceras versicolor from the Russian Cretaceous. Note on structure, R. Reyment (1).

†Spirogmoceras gen. nov. of Trachyceratidae genotype Protracheeras shastense Smith, N. J. Silberling, J. Pal. 30 5 1956 p. 1152.

†Stephanoceras cf. plicatissimum middle Jurassic, Miyagi Prefecture, Japan, T. Sato (1) p. 169 f. 2–5 pl. 13.

Strenoceras and Ammonites of the Dogger—sutures, figs., O. H. Schindewolf.

†Subtissotia peroni Hyatt 1903 = Tissotia africana Peron, 1897, H. Cruys.

†Synuraloceras gen. nov. Paragastrioceratidae p. 73 genotype: S. carinatum sp. nov. p. 75 pl. 5 f. 4-6 Sakmara deposit, Aktyubinsk Province. Kazakh, Russia. Permian?, V. Ruzhentzev (4). †Tabantalites gen. nov. p. 76 Marathonitidae: genotype: T. bifurcatus sp. nov. p. 77 pl. 6 f. 3-7 Sakmara deposit. Aktyubinsk Province. Kazakh, Russia Permian?, V. Ruzhentzev (4).

†Tetragonites crudus sp. nov. Lower Cretaceous. Crimea & Northern Caucasus, V. Drushchitz p. 105 pl. 8 f. 29.

†Thalassoceras ulkense sp. nov. Sakmara deposit, Aktyubinsk Province, Kazakh, Russia. Permian?, V. Ruzhentzev (4) p. 61 pl. 6 f. 2.

†Timorites sigillarius sp. nov. Khingan, ridge Great Churki, Russia upper Permian, Ruzhentzev, V. E. (1) C.R. Acad. Sci. Moscou 103 4 1955 p. 701.

†Tithopeltoceras gen. nov. type Aspidoceras moricanii Meneghini 1885 pl. xxii f. 2, W. J. Arkell (1) Geol. Mag. 90 l 1953 p. 39 pl. 1 f. 1.

†Tornoceras (Aulatornoceras) keyserlingi nom. nov. p. 49, for Goniatites cinctus Keyserling 1844, non Braun 1842; T. acutiforme Gürich = Dimeroceras lentiforme Sandberger (Jugendform) p. 55, 80, K. J. Müller.

†"Toxoceras" requienianum unituberculata var. nov. Cretaceous, south east France, S. Sarkar p. 56 pl. 5 f. 1a, b.

†Uddenites postsakmarensis p. 108 text-f. 53. Ural River; U. orenburgensis p. 109 text-f. 54. Orenburg; U. tuberculatus p. 111 pl. 5 f. 15, 16. Ural River; U. convexus p. 112 text-f. 56. Sakmara spp. nov. Carboniferous. Russia, V. Ruzhentzev (3).

†*Uddenoceras* gen. nov. for *Uddenites* oweni Miller and Furnish 1940, Miller & Furnish J. Paleont. 28 5 1954 p. 691.

†Uraloceras serpentinum sp. nov. Schwagerina horizon, Permian. Yurezan river, S. Urals, Russia, S. Maksimora p. 7 pl. 1 f. 1-5.

Utaturiceras gen. nov. Acanthoceratidae, subfam. Metoicoceratinae: genotype species: Ammonites vicinalis Stoliczka, 1864. Upper Cenomanian (Cretaceous), C. W. Wright p. 392.

†Valdedorsella gignouxi sp. nov. Cretaceous. Moghara Massif, Sinai, Egypt, I. G. Mahmoud p. 94 pl. 5 f. 15.

†Virgataxioceras gen. nov. type Virgatosphinates setatus Schneid 1914 p. 160 pl. v f. 5 pl. vi f. 4. Middle Kimeridginae, Bavaria, W. J. Arkell (1) Geol. Mag. 90 1 1953 p. 39.

†Virgatitacea superfam. nov. Ammonoidea. Virgatitidae Spath, Virgatosphinctidae Spath and Dorsoplanitidae Arkell, N. T. Sazonov p. 117.

†Walcottoceras helleri sp. nov. Ordovician Texas, Unklesbay (1) J. Paleont. 28 5 1954 pp. 637, 652 pl. 71 f. 5-7.

†Witchellia tibetica sp. nov. Bajocian Tibet, W. J. Arkell (2) Geol. Mag. 90 5 1953 p. 332 pl. xiv f. 7a, b.

†Yebisites gen. nov. p. 205 Psiloceratidae. Genotype: Y. onoderai sp. nov. p. 207 pl. 30, Shizukawa Group, Jurassic. Miyagi Prefecture, Japan, T. Matsumoto.

† Yokoyamaoceras ornatum p. 183 pl. 16 f. 3a, b, 4a-d, Abeshinai Valley; Y. minimum p. 185 pl. 16 f. 6a-c, Ikushumbets Valley spp. nov. Cretaceous, Hokkaido, Japan, T. Matsumoto (1). †Zieteniceras subgen. nov. of Hecticoceras, subgenotype: H. zieteni de Tsytovitch, 1911. Callovian Jurassic, Blumberg, Germany, A. Zeiss p. 24.

## INSERTAE SEDIS

Agasoma Gabb "[1869]", neuter gender, Direction 42.

†Cowwarrella gen. nov. Genotype: C. cylindrica sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 61 pl. 6 f. 12-14.

†Ostlerina gen. nov. Genotype: O. delicata sp. nov. Siluro-Devonian. Marble Creek, Thomson river, Victoria, Australia, Talent & Philip p. 59 pl. 6 f. 1-4 text-f. 2, 3.

†Stenothecoides, on the shell structure, F. Rosetti.

